

Contents

RECEIVED  
JUN 5 1972  
449-417



50 POUNDS

# CHLORDANE 10% GRANULAR

ACTIVE INGREDIENT:  
\*Technical Chlordane

10%

INERT INGREDIENTS

90%

100%

\*Equivalent to 6% Octachloro-4,7-Methanotetrahydroindane  
and 4% Related Compounds.

FOR CRABGRASS CONTROL AND INSECT CONTROL OF CERTAIN  
SOIL-INHABITING INSECTS

## DIRECTIONS FOR CRABGRASS CONTROL

Apply 15 lbs. Chlordane 10% Granular per 1000 sq. ft. of lawn or ornamental turf to suppress crabgrass seedlings before germination or sprouting occurs. It is for use as an early season preventive of established crabgrass plants. Timing of application is important and should be applied before any germination which in many areas is during the period April 15 to May 1. This treatment will not burn or injure established lawns. Do not apply to newly seeded lawns.

HELPFUL HINTS: After crabgrass seedlings have been suppressed by the Chlordane treatment, control may be assisted by:

1. Mowing the lawn to 1 1/2 to 2 inches.
2. Using an adequate quantity of fertilizer to maintain active growth.

After treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children or pets.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

## RECOMMENDATIONS

INSECT CONTROL ON ORNAMENTAL TURF, NURSERY AND GREENHOUSE SOILS: For the control of Ants, Armyworms, Chiggers, Chinch Bugs, Crickets, European Earwigs, Grasshoppers, Japanese Beetle Grubs, Rose Chafer, Sod Webworm, Spiders, Ticks, White-fringed Beetle, White Grubs, and Wireworms in Turf (such as Golf Courses, Athletic Fields, Large Estate Nurseries, Cemeteries and Park Sod), and in Nursery and Greenhouse soils where ornamental plants are grown, use at the rate of 50 lbs. of Chlordane 10% Granular per acre (1 1/4 lbs. to each 1000 sq. ft.) to cover surface uniformly, using a lawn fertilizer spreader. Treatment should be applied in the spring or when insects are observed to be troublesome. Do not treat soil when it is wet. Apply before rain or sprinkling. Do not apply to newly seeded lawns.

INSECT CONTROL ON HOME LAWNS: For the control of the above listed insects on home lawns apply 50 lbs. of Chlordane 10% Granular per acre (1 1/4 lbs. to each 1000 sq. ft.) to cover surface uniformly using a lawn fertilizer spreader. Treatment should be applied in the spring, or when insects are observed to be troublesome. Immediately after treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children and pets. Do not treat when soil is wet. Apply before rain or sprinkling. CAUTION: Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass into the soil by drenching or sprinkling with water and grass has completely dried.

CAUTION: Hazardous if swallowed, inhaled, or absorbed through skin! In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, flush with plenty of water for at least 15 minutes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. Keep away from children and domestic animals. Do not store near feed and foodstuffs. Do not apply to areas occupied by unprotected humans or beneficial animals.

This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label. Birds feeding on treated areas may be killed. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse empty container. Destroy by burying with waste or burning. Bury or discard in a safe place. Stay away from smoke or fumes.

NOTICE: Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured For

# WOODBURY CHEMICAL CO.

Division of TECHNE CORP.

ST. JOSEPH, MO. 64502; DENVER, COLO.

U.S. PAT. 2,811,100  
REG. U.S. PAT. OFF.