

LBS.


NET

ZINEB 75-W

A WETTABLE POWDER FOR USE IN WATER

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT:	
ZINEB (Zinc ethylene bisdithiocarbamate)	75%
(Zinc expressed as metallic . . . 17.7%)	
INERT INGREDIENTS	25%
TOTAL	100%



CAUTION

May cause irritation of nose, throat and skin. Do not breathe dust or spray mist. In case of prolonged or frequent use, wear clean protective clothing and bathe after each exposure. Keep containers closed. Store in dry place. Keep away from fire and sparks.

CAUTION

KEEP OUT OF REACH OF CHILDREN
DIRECTIONS FOR USE

Pour recommended amount of this material through screen on surface of water in nearly filled spray tank or pre-mix in a bucket half filled with water and pour mixture into nearly filled spray tank. Add balance of water to fill tank. Do not allow mixture to stand. Keep agitator running during filling, mixing and spraying operations. Do not use in low volume gear type spray equipment. This material is compatible with most of the commonly used insecticides, fungicides and nutritional sprays. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible. applications needed will vary with local conditions.

Consult State Agricultural Experiment Stations for specific information as the timing, number and rate of

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and do not apply closer to harvest than specified.

RECOMMENDATIONS

The dosage rates on this label indicate pounds of Zineb 75-W per 100 gal. of dilute spray.

FRUIT AND NUTS

APPLE: Apple Blotch, Black Rot, Bitter Rot, Botryosphaeria Fruit Rot, Brooks Spot, Cedar Rust, Fly Spock, Frog-eye, Quince Rust, Scab, Sooty Blotch—Use 1 to 2 lb. Begin applications in delayed dormant and continue through cover sprays. Use ½ to 1 lb. when combined with Captan. Do not use within 7 days of harvest.

APPLE, PEAR: Fire Blight—Use 2 lb. at 2 to 3 day intervals during periods of bloom-opening and wet weather. Maintain recommended sanitary pruning practices. Do not apply within 7 days of harvest.

APRICOT, PEACH: Coryneum Blight (Shot Hole), Peach Leaf Curl—Use 2 lb. in post-harvest or dormant sprays and in popcorn to jacket stages according to local spray program.

CITRUS: Rust Mite and associated Russetting and Greasy Spot—Use 1 lb. Apply with thorough coverage. Make application from late June through early August, or earlier if rust mite begins to show.

SOUR CHERRY: Leaf Spot or Shot Hole—Use 1 lb. Apply at petal fall and 10 and 20 days after petal fall, and immediately after harvest. Do not apply within 7 days of harvest.

CURRANT: Leaf Spot—Use 2 lb. Apply 2 to 3 weeks after bloom and again immediately after harvest.

GRAPE: Black Rot, Downy Mildew, Ripe Rot—Use 1½ lb. just before bloom, just after bloom, 10 days after bloom, and 2 to 3 weeks later. Do not apply within 7 days of harvest.

STRAWBERRY: Leaf Spot—Use 1½ to 2 lb. Begin applications when new growth starts in spring and repeat at 7 to 10 day intervals until fruit begins to form. Leaf Scorch—Use 1½ to 2 lbs. in new beds in late summer. Make 2 to 3 applications at 7 day intervals. Do not apply within 7 days of harvest.

VEGETABLES

Use 1½ to 2 lb. per 100 gal. of dilute spray. Begin applications at first signs of disease and repeat at 7 to 10 day intervals or as needed to maintain control of the diseases stated.

BEANS: Anthracnose—Begin when first blossoms appear. Downy Mildew, Rust—Begin when disease threatens. Do not apply within 7 days of picking.

BEETS, KALE, MUSTARD GREENS, SWISS CHARD, TURNIP: Cercospora Leaf Spot, Downy Mildew—Do not apply within 7 days of harvest.

CARROT: Alternaria, Cercospora, Leaf Spot—Do not apply within 7 days of harvest if tops are to be used as food or feed.

CELERY: Early Blight, Late Blight—Begin applications when plants emerge in the plant beds. Repeat at 3 to 5 day intervals until plants are set in field. Continue applications at 7 day intervals. Remove residues remaining at harvest by washing and stripping.


CUCURBIT'S (Cantaloupe, Cucumber, Melons, Squash): Alternaria (Macrosporium) Blight, Anthracnose, Downy Mildew, Gummy Stem Blight—Direct spray to underside of leaves. Do not apply within 7 days of harvest on summer squash.

LETTUCE: Downy Mildew—Do not apply within 5 days of harvest on head lettuce or 7 days on leaf lettuce.

ONION: Botrytis Leaf Blight (Blast), Downy Mildew, Purple Blotch—Do not apply within 7 days of harvest on green or spring onions.

SPINACH: Downy Mildew (Blue Mold), White Rust—Do not apply within 7 days of harvest.

TOMATO: Anthracnose, Cladosporium Gray Mold, Early Blight, Late Blight, Septoria Leaf Spot, Stemphylium Leaf Spot.



CONDITIONS OF SALE AND WARRANTY: The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application even though the product is used in accordance with Directions. All such risks shall be assumed by the Buyer.

Woodbury Chemical Co. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. Woodbury makes no express or implied warranty of fitness or merchantability or any other express or implied warranty. In no case shall Woodbury or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Woodbury and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of Woodbury Chemical Company.

Manufactured by
WOODBURY CHEMICAL COMPANY

St. Joseph, Mo.

Denver, Colo.

Orlando, Fla.

Lubbock, Texas

USDA Reg. No. 449- 431