

EACH QUART IS EQUIVALENT TO ONE POUND 50% WETTABLE METHOXYCHLOR POWDER

#### CHERRIES

For Cherry Fruit Fly, use 1 quart in 100 gallons of water. Make first application when flies appear and repeat applications at 7 to 14 day intervals until harvest. Do not apply within 7 days of harvest.

#### ALFALFA

For spittlebug, apply 2 to 4 quarts per acre in sufficient water to wet thoroughly the crowns of the plants. Make application in the spring when plants are 4 to 6 inches high, or in the fall during the period between the end of spittlebug migration into new fields and the beginning of egg-laying. For fall Armyworm, Lygus Bug nymphs, Alfalfa Weevil, Alfalfa Caterpillar and certain other insects, apply 4 to 5 quarts per acre. For Leafhoppers, apply 1 to 2 quarts per acre. Apply in sufficient water to give adequate coverage, when insect populations begin to build up. Do not apply within 7 days of harvest or grazing.

#### EMPTY GRAIN BINS

As an aid for control of Rice Weevil, Grain Weevil, Lesser Grain Borer, Angoumois Grain Moth and certain other insect pests of stored grain, during accessible or exposed stages, and as a treatment to help prevent infestation of newly stored grain, apply 1 gallon with 9 gallons of water. Make application to thoroughly cleaned surfaces at the rate of 2 gallons of spray mixture per 1,000 sq. ft. Allow sprayed surfaces to dry before storing grain.



SURE DEATH BRAND

# METHOXYCHLOR

## Emulsifiable Concentrate

Contains 2 Lbs. of Methoxychlor per Gallon

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

#### ACTIVE INGREDIENTS

Methoxychlor, Technical	24%
Methylated Naphthalenes	63%
Dimethyl Phthalate	10%
INERT INGREDIENTS	3%

U. S. Pat. 2,420,328

\*24% Methoxychlor, technical is equivalent to 21% 2,2-bis (p-methoxyphenyl) 1,1,1-trichloroethane and 3% other isomers and reaction products.

FOR USE AGAINST INSECTS AFFECTING DAIRY CATTLE, LIVESTOCK, BARN, WAREHOUSES

FOR SPRAYING BEEF CATTLE for horn flies, stableflies. Only supplementary control can be obtained — major control would be in the thorough spraying of barns and treatment of other breeding area and lice. Use 1 quart per 12 gallons water, using 2 quarts of the dilute solution per adult animal. Repeat as necessary 2-3 weeks.

#### FOR SPRAYING INTERIOR OF BARN

For House Flies, Stable Flies and Mosquitoes: use 1/2 gallon per 5 gallons of water. Spray thoroughly walls, ceilings, stalls, screens, manure piles, etc., using 1 gallon dilute spray to each 600-1,000 sq. ft. Repeat as necessary 3-4 weeks.

COVER TROUGHS, BINS OR OTHER FOOD OR WATER RECEPTACLES WHILE SPRAYING. Exclude dairy animals during treatment.

#### COMPATIBILITY

Do not use with materials containing sulfur or other substances incompatible with oil. Do not use on crops within 14 days of application of any material containing sulfur.

#### CAUTION

MAY CAUSE SKIN IRRITATION

Harmful if swallowed

Avoid contact with skin

In case of contact, wash with soap and water.

Avoid contamination of feed and food stuffs.


Do not use on humans or animals.

Do not store or use near fire or open flame.

This product is toxic to fish. Keep out of any body of water.

NOTICE: This product is a restricted use pesticide. It is to be used only by or under the direct supervision of a person licensed by the state in which it is used.

U. S. D. A. Reg. No. 449,451



Manufactured by  
WOODBURY CHEMICAL COMPANY  
St. Joseph, Mo.      Denver, Colo.      Orlando, Fla.      Lubbock, Tex.