

CONTENTS

POUNDS

Sure Death Brand

COPPER - SEVIN® - SULFUR 3.4 - 5 - 75 DUST

AN AGRICULTURAL INSECTICIDE - FUNGICIDE

ACTIVE INGREDIENTS:

Copper (in basic copper sulfate) (expressed as metallic)	3.4%
Carbaryl (1-Naphthyl N-methylcarbamate)*	5.0%
Sulfur	75.0%

INERT INGREDIENTS:

TOTAL

16.6%
<u>100.0%</u>

*SEVIN ® Trademark-Union Carbide Corporation

Sulfur fineness: 93% Passing No. 325 USS Sieve.

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

DO NOT BREATHE DUST
DO NOT CONTAMINATE FEED OR FOODSUFFES
AVOID SKIN CONTACT
WASH WITH SOAP AND WATER AFTER USING
DANGEROUS IF SWALLOWED

DESTROY EMPTY CONTAINERS--
BURY OR BURN. FLEE OUT OF
SMOKE.

OBSERVE THE FOLLOWING USE PRECAUTIONS:

Do not apply when weather conditions favor drift of dust from crop areas treated. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals, or onto adjacent food or forage crops about to be harvested. Contamination of productive waters may destroy essential fish foods. Do not apply when crops are in bloom. Remove or cover bees if dusting is necessary. Apply in early morning or evening when there is little or no wind. Do not use fruit and leaf injury to sulfur-sensitive crops. Do not apply to artichokes, comice pears, cranberries, cucumbers (for winter), eggplant, melons, squash, walnuts, or to other sensitive plants. Do not use on foliage and fruit during periods of high temperature and high humidity or matic conditions. Do not apply sulfur to sensitive crops when shade temperatures are exceeding or are likely to exceed 85°F. Do not use on citrus and other unless local use has proved safe. Do not use on plants with "bleeding" or "set official tolerance" for sulfur. Do not use on plants that residues of sulfur and forage crops. Do not use on plants that residues of sulfur are likely to exceed 100 ppm. Do not use on plants that residues of sulfur are likely to exceed 100 ppm. Do not use on plants that residues of sulfur are likely to exceed 100 ppm.

Sure Death Brand
COPPER - SEVIN® - SULFUR
3.4 - 5 - 75 DUST

AN AGRICULTURAL INSECTICIDE - FUNGICIDE

ACTIVE INGREDIENTS:

Copper (in basic copper sulfate) (expressed as metallic)

3.4%

Sulfur

75.0%

INERT INGREDIENTS:

16.6%

TOTAL

100.0%

*SEVIN ® Trademark-Union Carbide Corporation

Sulfur fineness: 93% Passing No. 325 USS Sieve.

CAUTION
KEEP OUT OF REACH OF CHILDREN
 SEE SIDE PANEL FOR ADDITIONAL
 PRECAUTIONARY LABELING

DO NOT BREATHE DUST
 DO NOT CONTAMINATE FEED OR FOODSTUFFS
 AVOID SKIN CONTACT
 WASH WITH SOAP AND WATER AFTER USING
 HARMFUL IF SWALLOWED

DESTROY EMPTY CONTAINERS--
 BURY OR BURN. KEEP OUT OF
 SMOKE.

OBSERVE THE FOLLOWING USE PRECAUTIONS:
 Do not apply when weather conditions favor drift of dust from crop areas treated. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals, or onto adjoining food or forage crops about to be harvested. Contamination of productive waters may destroy essential fish foods. To protect bees, do not apply when crops are in bloom. Remove or cover hives if dusting is necessary. Apply in early morning or evening when there is little or no wind. Sulfur will cause fruit and leaf injury to sulfur-sensitive crops. Do not apply to apricots, comice pears, cranberries, cucurbits (cucumbers, cantaloupes, melons, squash) walnuts, or to other sensitive plants. Sulfur may burn foliage and fruit during periods of high temperature and under certain climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures are exceeding or are likely to exceed 90°F. Do not use sulfur on any crop unless local use has proved safe. The Federal Insecticide, Fungicide, and Rodenticide Administration has set official tolerances for certain uses. In order that residues on food and forage crops shall not exceed established tolerances, use only at recommended rates, and do not exceed the maximum amount specified. Consult state Agricultural Experiment Station for information as to the timing, number of applications, and other details.

DIRECTIONS FOR USE

Apply when air is calm, using ground equipment or aircraft. Make thorough coverage of plants with a uniform deposit of dust. Apply at first sign of infestation and infection and repeat at 5- to 10-day intervals, depending upon severity of disease and insect conditions, but observe use limits given for specific crops. APPLY AT THE RATE OF 30 TO 40 POUNDS PER ACRE.

For the control of diseases persisting after the prescribed time limits have expired, an effective fungicide should be used alone, in a manner which will not leave residues in excess of established tolerances.

RECOMMENDATIONS

BEANS: Ascaris leaf spot, Anthracnose, Armyworm, Bean leaf beetle, cucumber beetle, Downy mildew, Flea beetle, Leafhopper, Velvetbean caterpillar, Rust, Stink bug, Tarnished plant bug.

CABBAGE, BROCCOLI, CAULIFLOWER: Armyworm, cabbage caterpillar, corn earworm, downy mildew, flea beetle, leaf spot. Do not apply within 3 days of harvest.
EGGPLANT, PEPPER: Armyworm, Colorado potato beetle, early and late blight, European corn borer, flea beetle, leafhopper, leaf spot, tomato fruitworm, tomato hornworm, stink bug.

POTATO, TOMATO: Armyworm, early and late blight, European corn borer, Flea beetle, leafhopper, stink bug, tomato fruitworm, tomato hornworm.
Do not apply Foliage application only for potatoes.

STRAWBERRIES: leaf spot, meadow spittlebug, strawberry leaf roller, strawberry weevil. Do not apply within 1 day of harvest.

PEACH NECTARINES: Brown rot, codling moth, cucumber beetle, European earwig, fruit tree leaf roller, Japanese beetle, June beetle, lesser peach tree borer, peach blight, peach leaf curl, peach twig borer, periodical cicada, plum curculio, red banded leaf roller, scales (Lecanium, olive, San Jose), tarnished plant bug. Apply November 15-30 and in Spring before buds swell and in full pink. Do not apply within 3 days of harvest on nectarines and one day of harvest on peaches.

NOTICE: Seller's responsibility is limited to accuracy of contents and confirmation to the chemical description. Buyer assumes all risk and liability for use, storage, and handling, and accepts and uses this product under these conditions.

Manufactured by

W. D. L. CHEMICAL COMPANY

St. Joseph, Mo.
USDA Reg. No. 449-

Denver, Colo.

Orlando, Fla.

Lubbock, Tx.

ACCEPTED

MAY 22 1967

U.S. FEDERAL AGENCIES
FUNCTIONS AND IDENTITIES ACT
FOR SOBERING OFFICE BOARD
SO ORDER NO. 42-100000
ATTACHED COPY...