



CONTENTS

WEIGHTS

Sure Death Brand

STROBANE® 8# E.C.

SOLUBLE FOR AGRICULTURAL INSECTICIDAL PURPOSES

ACTIVE INGREDIENTS:

Terpene Polychlorinates (65% Chlorine)*	72.9%
Petroleum Distillate	20.6%
INERT INGREDIENTS:	6.5%
TOTAL	100.0%

*Strobane, consisting of polychlorinates of camphene, pinene, and related terpenes.

CAUTION KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY LABELING

WARNING: May be fatal if swallowed. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after using. Do not breathe spray mist. Avoid contamination of feed and food stuffs. Keep this product and containers out of reach of children and animals. Do not use or store near heat or open flame.

Do not use on crops other than cotton and avoid spray drift to food crops and to pastures where animals graze. Do not feed trash from treated fields to meat or dairy animals.

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on this label. Rinse empty containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (Keep out of smoke.)

DIRECTIONS:

GENERAL: Mix this material with sufficient water to provide uniform coverage. May be applied by airplane or ground sprayer. **In areas where the bollworm is likely to be a serious problem or where resistant weevils are present, this material should be used in combination with DDT, following local recommendations from State Agricultural Extension Service.**

ACCEPTED
JUL 3 1957
UNDER THE FEDERAL REGISTRATION ACT

AMOUNT TO USE:

<u>Insect</u>	<u>Pints of 8% Concentrate Per Acre</u>	<u>Corresponding lbs. Strobane per acre</u>
<u>Boll Weevil</u>	2-4	2-4
<u>Bollworm</u>	2-4	2-4
<u>Cotton Leaf Worm</u>	2.5	2.5
<u>Cotton Pickworm</u>	2-4	2-4
<u>Stink Bugs and Other Mirids</u>	1-4	1-4
<u>Cotton Fleahopper</u>	1-4	1-4
<u>Thrips</u>	.8 - 2.25	1-4
<u>Cotton Leaf Perforator</u>	4	.8 - 2.25
<u>Stink Bugs</u>	4	4
<u>Cutworms</u>	2-4	4
<u>Fall Armyworm</u>	2-4	2-4
<u>Grasshoppers</u>	2-4	2-4

Use the higher rate for heavy insect infestations and on rank-growing cotton. Repeat applications as needed, usually at 5-day intervals.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured by
 FERTILIZER CHEMICALS COMPANY
 St. Louis, Mo. Denver, Colo. Orlando, Fla. Lubbock, Tex.

USDA REG. NO. 4492