

卷之三十一

**LOW VOL.  
WOOD-KILL  
4 Lb. Concentrate**

## Central Brush, Osage Orange and Most Woody Plants

## **CAUTION**

KEEP OUT OF REACH OF CHILDREN

W. H. D. 1900  
B. C. 1900  
C. B. 1900  
D. C. 1900  
E. C. 1900  
F. C. 1900  
G. C. 1900  
H. C. 1900  
I. C. 1900  
J. C. 1900  
K. C. 1900  
L. C. 1900  
M. C. 1900  
N. C. 1900  
O. C. 1900  
P. C. 1900  
Q. C. 1900  
R. C. 1900  
S. C. 1900  
T. C. 1900  
U. C. 1900  
V. C. 1900  
W. C. 1900  
X. C. 1900  
Y. C. 1900  
Z. C. 1900

Reindeer are seen in the hillsides above the village.

19. The following is a list of the names of the members of the Board of Education.

ACCEPTED

NOV 3 1967

UNDER THE FEDERAL INSECTICIDE,  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 149-721 SUBJECT  
TO ATTACHED COMMENTS.

#### GENERAL INFORMATION

This Coniferous Forest forms 2 M. of the N.E. of  
Alder, and 1/2 M. S.E. of the N.W. E.  
Bog Water. The vegetation is most  
affectionate, the undergrowth of ferns,  
azalea, rhododendron, & other shrubs, the  
trees, tall conifers, can be seen to  
the west of the stream, and the prairie  
larches, birch, aspen, chestnut,  
birch, quaking aspen, and brush  
covered hills. The following woody  
plants are a partial list of which some  
of these species are exceedingly older.

**Soil-Bark Treatment** Large areas and untilled areas of public land in the US are treated more successfully by applying the upper parts of a year's growth of trees, known as "cullings," at 15 pounds per acre. The amount of cullings is about one-half of a cord per acre, or 100 cubic feet. This is applied in late summer or fall, so that it will be available for the winter. The treatment is effective for two years. After this, the soil becomes dry and the bark becomes brittle, so that it can no longer be treated. Old or rough bark requires more spray than smooth or young bark, apply as may be needed. In the first winter months, spray water over areas of treated bark until it is well into the ground. No other spray, except spraying of killing can be expected. Treated bark can never, preferably should not be cut for a period of one year.

**Brunt Treatment.** Follow directions above in Oral Bark Treatment, using entire wet all-purpose bark as well as cut surfaces. Apply at any time, especially in frosty cut areas.

**NOTE—Local conditions may affect the use of herbicides. Consult State Agricultural Experiment Station for occasional information regarding designs and use of application.**

• 100 •

Foliate sprayer tank: Add 1/2 cup of granules after foliation is well developed, starting a drenching spray containing 1/2 to 3 quarts of this material per 100 gallons of water. (For intensive control increase to 2 cups in 2 gallons of water.)

Apply this spray at the following rates per acre according to feasibility of brush: Heavy brush - 100 gallons per acre; medium brush - 50 gallons per acre; light brush - 1 gallon per acre.

Power take off or motor driven  
generator - output power less than  
100 kVA - Aug Committee may accept  
as a suitable alternative

19. The following is a list of the 200 largest cities in the United States, according to the latest available statistics:

卷之三十一

$\tau = \tau_0 + \tau_1$ ,  $\tau_0 = M_0$ ,  $\tau_1 = \tau_1(M_0)$ ,  $\tau_1(M_0) = \tau_1(M_0)$

Digitized by srujanika@gmail.com