WOODBURY CHEMICAL COMPANY

CONTENTS
GALLON

SURE DEATH BRAND

D-P WOOD-KILL

FOR CONTROLLING MIXED BRUSH ON
RIGHT-OF-WAYS AND HIGHWAYS

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

ACTIVE INGREDIENTS

* 2 ethyl-4 methylpentanol ester of 2,4 dichlorophenoxyacetic acid	20.10%
2 ethylhexanol ester of 2,4 dichlorophenoxyacetic ester	13.40%
** 2 ethyl-4 methylpentanol ester of 2,4 dichlorophenoxypropionic acid	20.10%
2 ethylhexanol ester of 2,4-dichlorophenoxypropionic acid	13.40%
INERT INGREDIENTS	33.00%
TOTAL	100.00%

WOODBURY CHEMICAL COMPANY • St. Joseph, Mo.

WOODBURY CHEMICAL COMPANY

CONTENTS
GALLON

SURE DEATH BRAND

)-P WOOD-KILL

FOR CONTROLLING MIXED BRUSH ON
RIGHT-OF-WAYS AND HIGHWAYS

CAUTION
KEEP OUT OF REACH OF CHILDREN
PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

ACTIVE INGREDIENTS

*(2-ethyl-4 methylpentanol ester of 2,4 dichlorophenoxyacetic acid	20.10%
(2-ethylhexanol ester of 2,4 dichlorophenoxyacetic acid	13.40%
*(2 ethyl-4 methylpentanol ester of 2,4-dichlorophenoxypropionic acid	20.10%
(2-ethylhexanol ester of 2,4-dichlorophenoxypropionic acid	13.40%
INGREDIENTS	33.00%
TOTAL	100.00%

*Equivalent [illegible] of 2,4 [illegible] acid
[illegible] of 2,4 [illegible]

[illegible]

CAUTION

[illegible]

NOTE [illegible]

USDA Reg. No. 449-[illegible]

[illegible]

DBURY
MICAL
PANY St. Joseph, Mo.

This concentrate contains 2# of 2,4-D Acid 2# 2,4-D-P, Acid per gallon. This product an be used for the control of brush in such areas as fence rows, right-of-ways, ditch banks, roadsides, and brush covered lots. The following woody plants are a partial list of which one of their species are susceptible:

Alder	Fir	Red Elm
Aspen	Hemlock	Red Maple
Big Leaf Maple	Honeysuckle	Salmonberry
Birch	Oaks	Spruce
Black Cherry	Pine	Sycamore
Buckbrush	Poplar	Tulip Poplar
Black Locust	Red Elder	Willow

(and many other species)

Additional sprays may be necessary for some growth.

FOLIAGE SPRAY: Spray woody growth after foliage is well developed until just before plant goes dormant, using a drenching spray containing 1 to 1½ gallons of Woodkill to 100 gallons water. Use 200 to 600 gallons of spray mixture per acre depending on height and density of growth. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line. Some regrowth can be expected on resistant species such as ash, maple, oak and persimmon.

BASAL BARK TREATMENT: Large brush and small trees can sometimes be controlled more successfully by spraying the basal parts of brush stems and tree trunks to a height of 12 to 15 inches from the ground line. Use a solution of 4 gallons of this product in 96 gallons of diesel oil, fuel oil or similar oil. (1 pint in 3 gallons) Knapsack on power equipment may be used, but complete wetting of the indicated area is necessary, particularly at the ground line. This means spraying until run down or run-off to the ground line is noticeable. Old or rough bark requires more spray than smooth or young bark. Apply at any time, including winter months, except when snow or water prevents spraying to the ground line. Often delayed response of killing can be expected. Treated brush or trees should not be cut for a period of one year.

FOR STUMP TREATMENT. Follow directions above in Basal Bark Treatment being sure to wet all exposed bark as well as cut surface. Apply at any time, preferably to freshly cut stumps.

W
CH

D-

KE
SEE SIDE PANE

WOODE
CHEMIC
COMPA