



CONTENTS

POUNDS

Sure Death Brand

MSMA PLUS SURFACTANT

HERBICIDE FOR POST-EMERGENCE WEED CONTROL IN COTTON

ACTIVE INGREDIENTS:

Monosodium acid methanearsonate . . . 35.42%

INERT INGREDIENTS: 64.58%

TOTAL 100.00%

Total arsenic, all in water-soluble form, expressed as elemental 16.38%. (Contains 4.0 lbs. MSMA per gallon.)

CAUTION KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY LABELING

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas.

KEEP ENTIRE LABEL BEFORE USING THIS PRODUCT. Use of this material in a manner or at any time other than in accordance with the directions may produce plant injury, excessive residues, or other undesirable results.

DIRECTIONS FOR USE

COTTON: This herbicide is useful as a directed spray application to weeds in cotton for post-emergence control of weeds and grasses including Johnson grass, nut grass, water grass, goose grass, dallis grass, and cocklebur.

Mix at a rate of 2 quarts in 40 gallons of water. Apply at a broadcast rate of 40 gallons of this solution per acre. This herbicide is a combination of surfactant (wetting agent) and herbicide. It is unnecessary to add any other surfactant to the spray solution. For band application, apply at a rate of one gallon of this solution for each 1 inch of band width to be treated on cotton grown in 40 inch rows. A second application may be required during humid growing conditions. Make the second application if necessary about 1 to 3 weeks after the first application.

Apply only when cotton is 3 inches high to first bloom. Keep off cotton foliage. Do not apply after first bloom. Do not feed treated foliage to livestock or graze treated areas.

Slight buring and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

Do not contaminate waters used for domestic consumption or by animals, wildlife, and aquatic life or water used for irrigation purposes.

APPLICATION METHODS

This herbicide must be thoroughly dissolved. Fill the tank about half full with water, and with the pump operating add the required amount of herbicide. Finish filling the tank with water and apply.

If preferred, it can be premixed by dissolving the required amount in 3 to 5 gallons of water in an open head bucket. This premix should then be placed in the partially filled spray tank and the tank filled.

Do not store spray solution in tank for a prolonged period. Clean application equipment thoroughly after use by flushing with water.

NOTE: Keep from freezing. Solids may appear in solution under cold storage conditions. Holding solution at room temperature, 70° F., a short time redissolves all solids and solution is acceptable for use.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured by
WOODSBURY CHEMICAL COMPANY

St. Joseph, Mo. Denver, Colo. Orlando, Fla. Lubbock, Texas

USDA Reg. No. 442-