

USAGE WARNING

All fumigant gases, toxic to insects, are also toxic to humans. Use only with adequate ventilation provided for the operator. Do not breath vapor or spray mist. Avoid contact with eyes, skin, and clothing. In case of contact, immediately remove contaminated clothing and shoes. Flush affected skin areas for 15 minutes with water. Wash out eyes with water. Get medical attention. Dry clothing before re-use. Air shoes for at least one week. This fumigant is volatile. Store in a cool place out of direct sunlight. Keep out of reach of children. Do not re-enter bin without an air-line respirator or a full-face gas mask with double black canister for protection against organic vapors and gases. Flammable! Harmful if inhaled or swallowed.

EMERGENCY TREATMENT FOR GAS POISONING: Remove victim to fresh air immediately. Lay victim down, and keep warm and comfortable. Rest and quiet are essential to avoid hysteria. Apply artificial respiration only if breathing has stopped. Call a physician.

Do not reuse container. Destroy when empty.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured by

WOODBURY CHEMICAL COMPANY

St. Joseph, Mo.

Denver, Colo.

Orlando, Fla.


Lubbock, Tex.

USDA Reg. No. 443-

CONTENTS

Powder 30 (4-oz) bags

POUNDS



PIONEER IN ANTICOAGULANT RODENTICIDE!

**RōCON THROW BAGS
DRY MEAL BAIT—KILLS RATS & MICE**

U.S.D.A. Registration No. 449-

Multiple bags are necessary for rat control.

ACTIVE INGREDIENT:

WARFARIN (3-alpha-acetylbenzyl)-

4-hydroxycoumarin025%

QUINOXALINYL SULFANILAMIDE

(Sulfaquinoxaline)025%

INERT INGREDIENTS 99.950%

TOTAL PER CENT 100.000%

ACCEPTED
MAY 23 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 449-512 SUBJECT
TO ATTACHED COMMENTS.

CAUTION

KEEP OUT OF THE REACH OF CHILDREN, domestic animals, and pets. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS.

This bag contains 20 4 oz. Throw bags. These 4 oz.

Throw bags can be tossed into those inaccessible places frequented by rats and mice. Throw bags are their own

bait stations.