

Apply as a spray at the rate of 1/2 to 1 pt. per acre in sufficient water to thoroughly cover plant surfaces, including the leaf whorls.

SWEET CORN: Make first application when 25% of the plants show evidence of leaf feeding and continue at 5-day intervals as long as leaf feeding is evident. From one to three treatments may be required. For control of second brood borers, apply when eggs begin to hatch.

FIELD CORN: Apply 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is present. For control of second brood borers, apply when eggs begin to hatch.

WEAPON 4-E: Recommended for use on corn to control corn rootworm (adult) to protect the silks from damage. Apply as a spray at the rate of 1/2 to 1 pint per acre in sufficient water to thoroughly cover plant surfaces, including the leaf whorls. For aircraft application, use 2 to 10 gallons of water per acre.

SWEET CORN AND FIELD CORN: Apply when insects appear in sufficient number to damage silks and prevent pollination.

NOTE: Do not apply within 14 days of harvest.

IMPORTANT

Keep animals and unprotected persons away during treatment and while there is danger from drift or residues. Bury spillage, clean up area with strong lye solution.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not reuse empty drum. Return to drum re-conditioner or destroy by perforating or crushing and burying in a safe place.

Store in a cool dry place, away from feed and foodstuffs and out of reach of children and domestic animals. Keep container closed.

DANGER !!

Poisonous by skin contact, inhalation, or swallowing. Rapidly absorbed through skin. Repeated exposure may, without symptoms, be increasingly hazardous.

Do not get in eyes, on skin, on clothing. Wear clean, heavy rubber gloves, rubber boots, and clean, waterproof or freshly laundered protective clothing.



CONTENTS
GALLON

Sure Death Brand

WEAPON 4-E

FOR CONTROL OF EUROPEAN CORN BORER

ACCEPTED
9-23-68
449-517

DANGER!!
KEEP OUT OF REACH OF CHILDREN



SEE SIDE PANEL FOR ANTIDOTE STATEMENT AND ADDITIONAL PRECAUTIONARY LABELING

ACTIVE INGREDIENTS:

O-Ethyl O-p-nitrophenyl phenyl phosphonothioate	46.7%
Xylene	47.3%
INERT INGREDIENTS:	6.0%
Total	100.0%

CONTAINS 4 LBS. EPN PER GALLON.
U. S. Pat. 2,503,390

WOODBURY

CHEMICAL

Wash hands, arms, and face thoroughly with soap and warm water after handling and before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and warm water; for eyes, flush with water for 15 minutes and get medical attention.

FIRST AID

Atropine is an antidote. Consult physician for emergency supply. Call a physician at once in all cases of suspected poisoning. Never give anything by mouth to an unconscious person.

Antidotes

INTERNAL: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet.

EXTERNAL: Immediately remove all contaminated clothing and wash skin thoroughly with soap and warm water.

SYMPTOMS OF POISONING

The symptoms of poisoning by organophosphorous insecticides, including EPN, are likely to be as follows:

1. — Digestive upsets; nausea, vomiting, cramps, feeling of pressure under breast bone, diarrhea.
2. — Sweating and excessive salivation.
3. — Restlessness, giddiness, headache.
4. — Muscle tremors, twitching of the facial and eye muscles.

If any of these symptoms are felt, discontinue operation at once; remove person from area; give two atropine tablets (each 1/100 gr.) and call a physician. Do not resume operations with any organophosphorus insecticide, including EPN, until the physician has tested a sample of your blood for organophosphorus effect and tells you it is safe to do so.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured by
WOODBURY CHEMICAL COMPANY
St. Joseph, Mo.; Denver, Colo.; Orlando, Fla.;
Lubbock, Texas

FIELD CORN:

WEAPON 4-E:

SWEET CORN AND FIELD CORN:

NOTE:

IMPORTANT

DANGER !!

ACCEPTED
9-23-68
449-517

CONTENTS
GALLON

Sure Death Brand

WEAPON 4-E

FOR CONTROL OF EUROPEAN CORN BORER

ACTIVE INGREDIENTS	
O-Ethyl O-p nitrophenyl	
diethyl phosphonate	46.7
Xylene	47.3
INERT INGREDIENTS	
	6.0
Total	100.0

**WOODBURY
CHEMICAL
COMPANY • St. Joseph, Mo.**

FIRST AID

Antidotes

INTERNAL:

EXTERNAL:

SYMPTOMS OF POISONING

call a physician.

NOTICE:

Manufactured by
WOODBURY CHEMICAL COMPANY
St. Joseph, Mo., Denver, Colo., Orlando, Fla.,
Lubbock, Texas