

MIXING DIRECTIONS

2 Tablespoonfuls A and
2 Tablespoonfuls B 1 Gallon Water
1 Lb. 13 oz. each A and B: 13 Gallons Water
On items marked B, use only B Formula
mixed 3 tablespoonfuls per gallon of water.

ORCHARD USES

APPLICATION: Apply thoroughly with hand pressure or powder sprayer until leaves are wet on top and bottom to the point of run-off.

SPRAY SCHEDULES

APPLES: Control Scab, Blotch, Sooty Blotch, Fly Speck, Cedar rust, Codling Moths, Curculio, Leaf Hopper, European Red Mite, Red Spider, Red Banded Leaf Roller. Spray when blossoms are pink, when most petals have fallen; ten days after petals fall and at 14 day intervals until early July. Usually six sprayings are required. Do not spray within 15 days of harvest. May injure foliage on some varieties.

PEACHES, PLUMS, APRICOTS: Controls Peach Scab, Curculio, Oriental Fruit Moth, Red Spider, European Red Mite, Red Banded Leaf Roller. Spray with microfine wettable sulfur when 1/4 of flowers are open. Use G. O. F. Garden Spray when petals fall. Follow by three sprayings 7 to 10 days apart, then one in late June and one in mid-July. To prevent harmful residues, use microfine wettable sulfur for Brown Rot control one week before harvest. Do not apply to apricots after petal fall. Do not apply to peaches within 30 days before harvest. Do not apply to plums within 30 days of harvest.

CHERRIES: Controls Brown Rot, Yellow Leaf Spot, Curculio. Apply one spray at petal fall and two additional ones 8 days apart. Spray immediately after harvest, and again two weeks later. Do not apply within 7 days of harvest. May injure foliage on some varieties.

RASPBERRIES: Controls Red Spider, European Red Mite. Spray when first leaves show; repeat at 10 day intervals until blooming begins. Do not spray after fruit development begins.

GRAPES: Controls Aphids, Leaf Hoppers, Berry Moth, Black Rot. Spray when new shoots are 1/2 to 1" long, and spray at 14 day intervals until fruit is about pea size. Do not apply within 14 days of harvest. Emulsion may cause injury to foliage on some varieties.

STRAWBERRIES: Controls Strawberry Leaf Roller, Strawberry Weevil, Spittle Bugs, Strawberry Leaf Beetle. Do not apply within 7 days of harvest. Remove residues at harvest time.

GARDEN USES: Mix according to directions. Spray thoroughly, covering both sides of leaves and stems. Apply when insects or diseases are likely to appear. Repeat every 7 to 10 days as needed. **DO NOT SPRAY WITHIN 10 DAYS OF HARVEST.** Remove residue remaining at harvest time by washing, brushing, wiping, or other effective means.

Will control the following insects and diseases:

BEANS: Aphids, Leaf Hopper, Bean Leaf Beetle, Mexican Bean Beetle, Thrips, Downy Mildew and Red Spider.

CABBAGE, CAULIFLOWER, BROCCOLI,

***TURNIPS, RADISHES:** Cabbage Worms, Diamond Back Moth, Cabbage Looper, Cabbage Root Fly, Cabbage and Turnip Aphid, Cabbage Flea Beetle, Imported Cabbage Worm. Do not apply within 14 days of harvest on turnips and radishes and broccoli.

TOMATOES: Leaf Spot, Anthracnose, Early and Late Blight, Tomato Hornworm, Flea Beetle, Aphids, Leaf Hopper, Psyllid. Start sprays as soon as plants are set, and repeat sprays every two weeks until fruit begins to color. Do not spray within 10 days of harvest. Remove remaining residues at harvest time by washing, brushing, wiping, or other effective means.

FLOWER USE

Roses: Black Spot, Aphid, Rose Chafer, Rose Leaf Roller, Rose Midge, Rose Slug. Start sprays when first leaves form, and repeat every two weeks. Surface of beds should be sprayed.

Flower Insects: Chrysanthemum, Thrips and Tier, Climbing Cut Worm, Tussock Moth, Japanese Beetle, Thrips on Snapdragon, White Fly Sowbugs, Aphid, Mites, and certain other insects.

Trees and Shrubs: Elm Leaf Beetle, Catalpa Sphinx, Leaf Hopper, Cankerworms, Maple Worms, Boxwood Leaf Miner, Minosa Webworm, Tent Caterpillar, Bagworms, Japanese Beetle. Apply when insects appear and repeat at 7 to 10 day intervals.


CAUTION

Keep out of reach of children. Avoid contact with skin. Avoid contamination of feed and foodstuffs. Do not breathe dust or spray mist. May cause irritation of nose and throat. Wash thoroughly after using.

This product will kill fish. Keep out of any body of water. Do not reuse container. Destroy when empty.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

U. S. D. A. Reg. No. 449  569

523

↑ A ↑
8 OUNCES FORMULA

Garden Orchard

Flowers

G. O. F.

GARDEN SPRAY

WETTABLE POWDER

Complete Insecticide-Fungicide

A FORMULA

ACTIVE INGREDIENTS:	
*Captan	16.66%
**Malathion	16.66%
INERT INGREDIENTS	66.68%
Total	100.00%

B FORMULA

ACTIVE INGREDIENTS	
***Methoxychlor	30.70%
Zineb (Zinc Ethylene Bisdithiocarbamate)	25.00%
INERT INGREDIENTS	44.30%
Total	100.00%

*N-Trichloromethyl-N-(4-cylophenoxy)-2,2-dimethylacetamide
**O,O-Dimethyl-dithiophosphate of diethylmercaptosuccinate
***27.02% 2,2-bis-(4-methoxyphenyl)-1,1,1-trichloroethane and 3.68% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Precautionary Labeling

8 OUNCES FORMULA

↓ B ↓

Manufactured by
WOODBURY CHEMICAL
COMPANY

St. Joseph, Mo., Denver,
Colo.; Goulds, Fla.;
Lubbock, Texas

ACCEPTED
4-16-70
449-523
UNDER THE FEDERAL INSECTICIDE ACT
FOR ECONOMIC POISON REGISTERED
EPA REG. NO. 101-101-01 SUBJECT
TO APPLICABLE COMMENTS.