

ACCEPTED
MAR 5 1975

DIRECTIONS

FOR HORN FLY CONTROL ON DAIRY CATTLE:

Use 3 to 5 tablespoonfuls per animal. Dust on poll, neck, back and upper sides of the animal. Rub lightly to insure that the material is worked down beneath the hair. Repeat application once every three weeks.

FOR HORN FLY, LICE AND FLEA CONTROL ON BEEF CATTLE AND HOGS:

Apply 5 to 10 tablespoonfuls per animal. Rub in lightly to insure that the material is worked down beneath the hair. Provide thorough coverage of the animal.

FACE FLY ON BEEF CATTLE AND HOGS:

Direct dust over the face of the animal, repeating treatment daily or as necessary.

Manufactured For

TECHNE CORPORATION

St. Joseph, Mo 64502; Denver, Colo 80216

EPA Reg No. 449-525

Techne dairy dust

A Powder Control for Face Flies, Horn Flies,
Lice, Ticks on Various Animals.

ACTIVE INGREDIENTS:

*Technical Methoxychlor	10.99%
**Technical Piperonyl Butoxide60%
Pyrethrins06%
INERT INGREDIENTS	88.35%
Total	100.00%

*Equivalent to 9.67% 2,2-bis (p-methoxyphenyl) 1,1,1-trichloroethane and 1.32% related compounds.
**Equivalent to 0.48% (butylcarbityl) (6-propyl-piperonyl) ether and 0.12% related compounds.

FOR SHEEP:

Direct dusting after shearing will give temporary relief from ticks (keds). Dust thoroughly on neck, back and upper sides of animal. Use 3 to 5 tablespoonfuls per animal. Rub lightly to insure that the material is worked down beneath the hair. Repeat application once every three weeks.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams.

Do not reuse empty container. Destroy by perforating or crushing and bury in a safe place, away from water supplies.

NOTICE: Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY LABELING