


449-532 Comments



CONTENTS
GALLON

8 LB. TOXAPHENE EMULSIFIABLE

Contains 8 Pounds Toxaphene per Gallon

ACTIVE INGREDIENTS:	
Toxaphene (technical Chlorinated Camphene with 67% to 69% Chlorine)	70%
INERT INGREDIENT	30%
TOTAL	100%

MIXES WITH WATER IN ALL PROPORTIONS
CAUTION: NOT FOR USE OR STORAGE IN
THE HOME ENVIRONMENT.
DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME.

WOODBURY
CHEMICAL
COMPANY • St. Joseph, Mo.

WARNING KEEP OUT OF REACH OF CHILDREN SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

ACCEPTED WITH COMMENTS

DIRECTIONS

USE ONLY A SINGLE APPLICATION

GRASSHOPPERS ON CORN AND COTTON: Under average conditions of rainfall and vegetation, apply Toxaphene at the rate of $1\frac{1}{2}$ pints per acre on corn, 2 pints per acre on cotton, diluted with the volume of water applied per acre by the sprayer. (10 to 30 gallons of water for ground application.) For small areas, use 1 tablespoon in 2 gallons of water per 1,000 square feet. When vegetation is dense and or grasshoppers are adult, increase the dosage to 3 pints per acre.

For Corn: Do not feed treated forage to dairy animals or animals being finished for slaughter.

For Cotton: Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

BAGWORMS ON ORNAMENTALS: Use 1 quart to 100 gallons of water (2 teaspoonfuls per gallon). Spray very thoroughly.

FOR CUTWORM ON CORN: Use 2 pints per acre when applied directly to row; 2 pints per acre when applied over entire area. Mix with sufficient water (10 to 50 gallons water for ground application.) Do not feed treated forage to dairy animals or animals being finished for slaughter.

FOR FALL ARMYWORM ON CORN: Use at the rate of $2\frac{1}{2}$ pints per acre in sufficient water to give good coverage. Do not feed treated forage to dairy animals or animals being finished for slaughter.

FOR YELLOW STRIPED ARMYWORM ON COTTON: As soon as the larvae appear, apply at the rate of 2 pints per acre in sufficient water to give good coverage. (10 to 50 gallons per acre for ground applications). Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

PLANT BUG AND ALFALFA WEEVIL ON SEED ALFALFA: Apply in the early spring or after cutting before new growth is 4 inches tall as insects appear at the rate of 2 pints per acre. In especially dense vegetation, 3 pints per acre may be required. Do not feed treated forage to dairy animals or animals being finished for slaughter. Birds feeding on treated areas may be killed.

WARNING

MAY BE FATAL IF SWALLOWED.

Do Not Contact The Skin.

Do Not Inhale Mist From Sprays.

Do Not Get In Eyes.

Use of a respirator of a type approved by the U. S. Bureau of Mines and goggles is advisable. Wear clean clothing. In case of contact, wash thoroughly with soap and water before eating or smoking and after using. Keep away from children, domestic animals and food. Do not get on clothing.

Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not store near or contaminate feed or foodstuffs. Do not apply to the edible portions of crops within 30 days of harvest or to forage crops if it is to be fed to dairy or beef animals. If applied by aircraft use as indicated above in 3 to 10 gallons of water per acre.

This product will kill fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by perforating or crushing and burying with wastes.

Do not apply when weather conditions favor drift from areas being treated. When treating areas frequented by bees, treat only during those periods of the day when they are not flying. This product is toxic to bees and should not be applied when bees are actively visiting the area. Do not apply where runoff is likely to occur.

NOTE: Consult State Agricultural Experiment Stations for local recommendations.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Manufactured For

WOODBURY CHEMICAL COMPANY

Division of TECHNE CORP.

St. Joseph, Mo. 64502; Denver, Colo.

U.S.D.A. Reg. No. 449-532

449-532