



DIRECTIONS

USE AGAIN INSECT-AFFECTING BEEF CATTLE BARN & STABLES

FOR SPRAYING BEEF CATTLE for horn fly, stable flies, and mosquitoes control. Use 1 quart per 1000 square feet of area. Control would be obtained through spraying of stalls and heat mats. If other spraying areas are used, use 1 quart per 1000 square feet. Use 1 quart per 1000 square feet per adult animal. Repeat as necessary (2-3 weeks).

FOR SPRAYING INTERIOR OF BARN, STABLES, STABLE FLIES, AND MOSQUITOES. Use 1/2 gallon per 1000 square feet of area. Spray thoroughly walls, ceilings, stalls, and manure piles, etc. using 1 gallon of water per 1000 square feet. Repeat as necessary (4-6 weeks). Cover troughs, bins, or other food or water receptacles while spraying. Exclude dairy animals from barns during spraying.

CAUTION!

Avoid contact with eyes. Avoid inhalation of spray mist. May cause skin irritation. Avoid contact with skin. In case of contact, wash with soap and water. Do not use on humans or animals.

Avoid contact with feeds and food stuffs. Do not store near fire or petroleum.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

Do not use on empty stomach. Return to drug dealer for proper disposal, preferably by incineration and burning in a safe place.

Do not use on any type of storage or handling of this material in a structure or area where there is a fire hazard.

Manufactured for
WOODBURY CHEMICAL CO.
Division of TECHNE CORP.
12501 E. Hampden, Denver, CO
U.S.A. Reg. No. 447,811

EACH QUART IS EQUIVALENT TO ONE
POUND 50% WETTABLE
METHOXYCHLOR POWDER

CHERRIES

For cherry Fruit Fly, use 1 quart per 1000 gallons of water. Make first application when flies appear and repeat applications at 7 to 14-day intervals until harvest. Do not apply within 7 days of harvest.

ALFALFA AND CLOVER

For the control of the alfalfa caterpillar, alfalfa webworm, and yellow leg borer, Japanese beetle, grasshopper, etc., apply 2 to 3 quarts per 1000 gallons of water per acre. Apply this product to alfalfa prior to cutting. Do not apply within 7 days of harvest or grazing. For alfalfa spraying, apply 2 to 3 quarts per 1000 gallons of water per acre. Use the care of water tests. Do not use in pastures with a high percentage of alfalfa.

EMPTY GRAIN BINS

As an insecticide for Rice, Wheat, Granary Weevil, Lesser Grain Borer, and other insects. Mix this product with the insecticide mixture, prior to application. It is possible to use this product as a preventive treatment to help prevent infestation of newly stored grain. Apply 2 quarts per 1000 gallons of water. Make application to thoroughly cleaned surfaces of the bin. Use 2 quarts of spray mixture per 1000 square feet. Allow sprayed surface to dry before storing.

COMPATIBILITY

Do not use with materials containing sulfur. Do not use with materials incompatible with it. Do not use with materials within 14 days of application of materials containing sulfur.



CONTENTS
GALLON

METHOXYCHLOR

Emulsifiable

Contains 2 Lbs. of Methoxychlor per Gallon

CAUTION
KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

ACTIVE INGREDIENTS	
*Methoxychlor, Technical	24%
Methylated Naphthalenes	63%
Dimethyl Phthalate	10%
INERT INGREDIENTS	
Total	3%
	100%

*24% Methoxychlor, technical is equivalent to 21% 2,2-bis(p-methoxyphenyl)1,1,1-trichloroethane and 3% other isomers and reaction products

**WOODBURY
CHEMICAL
COMPANY • St. Joseph, Mo.**