[449-538]

WARNING

May be fatal if swallowed. Absorbed through skin. Do not breathe vapor or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near food and feed products.

USAGE CAUTIONS

Do not use if material does not emulsify.

Workers entering treated fields within 24 hours should wear protective clothing.

Do not use or store near heat or open flame.

Do not store at temperatures below 0° F.

This product is toxic to fish and wildlife. Keep out of any body of water.

This product is toxic to bees and should not be applied when bees are actively visiting the area. Birds feeding on treated areas may be killed.

Apply this product only as specified on this label.
Rinse equipment and containers and DISPOSE OF WASTES by burying at least 18 inches deep in non-crop lands away from water supplies. Do not re-use containers, smash or punch holes in it and bury with wastes.

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

DIRECTIONS FOR USE

Apply the listed amount per acre when insects first appear and repeat as required to maintain effective control. Use in sufficient water for thorough coverage, unless otherwise noted for listed crops. Coverage of upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use pattern, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in ( ) following the crop.

When applying this material by aircraft, mix the recommended amount with sufficient water to provide a minimum of 1 gallon finished spray per acre. Where more than 2 quarts are recommended mix with sufficient water to provide a minimum of finished spray equal to twice the amount of Ethion Miscible used. Do not use in undiluted form.

BEANS (SUCCULENT AND DRY) (2). Bean Leaf Beetle, Mexican Bean Beetle, Two-spotted Spider Mite — 0.5 to 1 pt. per acre. Potato Leafhopper — 1 pt. per acre. Do not feed treated foliage to livestock.

BEANS (SUCCULENT AND DRY) (4). Lygus Bugs — 1 pt. per acre. Do not feed treated foliage to livestock.

CORN FIELD (50). Pacific Spider Mite, Two-spotted Spider Mite — 1 pt. per acre. Do not make more than 1 application after ear formation. Do not feed treated fodder to livestock.

COTTON. Aphids — 1 to 2 pts. per acre. Cotton Leafworm — 1 pt. per acre. Spider Mites — 0.5 to 1 pt. per acre. Spider Mites, Lygus Bugs and Other Mirrids on West Coast only — 2 to 3 pts. per acre.

Do not apply after bolls open. Do not graze dairy or meat animals in treated fields.

CUCUMBERS (0). SUMMER SQUASH (0). Serpentine Leaf Miner, Two-spotted Spider Mite — 1 pt. per acre.

EGGPLANT (0). Two-spotted Spider Mite — 0.5 to 1 pt. per acre.
MELONS (0): Melon Leaf Miner, Serpentine Leaf Miner, Two-spotted Spider Mite — 1 pt. per acre. Melon Leafhopper — 1.5 to 2 pts. per acre. Petrobia Mite (California only) — 1 to 2 pts. in a minimum of 10 gals. water per acre.

PEPPERS (0). Two-spotted Spider Mite — 1 to 2 pts. per acre. Do not allow livestock to graze on treated area before harvest. Do not feed treated forage to livestock.

WOODBURY CHEMICAL COMPANY

CONTENTS
GALLON

ETHION 4 EC

(Contains 4 Lbs. Ethion per Gallon)

ACCEPTED

ACTIVE INGREDIENTS:

*Ethion (O,O,O',O'-tetraethyl S,S'-methylene bisphosphorodithioate)	47.1%
Petroleum Distillate	45.0%
INERT INGREDIENTS:	7.9%
	100.0%

WARNING

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

WOODBURY CHEMICAL COMPANY • St. Joseph, Mo.

SORGHUM. [illegible] Mites — [illegible] pts. per acre [illegible] 30 [illegible] dormant use.

STRAWBERRIES. [illegible] Red Mite, Rose Aphid — [illegible] pts. per acre [illegible] dormant use.

TOMATOES. [illegible] Mites — [illegible] pts. per acre [illegible] dormant use.

APPLES. [illegible] Red Mite, Rose Aphid, San Jose Scale — [illegible] pts. per acre [illegible] dormant use.

CHERRIES. [illegible] Aphid — [illegible] pts. per acre [illegible] dormant use.

NECTARINE, PEACH, PLUM, PRUNE. Clover Brown Aphid, Mites, European Red Mite [illegible] Aphid and Leaf Curl Plum Aphid, Mites, Plum Aphid [illegible] pts. per acre [illegible] dormant use. Olive (Pulvinaria) Scale — [illegible] pts. per acre [illegible] dormant use. San Jose Scale — [illegible] pts. per acre [illegible] dormant or delayed dormant use.)
NECTARINE, PEACH, PLUM, PRUNE. Clover Brown Aphid, Mites, European Red Mite [illegible] pts. per acre [illegible] water or 2.5 to 3.5 pts. per acre [illegible] dormant use.

PEARS. [illegible] pts. per acre [illegible] dormant use.

HOME GROUNDS. [illegible] TOP SOIL [illegible] 2500 sq. ft. [illegible] soil [illegible] tion from the bottom [illegible] ars: 5 to 25 [illegible] Keep children, pets and livestock off treated areas until spray has dried.

ORNAMENTALS (GREENHOUSE). [illegible] Use 10 [illegible] pts. per 100 gals. [illegible] per square foot. Wait [illegible] Ethion protection.

ORNAMENTALS (OUTDOOR). [illegible] pt. per 100 gals. [illegible]
TURF (COMMERCIAL USE). [illegible] 3.5 to 0.5 pt. per 100 sq. [illegible] to 1.67 to 2.5 pts. in 100 gals. [illegible] treat one acre. [illegible] It is important that [illegible] to reach the [illegible] to treatment.

CAUTIONS. [illegible]

TERMS. [illegible]

WOODBURY CHEMICAL COMPANY
[illegible] 64502 [illegible]