



TAKE TIME
OBSERVE LABEL
DIRECTIONS

WEED-KILL



Green With **ENVY** Lawn Food



ACTIVE INGREDIENTS

2,4-Dichlorophenoxyacetic acid	100%
2,4,5-Trichlorophenoxyacetic acid	100%
Dimethylamine Salt of 2,4,5-Trichlorophenoxyacetic acid	25%
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid	04%
INERT INGREDIENTS	98.71%
TOTAL	100.00%

*Equivalent to 85% 2,4,5-Trichlorophenoxyacetic acid
 *Equivalent to 20% 3,6-Dichloro-o-anisic acid (dicamba)
 EPA Reg. Number 449539

Net Weight 40 Lbs Covers 7,000 Sq Ft

ENVY LAWN FOOD
Pelletized for easy spreading


GUARANTEED ANALYSIS

Total Nitrogen (N)	14.0%
Available Phosphoric Acid (P ₂ O ₅)	14.0%
Soluble Potash (K ₂ O)	14.0%
Iron (FE) (Elemental)	14.0%
Sulfur (Combined chemicals)	14.0%

(Iron, Nitrogen and Sulfur derived from chemically combined Ferric Ammonium Sulphate)
 Calcium Carbonate (lime) Equivalent per ton 1,700 lbs.

JAN. 18, 1971

WOODBURY CHEMICAL COMPANY ST. JOSEPH, MISSOURI
 DIVISION OF TECHNE CORP DENVER, COLORADO 80216



WEEDS CONTROLLED

Dandelion	Ragweed	Lambsquarters	Puncture
Morning Glory	Mustard	Canada Thistle	Peppercorn
(Windweed)	Smartweed	Spanish nettle	Furze
Chickweed	Prostrate Spurge	Poison	English Daisy
Annual Clover	Veronica	Red Top	Buckhorn
Knockout	Thymus	St. John's	Mullein
Black Medick	Purslane	White Top	Pigweed
Pennycress	Runaway	White Top	Groundsel
Chickweed			St. John's

USES: This is a pre-emergent herbicide. It is used to prevent the germination of weeds and grasses. It is applied to the soil before the weeds and grasses have a chance to grow. It is most effective when applied to the soil before the weeds and grasses have a chance to grow. It is most effective when applied to the soil before the weeds and grasses have a chance to grow.

APPLY: at the rate of 40 lbs per acre. Apply to the soil before the weeds and grasses have a chance to grow.

SPREADER SETTINGS


Scott's	4
Dynalene	4
Ward	3
Ward	4
Ward	4
Ward	4
Ward	4
Ward	4
Ward	4

For best results, apply this product to the soil before the weeds and grasses have a chance to grow. It is most effective when applied to the soil before the weeds and grasses have a chance to grow.

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes, clothing, do not store near seeds. Avoid application in vegetable gardens, lawns, or ornamental plants other than turf. Avoid application to the soil of trees or shrubs. Do not contaminate water used for irrigation or domestic purposes.

WARNING: This is an effective product. To avoid injury to your lawn, it must be used according to directions.

FREIGHT AND POSTAGE PAID BY ADDRESSEE
 APPLICABLE FREIGHT AND POSTAGE
 GREAT PLAINS CHEMICAL CORP.
 DES MOINES, IOWA 50319



NET WEIGHT 40 LBS.

COVERS
7,000 SQ FEET

6.89

WEED-KILL

+



EMMY is a registered trademark of EMMY Chemicals, Inc. © 1968 EMMY Chemicals, Inc. All rights reserved. EMMY is a registered trademark of EMMY Chemicals, Inc. All rights reserved.

Contains Dicamba and 2-4-D

CAUTION
KEEP OUT OF REACH OF CHILDREN

14-0-0

Plus 3% Iron and
15% Sulphur



TAKE TIME
OBSERVE LABEL
DIRECTIONS

WEED-KILL

+

Green With **ENVY** Lawn Food



ACTIVE INGREDIENTS

lauroyl Ester of 2,4-dichlorophenoxyacetic acid*	1.00%
Dimethylamine Salt of dicamba** (3,6-dichloro-o-anisic acid)	.25%
Dimethylamine Salt of related acids	.04%
INERT INGREDIENTS	98.71%
TOTAL	100.00%

*Equivalent to 65% 2,4-dichlorophenoxyacetic acid
**Equivalent to 206% 3,6-dichloro-o-anisic acid (dicamba)

EPA REG. NUMBER 449539

Net Weight 40 Lbs. Covers 7,000 Sq. Ft.

ENVY LAWN FOOD
Pelletized for easy spreading



GUARANTEED ANALYSIS

Total Nitrogen (N)	14.0%
Available Phosphate Acid (P ₂ O ₅)	.0%
Soluble Potash (K ₂ O)	.0%
Iron (FE) (Elemental)	3.0%
Sulfur (Combined chemically)	15.0%

(Iron, Nitrogen and Sulfur derived from chemically combined Ferric Ammonium Sulphate)
Calcium Carbonate (lime) Equivalent per ton 1,700 lbs.

MANUFACTURED FOR
WOODBURY CHEMICAL COMPANY ST. JOSEPH, MO. 64502
DIVISION OF TECHNE CORP. DENVER, COLORADO 80216

449-539



WEEDS CONTROLLED

Dandelion	Ragweed	Lambquarters	Plantain
Broomrape	Mustard	Canada Thistle	Peppercorn
(bindweed)	Smartweeds	Spanish nettle	Furlydock
Chickweed	Prostrate Sours	Poorjon	English Daisy
Annual Clover	Velvetleaf	Heat-Set	Blacktop
Knotted	Dollarweed	Herb	Maintained
Black Medic	Purslane	Waterwing	Plowseed
Pennywort	Sunflower	Whitecockle	Knave (German Moss)
Cocklebur			

USES: Use on ornamental turf only. Use when any of the above weeds are young and growing rapidly. Old established full grown weeds are hard to kill. Safe on most cool and warm season grasses such as bluegrass (Kentucky, Merion and annual), fescues, perennial ryegrass, red top, Bermudagrass, Zoysia, and others. Do Not use on centipede grass, St. Augustine grass, dichonora or where desirable bentgrass or clovers are present. Avoid application under dense shade of trees and bushes. Do Not use on vegetables or ornamentals or on newly seeded turf. Cut new grass three to four times before making application. Wait three weeks after application before reseeding or spot seeding established turf.

APPLY at the rate of 40 lbs. per 7,000 sq. ft. See spreader settings below.

SPREADER SETTINGS

Scott's	4
Cyclone	4
Ward's	2 1/2
Imperial Husky C1667	4
Imperial Suburban C1662C	4
Imperial C1665	4
Imperial Professional	4
True Temper	4
Jackson	4

For twice over or crisscross application, use one-half (1/2) normal rate. Spreader settings should always be checked for accuracy before application on entire yard. Sweep pellets from sidewalk before watering. Iron will stain concrete. Do not apply more than recommended amount. Water lawn thoroughly after application but do not flood lawn.

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes, or clothing. Do not store near seeds. Avoid application on vegetable gardens, flower, or desirable plants other than turf. Avoid application on drip zone of trees or shrubs. Do not contaminate waters used for irrigation or domestic purposes. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not store near other fertilizers, seeds, insecticides or fungicides.

WARNING: Weed Kill + Envy is an effective product to kill most weeds but must be used according to directions.
Do Not Reuse Container. Destroy by burying in a safe place. Do Not Burn.

FREIGHT SHIPPING BAG
Meeting Requirements of
APPLICABLE FREIGHT CLASSIFICATION
Guaranteed By
GREAT PLAINS BAG CORP
DES MOINES, IOWA & JACKSONVILLE, ARIZONA

A-4133



TAKE THE
SERIOUS LABEL
DIRECTIONS

WEED-KILL



Green With

ACTIVE INGREDIENTS
 *s-octyl Ester of 2,4-dichlorophenoxyacetic acid**
 Dichlorophenoxyacetic acid
 Dimethylamine Salt of dicamba** (3,6-dichloro-o-anisic acid)
 Dimethylamine Salt of related acids

INERT INGREDIENTS

TOTAL
 **Equivalent to .88% 2,4-dichlorophenoxyacetic acid
 *Equivalent to .208% 3,6-dichloro-o-anisic acid (dicamba)

Net Weight 40 Lbs. Cover

ENVY LAWN FOOD
 Pelletized for easy spreading

GUARANTEED ANALYSIS

Total Nitrogen (N)
 Available Phosphoric Acid (P₂O₅)
 Soluble Potash (K₂O)
 Iron (Fe) (Elemental)
 Sulfur (Combined chemically)

(Iron, Nitrogen and Sulfur derived from chemical
 Ferrous Ammonium Sulphate)
 Calcium Carbonate (lime) Equivalent per ton 1

MANUFACTURED FOR
WOODBURY CHEMICAL COMPANY ST. JOSEPH, MO.
DIVISION OF TECHNIE CORP. DENVER, CO.

WEEDS CONTROLLED

- | | | |
|-----------------------------|------------------|----------------|
| Dandelion | Ragweed | Lambquarter |
| Flaming Glory
(bindweed) | Mustard | Canada Thistle |
| Chickweed | Smartweeds | Spanish nettle |
| Annual Clover | Prostrate Spurge | Pooryes |
| Knottedweed | Velvetleaf | Heal All |
| Black Magic | Dollarweed | Henbit |
| Pennywort | Purslane | Waterhemp |
| Cocklebur | Sunflower | Amaranth |

USES: Use on ornamental turf only. Use when any of the above weeds are growing rapidly. Old established full grown weeds are hard to kill. Warm season grasses such as bluegrass, Kentucky, Marion and bermudagrass, red top, bermudagrass, zoysia, and others, are not controlled. St. Augustine grass, perennials, and where desirable present. Avoid application on ornamentals, shrubs, and bushes or ornamentals or on newly seeded turf. Cut new grass before making application. Wait three weeks after application before establishing turf.

APPLY at the rate of 40 lbs. per 7,000 sq. ft. See spreader setting.

SPREADER SETTINGS

- Scott's
 Cyclone
 Ward's
 Imperial Mustang C1667
 Imperial Suburban C1662C
 Imperial C1685
 Imperial Professional
 True Temper
 Jackson

For twice over or crescent application, use one-half (1/2) rate. Accuracy should always be checked before application on concrete. Iron will stain concrete. Do not contaminate waters used for irrigation or domestic purposes. Water lawn thoroughly after application.

CAUTION: KEEP OUT OF REACH OF CHILDREN. Avoid contact with clothing. Do not store near seeds. Avoid application on vegetables, desirable plants other than turf. Avoid application on drift. Do not contaminate waters used for irrigation or domestic purposes.

This product may injure susceptible plants in the immediate vicinity of fertilizers, seeds, insecticides or fungicides.

WARNING: Weed Kill + Envy is an effective product to kill weeds according to directions.

Do Not Reuse Container. Destroy by burying in a safe place.

FREIGHT SHIPPING BAG
Meeting Requirements of
APPLICABLE FREIGHT CLASSIFICATION
 Guaranteed By
GREAT PLAINS BAG CORP.
 205 NORTH 10TH & JACKSONVILLE, ARLINGHAM, MASS.

NET WEIGHT 40 LBS.

COVERS
7,000 SQ FEET

6.89

WEED-KILL



ACTIVE INGREDIENTS

Isopropyl ester of 2,4-dichlorophenoxyacetic acid	1.00%
Dimethylamine Salt of the ester (2,4-Dichloro-D-amine salt)	25
Dimethylamine Salt of related acids	26

INERT INGREDIENTS	68.71
TOTAL	100.00%

Equivalent to 66% 2,4-Dichlorophenoxyacetic acid
Equivalent to 200% 2,4-Dichloro-D-amine salt (66.71%)

Contains Dicamba and 2-4-D

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

14-0-0

Plus 3% Iron and
15% Sulphur

ACTIVE INGREDIENTS

Isocetyl Ester of 2,4-dichlorophenoxyacetic acid	1.00%
Methylammonium Salt of Dicamba* (3,6-dichloro-o-anisic acid)	.25%
Methylammonium Salt of related acids	.24%
INERT INGREDIENTS	98.71%
TOTAL	100.00%

* Equivalent to .66% 2,4-dichlorophenoxyacetic acid
* Equivalent to .200% 3,6-dichloro-o-anisic acid (Dicamba)

Cameraman's Note

Poor Copy

Contains Dicamba and 2-4-D

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

14-0-0

**Plus 3% Iron and
15% Sulphur**