


NET WEIGHT

ACCEPTED
May 31 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PESTS REGISTERED
 UNDER NO. *419-541*



50 POUNDS

TOXAPHENE

15% GRANULAR INSECTICIDE

AN INSECTICIDE CONCENTRATE ESPECIALLY PREPARED FOR
 EUROPEAN CORN BORER CONTROL.

ACTIVE INGREDIENT:	Toxaphene (technical chlorinated camphene containing 67-69% chlorine)	by Wt.
		15
INERT INGREDIENTS:		85
		<hr/>
TOTAL		100

WARNING: May be fatal if swallowed. Do not get on skin, in eyes or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe dust. Avoid contamination of feed or foodstuffs. Do not use around dairy barns. Do not use on cucurbits.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label. Do not apply where runoff is likely to occur. Dispose of wastes by burying on noncrop lands away from water supplies. Dispose of containers by burning. Keep out of smoke.

NOTICE: This product is toxic to bees and should not be applied when bees are actively visiting the area.

ANTIDOTE: Internal — Give sodium chloride 15g. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately. External — Wash with soap and water.

DIRECTIONS FOR USE: European Corn Borer Control — Apply 13½ pounds per acre by either ground equipment or by airplane. A second application may be required 7 to 10 days later. A third application may be required for control of second-brood borers. Consult State Extension Service specialists for most effective application dates.

Do not feed ensilage made from treated corn plants to dairy animals or animals being finished for slaughter. Do not allow dairy animals or animals being finished for slaughter to graze or feed on treated stover.

NON-WARRANTY: Woodbury warrants that this insecticide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this insecticide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

WARNING
 KEEP OUT OF REACH OF CHILDREN



WOODBURY CHEMICAL CO.
 Division of TECHNIE CORP

St. Joseph, Mo. 64502
 Denver, Colo.

Manufactured for:

EPA Reg. No. 449-541
 Rev. No. 1

NET WEIGHT



50 POUNDS

TOXAPHENE

15% GRANULAR INSECTICIDE

AN INSECTICIDE CONCENTRATE ESPECIALLY PREPARED FOR
EUROPEAN CORN BORER CONTROL.

ACTIVE INGREDIENT:	% by Wt.
Toxaphene (technical chlorinated camphene containing 67-69% chlorine)	15
INERT INGREDIENTS:	85
TOTAL	100

WARNING: May be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Can be absorbed through skin. Wash thoroughly after using. Do not breathe dust. Avoid contamination of feed or foodstuffs. Do not use around dairy barns. Do not use on cucurbits.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label. Do not apply where runoff is likely to occur. Dispose of wastes by burying on noncrop lands away from water supplies. Dispose of containers by burning. Keep out of smoke.

NOTICE: This product is toxic to bees and should not be applied when bees are actively visiting the area.

ANTIDOTE: Internal — Give sodium chloride 15g. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear. Call a doctor immediately. External — Wash with soap and water.

DIRECTIONS FOR USE: European Corn Borer Control — Apply 13 1/2 pounds per acre by either ground equipment or by airplane. A second application may be required 7 to 10 days later. A third application may be required for control of second-brood borers. Consult State Extension Service specialists for most effective application dates.

Do not feed ensilage made from treated corn plants to dairy animals or animals being finished for slaughter. Do not allow dairy animals or animals being finished for slaughter to graze or feed on treated stover.

NON-WARRANTY: Woodbury warrants that this insecticide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this insecticide is reasonably fit for its intended purpose. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

WARNING
KEEP OUT OF REACH OF CHILDREN

WOODBURY CHEMICAL CO.

Division of TECHNE CORP.

Manufactured for:

St. Joseph, Mo. 64502;
Denver, Colo.

EPA Reg. No. 449-541
Rev. No. 1