

NET CONTENTS

GALLONS



STOX-SPRAY

CONCENTRATE

For Use On Beef Cattle, Sheep and Swine. Controls Horn Flies, Lice, Sarcoptic and Psoroptic Mange, Ticks.

ACTIVE INGREDIENTS

Toxaphene (Technical Chlorinated Camphene containing 67% to 69% Chlorine)	42.0%
Gamma Isomer of Benzene Hexachloride from Lindane ..	1.9%
Xylene Range Aromatic Petroleum Solvent	49.7%
INERT INGREDIENTS	6.4%

Total.....100.0%

Protect From Temperatures Below 32° F.

WARNING: Keep Out of Reach of Children. See other warnings on back panel.

MANUFACTURED FOR **MILLER CHEMICAL CO.** QUALITY PRODUCTS
OMAHA, NEBRASKA

DIRECTIONS FOR USE

BEEF CATTLE — For control of **HORN FLIES, LICE, TICKS, SARCOPTIC MANGE.** Use 1 qt. STOX-SPRAY to 26 gallons water (7 teaspoonfuls to one gallon water). Apply as a spray treating animals thoroughly. Repeat after 2 to 3 weeks if needed. Do not use on dairy animals or animals under 3 months of age. Do not treat within 30 days of slaughter.

SHEEP — For control of **LICE, KEDS, TICKS, PSOROPTIC SCAB OR SARCOPTIC SHEEP SCAB.** Use 1 qt. STOX-SPRAY to 26 gallons water (7 teaspoonfuls to one gallon water). Apply as a spray treating animals thoroughly. Repeat after 2 to 3 weeks if necessary. Do not treat within 30 days of slaughter. Do not treat animals under 3 months of age.

SWINE — For control of **LICE, SARCOPTIC MANGE.** Use 1 qt. STOX-SPRAY to 26 gallons water (7 teaspoonfuls to one gallon water). Apply as a spray treating animals thoroughly. Repeat after 2 to 3 weeks if necessary. Do not treat within 30 days of slaughter. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.

RESIDUAL SPRAY — For control of **HOUSE FLIES, STABLE FLIES, MOSQUITOES.** Use 3 gallons STOX-SPRAY to 25 gallons water (1 pint to one gallon water). Apply spray at the rate of 1/2 to 1 gallon per 1000 sq. ft. of surface. Thoroughly treat exteriors of barns, stock sheds and other farm buildings as well as

DIRECTIONS FOR USE

manure, vegetation and refuse. Do not use in dairy barns, milk rooms or poultry houses. Do not allow livestock to graze on treated vegetation. Repeat as needed.

BACKRUBBERS — For control of **HORN FLIES** and as an aid in reducing **LICE** on beef cattle. Mix 1 gallon of STOX-SPRAY with 10 gallons diesel fuel or kerosene. Use 1 gallon of this mixture to saturate 20 ft. of cable. Repeat as necessary to keep cable dampened. Do not use on dairy animals. Do not treat beef cattle within 30 days of slaughter, or animals under 3 months of age.

WARNING: May Be Fatal If Swallowed! May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. In case of contact, wash immediately with soap and water.

Do not store near or contaminate feed and food stuffs.

Do not use or store near heat or open flame.

Do not treat emaciated or lactating animals.

Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

Rinse spray equipment and containers and dispose of liquid wastes in a pit in non-crop lands located away from water supplies.

U.S.D.A. REG. NO. 450-133

2-67D

