

ACCEPTED
 JUN 12 1967
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 456-4 SUBJECT
 TO ATTACHED COMMENTS.

NET WEIGHT

2 LBS.



CHEMFORM COPPER 53

U.S.D.A. Reg. No. 456-4

ACTIVE INGREDIENTS:

Basic Copper Sulfate	98.0%
Containing 53.0% Copper as Metallic Fixed	
ERT	2.0%
<hr/>	
Total	100.0%

CAUTION: Poisonous if taken internally. Give an emetic of mustard and warm water. Call a Physician.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

CHEMICAL FORMULATORS, INC.

USE AS A SPRAY FOR THE CONTROL OF:

Early Tomato Blight
 (Macrosporium)
Tomato Leaf Spot
 (Septoria)
Anthracnose
Late Tomato Blight

Downy Mildew
Grape Black Rot
Bacterial Wilt on
Cucumbers
Bitter Rot on Apples
Potato Blight

DIRECTIONS

For use against the above diseases and others which are controlled by the use of Fixed Copper. Use 1 lb. in 25 gals. of water or 1 tablespoon per gallon. Use hand or tank type sprayer. Apply to foliage and stem of plants covering as much of the surface as possible. Spray every 7 to 10 days during the season, oftener during wet weather or when fungi are most active.

CHEMFORM COPPER 53 is compatible with lime, DDT, sulphur, Rotenone and most other insecticides and may be mixed with them. For cucumbers it is recommended that one half lb. of hydrated lime be added with each 1 lb. of CHEMFORM COPPER 53.

