

NET CONTENTS

ONE QUART

DIOXY²⁻²⁸⁻⁶⁸ WEED KILLER⁴⁵⁶⁻¹⁶

(FARM AND HOME)

For control of certain species of broad leaved weeds in lawns, pastures and cultivated crops.

Active Ingredients:

*Dimethylamine Salt of 2, 4-Dichlorophenoxyacetic acid 49.4%

Inert Ingredients:

TOTAL 50.6%
100.0%

*Equivalent to 41.03% of 2, 4-Dichlorophenoxyacetic acid

Manufactured by

CHEMICAL FORMULATORS, INC.

NITRO, W. VA.

KNOXVILLE, TENN.

WARNING

1. Apply this weed killer directly to, or permit to come in contact with vegetables, nut trees, grapes, flowers, fruit trees, alfalfa, cotton, ornamentals or any other desirable plants. Do not spray on windy days since drift is likely to be injurious to nearby sensitive plants. Use a coarse spray instead of a fine mist. Do not use on bent grass lawns, freshly seeded lawns until grass is well established. Legumes and white clover are highly damaged from 2, 4-D materials. Do not contaminate streams or irrigation ditches. Excessive amounts may cause temporary unproductivity of soil and inhibit seed germination. Because of the danger of contamination, it is recommended that the material not be used for applying this weed killer nor be used for any other purpose. Do not use near seed fertilizers, insecticides or fungicides. Do not reuse empty containers. Discard in a safe place.

DIRECTIONS

Chemform Farm and home Weed Killer is an effective 2,4-D selective weed killer, designed to control broad-leaved weeds occurring in lawns, pastures and corn fields.

CONTROLS

broad leaved	poison ivy	Wild Grapes
Plantain	Ground Ivy	Mustard
Fieldweed	Smartweed	Yellowwort
Wild Cornish	Black	Poison Oak
Glory	Angelica	Wild Honey Suckle
San Joaquin	Chicweed	Virginia Creeper

Lawns, PASTURES

For control of weeds in drives and pastures, use 1/2 to 1 pint per acre in 1 gallon of water and spray weed, spray evenly to cover 500 sq. ft. Best results if weeds are young and making vigorous growth. In 4-6 weeks, if required. Do not graze dairy on treated areas for 7 days after treatment.

CORN AND SORGHUM

For control of weeds, use 1/2 to 1 pint per acre in sufficient water for coverage (30-50 gallons). When corn is 6 to 16 inches tall. For sorghum use 1/2 to 1 pint per acre as tall and spray. Use drop nozzle when corn or sorghum is over 16 inches tall.

CAUTION

May cause skin irritation. Avoid contact with skin, and clothing. Harmful if swallowed. Do not use or spray mist. Wash off with water, e.

Manufacturer generally warrants material to be effective for indicated uses. Failure of buyer to properly handle, and store said material voids all warranties.

ONE QUART

DIOXY ED KILLER

FARM AND HOME

ACCEPTED
2-28-68
456-16

of certain species of broad-leaved
lawn, pastures, and cultivated crops:

Ingredients:
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid 49.4%

Ingredients: 50.6%
TOTAL 100.0%

Content to 41.03% of 2,4-Dichlorophenoxyacetic acid

Manufactured by

AL FORMULATORS, INC.

VA KNOXVILLE, TENN

WARNING

Do not apply this weed killer directly to, or permit it to come in contact with vegetables, fruits, flowers, shrubs, cotton, ornamentals or any other desirable plants. Do not spray on windy days since drift is likely to be injurious to nearby sensitive plants. Use a coarse spray instead of a fine mist. Do not use on bent grass lawns. Freshly seeded lawns until grass is well established. Legumes and white clover are usually damaged from 2,4-D materials. Do not contaminate streams or irrigation ditches. Excessive amounts may cause temporary, unproductivity of soil and inhibit seed germination. Because of the danger of contamination, it is recommended that the container used for applying this weed killer not be used for any other purpose. Do not use near seed, fertilizers, insecticides or fungicides. Do not reuse empty containers. Discard in a safe place.

DIRECTIONS

Chlorom Form and Home Weed Killer is an effective 2,4-D selective weed killer, designed to control most broad-leaved weeds occurring in lawns, pastures and corn fields.

CONTROLS

broad leaved	Poison Ivy	Wild Grapes
Plantain	Ground Ivy	Mustard
Red-top	Quack	Pennywort
Wild Cornin	Yuck	Poison Oak
Glory	Language	Wild Honey Suckle
Don't mistle	Chickweed	Virginia Creeper

LAWNS, PASTURE

For control of weeds in drives and pastures, mix 1 tablespoon in one gallon of water and spray weedy areas. Spray evenly to cover 500 sq. ft. Best results obtained if weeds are young and making vigorous growth. Repeat in 4-6 weeks, if required. Do not graze dairy animals on treated areas for 7 days after treatment.

CORN AND SORGHUM

For control of weeds, use 1/2 to 1 pint per acre in sufficient water for coverage (30-50 gallons). Spray when corn is 6 to 18 inches tall. For sorghum use 1 qt. per acre as fall and spring sprays. Use drop nozzles when corn or sorghum is over 10 inches tall.

CAUTION

May cause skin irritation. Avoid contact with eyes, skin, and clothing. Harmful if swallowed. Do not breathe fumes or spray mist. Wash off clothing.

(27A)

28

Manufacturer guarantees material to be effective for indicated uses. Failure of buyer to properly use, handle, and store said material voids all warranties.