



Net Contents

1 Gallon



# Methoxychlor

## 2 EC

U.S.D.A. Reg. #456-93

INSECTICIDE FOR CONTROL OF VARIOUS INSECTS ON SHADE TREES, FORAGE CROPS AND LIVESTOCK.

Active Ingredients:	
Methoxychlor, Technical* -----	25.0%
Petroleum Distillate (Xylene Bottoms) -----	70.0%
Inert Ingredients:	
Alkyl Aryl Sulfonate (Wetting Agent) -----	5.0%
	<hr/>
	100.0%

\*Equivalent to 22% 2,2 bis (p-methoxyphenyl)—1,1,1—trichloroethane and 3% other isomers and reaction products.

### CAUTION:

Keep Out of Reach of Children  
See Other Warnings on Back Panel.

CHEMICAL FORMULATORS, Inc.

P. O. BOX 26 — VISCOSE ROAD  
NITRO, WEST VIRGINIA 25143

Nov. 15, 1971

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 451-43. SUBJECT  
TO ATTACHED COMMENTS.

### DIRECTIONS

#### FORAGE CROPS:

On Alfalfa, Clover, Grass, Soybeans apply Chemform Methoxychlor 2 E. C. at the rate of 2-3 quarts (1 to 1½ lbs. of active) per acre for the control of Alfalfa Weevil (Larvae), Alfalfa Webworm, Fall Armyworm, Clover Leaf Aphid, Pea Weevil, Soybean Caterpillar and Velvet Bean Caterpillar.

(May be used in conjunction with Malathion for Alfalfa Weevil control)

Mix with water (both high and low volume spraying) to give correct coverage per acre. Start applications at first sign of infestation and repeat at 7-10 day intervals as needed for control. Do not apply later than 7 days before harvest or grazing.

#### LIVESTOCK:

For control of Hornflies mix 1 quart in 12½ gallons of water and apply two (2) quarts of spray to each mature animal. For control of lice apply as drenching spray.

For Control of Hog Lice, mix 1 quart in 12½ gallons of water and apply as a drenching spray.

**DO NOT USE ON DAIRY ANIMALS OR IN DAIRY BARNs. DO NOT USE IN DIPPING VATS.**

#### SHADE TREES, ORNAMENTALS:

For control of Japanese Beetles, Leaf Hoppers, Tent Caterpillars, Cankerworms and Rose Chafer apply Methoxychlor 2 E.C. at the rate of 2-3 quarts (1-1.5 lbs. Actual) in 100 gallons of water (3-5 tablespoons in 1 gallon of water) Spray foliage thoroughly. Repeat applications every 7-14 days.

For ELM EARWORM BEETLE (Assists in control of Dutch Elm Disease) apply at the rate of 4-6 quarts per 100 gallons of water. Spray trunk and limbs thoroughly. Repeat every 7-14 days thru growing season.

CAUTION: Avoid contact with eyes, skin or clothing.

Avoid prolonged or repeated breathing of spray mist.

Avoid contamination of feed or foodstuffs.

Harmful if swallowed.

This product is toxic to fish. Keep out of lakes, streams and ponds.

Apply this product only as specified on this label. Do not apply when

weather conditions favor drift of spray from areas treated. Rinse

equipment and containers and dispose of wastes by burying in non-

crop lands away from water supplies. Containers should be disposed

by punching holes in them and burying with wastes, or by burning

(Keep out of the smoke).

#### NOTICE:

Do not mix this material with other materials not compatible with oil.

Observe precaution when applying oil-based insecticides to crops dur-

ing high temperatures since some crops are then more susceptible to

injury. Do Not apply within 14 days of Sulfur or Sulfur containing

sprays. Consult your county agent for proper dosages when applying

by aircraft