



READ THE LABEL

DURSBAN[®] M INSECTICIDE

Controls Numerous Pests of Households, Turf, and Ornamental Plants.

Also Recommended for Mosquito Control and Area Control of Ticks and Chiggers.

To Be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, Mosquito Abatement Districts, and Other Trained Personnel Responsible for Insect Control Programs.

chlorothioate] 41.2%
 29.5%
 29.3%

E.P.A. Est. 464-MI-1

ICAL
Is, or Clothing

s, no use este producto
nte.
English, do not use this

IREN
BE ABSORBED
EYES AND SKIN

Wash thoroughly after
• Flush contaminated
tion • Avoid breathing
Keep away from food,
inner closed • DO NOT
• Handle concentrate

symptomatically. Atropine is

pests indicated in the areas
 ty. Use the 0.25% spray to
 y infestations or as directed.
 ole base oil[†] as shown in the
 |.
 applied in food handling

Note: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrins may be added to the spray mixture containing DURSBAN M insecticide. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays; use pyrethrins at 0.25% in water-base and 0.05% in oil-base sprays. Consult the labeling for the dichlorvos or pyrethrins product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products. Do not use DURSBAN M in combination with dichlorvos (DDVP) or pyrethrins for treatment of food areas of food handling establishments.

Application Within Food Handling Establishments (Places Other Than Private Residences in Which Food is Held, Processed Prepared or Served) Including, But Not Limited to, Restaurants, Grocery Stores, Bakeries, Bottling Plants, Canneries and Grain Mills: Use as a spot treatment to control cockroaches (including strains resistant to certain other insecticides), ants, crickets, firebrats, silverfish and spiders in and around both food and nonfood areas by application as a coarse, low pressure (20 psi or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated using special care to avoid unnecessary runoff. Do not introduce the spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners of rooms and closets; along and behind baseboards; beneath and behind stoves, refrigeration units and similar food preparation areas and equipment; floor drains; and around plumbing and other utility installations. For ants apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and bases or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a pin stream of insecticide should be used. Repeat treatment as needed but not more often than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back), DURSBAN M may be applied after 2 days from last treatment but emergency use should be limited to once per month.

Application of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

Application Within Residential Buildings Including Homes and Apartment Buildings and Within Nonfood Areas of Industrial, Institutional, and Commercial Buildings Including Hospitals, Stores, Manufacturing Plants and Warehouses: For Spot Treatment Only — Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets; cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators and cabinets; around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For clover mites use a 0.5% spray only. Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

Pests Out-of-Doors

General Information: Use DURSBAN M for control of certain pests of lawns and ornamental plants, for area control of ticks and chiggers and for mosquito control. Consult the following directions for dosage and application details to control the pests listed.

Pests of Lawns and other Ornamental and Recreational Turf Grass Areas: Use DURSBAN M to control the pests listed in the following table by application at the recommended dosages and in accordance with the directions given below. Dilute DURSBAN M in enough water to obtain complete and uniform coverage of pest infested areas and apply as a coarse, low-pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf, except as noted. For best results, the lawn should be moist at time of treatment.

DOSAGES and DIRECTIONS for USE on TURF

Pest	Dosage DURSBAN M per 1000 sq. ft.	Remarks
Ants Armyworms Brown dog tick Chiggers Chinch bugs Clover mite Crickets Cutworms Earwigs Fleas Grasshoppers Sod webworms	¾† fl. oz.	Spray when pests first appear; retreat when needed. For sod webworms delay watering or mowing the treated area for 12 to 24 hours after treatment.
Turfgrass (Hyperodes) weevil	1½ fl. oz.	Spray suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service specialist for your area.
White grubs (European chafer and Japanese beetle larvae only)	1½ to 3 fl. oz.	For white grubs spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying, irrigate

DURSBAN M to Use



SPECIMEN LABEL
REDUCED TO 73%

9

4641

FORMULA 40*

HERBICIDE

CONTAINS ALKANOLAMINE SALTS† OF 2,4-D
Acid Equivalent: 4 Pounds per Gallon

For the Selective Control of Many Broadleaf Weeds in Non-Crop Areas, Grass Pastures, Rangelands and in Certain Crops. Also for Control of Trees by Injection

ACTIVE INGREDIENTS

Alkanolamine Salts (of the Ethanol and Iso-propanol series) of 2,4-Dichlorophenoxyacetic acid	59.7%
INERT INGREDIENTS	40.3%
2,4-Dichlorophenoxyacetic Acid Equivalent	38.6%
E.P.A. Registration No. 464-1-AA	E.P.A. Est. No. 464-MI 1

Salts are the least volatile form of 2,4-D and do not release enough vapors from treated areas to require special adjacent susceptible crops.

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, or Clothing

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION: (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

CAUTION

KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
CAUSES IRRITATION OF SKIN AND EYES
Do Not Get in Eyes, on Skin, or on Clothing

In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention; wash skin with soap and plenty of water. Remove and wash contaminated clothing before re-use. Do not wear contaminated shoes.

18.9 L / 5 GAL

86-1142 PRINTED IN U.S.A. IN OCTOBER, 1975
REPLACES SPECIMEN LABEL 86-1142 PRINTED IN SEPTEMBER, 1974
DISCARD PREVIOUS SPECIMEN LABELS.
REVISIONS INCLUDE: (1) USE ON APPALACHIAN ADDED; (2) USE ON STEWBERRIES ADDED; (3) PREPARATION APPLICATION OF CORN ADDED.

SPECIMEN LABEL
(BACK)
REDUCED TO 73%

Label Copy
FORMULA 40
October 31, 1975

Section B.3(c) - November 1975

Revisions to be made when
tolerances are established
and all registrants are
required to add limitations.

- Insert (A) Do not forage or feed treated corn fodder
for 7 days following application.
- Insert (B) Do not use in rice paddies where shellfish
are of economic importance or where flood
water is used for irrigation of other crops.
- Insert (C) See Section B.4 (b) for sugarcane.
- Insert (D) Do not graze dairy cattle in treated areas
for 14 days after application. Remove meat
animals from freshly treated areas for 7 days
before slaughter. Withdrawal is not needed
if 2 weeks or more has elapsed since the
treatment was applied. Do not cut treated
grass for hay within 30 days after application.
- Insert (E) On rangeland, apply at a rate of 6 quarts of
FORMULA 40 per acre per application, as a
spot treatment so the overall rate is no
more than 3 quarts per acre.

Lak
FOR
Aug
Pag

WE
and
pos
to
rec
ha
gra
rea
pro

NO

SU