

ACCEPTED
JAN 8 1973
UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT FOR ECONOMIC PESTS REGISTERED UNDER NO. 464-150 SUBJECT TO ATTACHED COMMENTS.



SPECIMEN LABEL

ESTERON*

76-E

HERBICIDE

Contains Isopropyl and Butyl Esters of 2,4-D • 2,4-D Acid Equivalent: 6 pounds per gallon

ACTIVE INGREDIENTS:	
2,4-Dichlorophenoxyacetic Acid, Butyl Esters	39.7%
2,4-Dichlorophenoxyacetic Acid, Isopropyl Ester	37.7%
INERT INGREDIENTS:	22.6%
2,4-D Acid Equivalent 63.5%—6 pounds per gallon	
E.P.A. Registration No. 464-150	

CAUTION
KEEP OUT OF REACH OF CHILDREN
Read Complete Precautions on Rear Panel

THE DOW CHEMICAL COMPANY

AND SUBSIDIARIES

MIDLAND, MICHIGAN 48640, USA ZURICH, SWITZERLAND HONG KONG, BCC
CORAL GABLES, FLORIDA 33134, USA SARNIA, ONTARIO, CANADA

* Trademark of THE DOW CHEMICAL COMPANY

86-1058 PRINTED IN U.S.A. IN NOVEMBER 1972

REPLACES SPECIMEN LABEL 86-1058 PRINTED IN NOVEMBER 1967
REVISIONS INCLUDE: ALL SECTIONS OF LABEL REVISED EXCEPT
COMPANY IDENTITY AND DISCLAIMER NOTICE.

SPECIMEN LABEL

ESTERON 76-E HERBICIDE

For the Control of Many Broadleaf Weeds, Herbaceous Perennials and Woody Plants Susceptible to 2,4-D in Grass Pastures, Certain Crops and Non-Crop Areas.

WEED LIST

ESTERON 76 E herbicide is recommended for control of numerous broadleaf weeds and certain 2,4-D susceptible woody plants without injury to most established grasses. Use in small grains, corn, sorghum, grass seed crops, pastures, rangeland and in non-crop areas. Species controlled include the following, plus many others:

bitterweed	galinsaga	pigweed	sumac
blueweed	goatsbeard	plantains	sunflower
broomweed	halogeton	radish, wild	sunflower
buckbrush	hemp, wild	ragweed	tanymustard
burdock	lechua	rosetem	tanysagwort
carpetweed	lambquarters	sage, coastal	thistle, bull
chicory	locust	sagebrush, sand	thistle, musk
coffeeweed	marshelder	sandshinnery oak	thistle, Russian
craton	milkvetch	shepherdspurse	tumbleweed
dandelion	morningglory, annual	smartweed	witch
docks	mustards	sneezewood, bitter	willow
elderberry	nettles	sawthistle, annual	witchweed
fanweed	pepperweed, field	spanishneedles	

USE DIRECTIONS

Apply ESTERON 76 E as water or oil spray during warm weather when weeds or brush are actively growing. Application under drought conditions often will give poor results. Use low spray pressure to minimize spray drift. On cropland and along roadsides, do not exceed 20 psi pressure. Apply enough spray volume to provide uniform coverage of weeds and brush, usually 5 to 20 gallons per acre by ground equipment and 3 to 5 gallons by aircraft. Higher gallonage may be used if desired to improve spray coverage and to reduce the hazard from spray drift. Generally the lower dosages recommended on this label will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult the higher dosages will be needed. For crop uses, do not mix with oil or other adjuvants unless specifically recommended on this label. Deep rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for maximum control. Do not

apply ESTERON 76 E where spray drift may contact nearby susceptible crops or other desirable plants or may contaminate water for irrigation or domestic use. Do not apply in the vicinity of 2,4-D sensitive crops or ornamental plants since vapors from this product may cause injury to such crops or plants. Read and follow all Use Precautions given on this label.

NOTE: If there are uncertainties concerning special local use situations or specific crop variety tolerances to 2,4-D, consult your State Agriculture Experiment Station or Extension Service Weed Specialists for advice.

TO PREPARE THE SPRAY: (1) Fill the spray tank about half full with water, then add the required amount of ESTERON 76 E, with agitation, and finally the rest of the water. **NOTE:** ESTERON 76 E in water forms an emulsion which tends to separate unless the mixture is kept agitated. (2) If oil is added, first mix the ESTERON 76 E and the oil and then add this mixture to the water with agitation. However, with adequate agitation, the oil can be added after the ESTERON 76 E is mixed in the water if strong agitation is provided. (3) If straight oil is used, a solution is formed and separation does not occur. Do not allow any water to get into the oil herbicide mixture to avoid formation of an invert emulsion.

WEED CONTROL IN SMALL GRAINS NOT UNDERSEEDED WITH A LEGUME: Water is recommended to make up the spray. If oil is used, there is greater risk of crop injury and of spray drift. Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 2 weeks after treatment.

Spring Wheat and Barley: Apply 1 1/2 to 2 1/2 pint per acre by air or ground equipment. A 2 pint per acre rate of ESTERON 76 E is an average dosage, effective on many weeds. Spray when grain is in full tiller stage (usually 4 to 8 inches tall) but before the boot stage and when weeds are small. Do not apply before the tiller stage nor from early boot to the dough stage. Use of higher rates (up to 1 1/2 pints per acre) may be required to control certain weeds but crop injury may result.

Winter Wheat and Rye: Apply 1 1/2 to 2 1/2 pint per acre in the spring at the full tiller stage but before the early boot stage. See more complete use directions under Spring Wheat and Barley.

Spring Seeded Oats: Apply 1 1/2 pint per acre at the full tiller stage but before the early boot stage. Oats are less tolerant to 2,4-D than wheat or barley and are more likely to suffer some injury, especially if higher rates (1 1/2 to 2 1/2 pint) are used to control difficult weeds.

Preharvest Treatment: Apply 2 1/2 to 1 1/2 pints per acre when grains are in the hard dough stage to control large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to cause succulent weed growth. **NOTE:** Do not feed treated straw to livestock.

WEED CONTROL IN CORN: Use one of the following three programs. **Preemergence:** Apply 1 1/2 to 2 1/2 pints per acre to soil anytime after planting but before corn emerges. Do not use on light sandy soil. **Emergence:** Apply 2 1/2 pint per acre just as corn plants are breaking ground. **Postemergence:** After emergence of corn, use 1/2 pint per acre. Application of 1/2 to 2 1/2 pint per acre may be needed for maximum control of some weeds but such rates are more likely to injure the corn. If corn is over 8 inches tall, use drop nozzles to keep the spray off the corn foliage as much as possible. Do not apply from the tasseling to dough stage. Do not use with oil, atrazine or adjuvants. Crop injury is more likely to occur if corn is growing rapidly under high temperature and high soil moisture conditions. To reduce breakage of stalks from temporary brittleness caused by 2 1/2 pint per acre, delay cultivation for 8 to 10 days after treatment. **NOTE:** Hybrids vary in response to 2,4-D and some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Contact seed company and Extension Service Weed Specialists for this information.

WEED CONTROL IN SORGHUM (MILO): Apply 1 1/2 pint per acre when sorghum is 5 to 15 inches tall. A higher rate of 1/2 to 2 1/2 pint per acre may be needed to control some weeds but the chance for crop injury is likewise increased. Do not use with oil or other adjuvants. Do not treat before the sorghum is 5 inches tall nor during the boot, flowering or early dough

stages. If sorghum is taller than 8 inches, use drop nozzles to keep the spray off the foliage as much as possible. Temporary crop injury may occur under conditions of high soil moisture and high air temperatures. Varieties vary in tolerance to 2,4-D and some hybrids are quite sensitive. Spray only varieties known to be tolerant to 2,4-D. Contact seed company and Extension Service Weed Specialists for this information. Do not apply in the vicinity of cotton, soybeans or other 2,4-D susceptible plants.

WEED CONTROL IN GRASS SEED CROPS: Use 3/4 to 1 pint per acre in the amount of water required for uniform application by air or ground equipment. Apply to established stands in spring from the tiller to early boot stage. Do not spray in boot stage. New spring seedlings may be treated with the lower rate after the grasses have at least five leaves. Perennial weed regrowth may be treated in the fall.

WEED AND BRUSH CONTROL IN RANGELAND AND GRASS PASTURES BY AIR OR GROUND EQUIPMENT: **NOTE:** Do not graze dairy animals on treated areas within 7 days after application. Do not apply to bent or established grass pastures. Do not apply to newly seeded areas until the grass is well established. Do not apply to grass in the boot to milk stage where grass seed production is desired. The following treatments will injure or kill legumes so use only where loss of legumes can be tolerated.

Broadleaf Weeds: To control bitterweed, broomweed, craton, docks, Kochia, marshelder, muskthistle and others, use 2 1/2 to 3 pints of ESTERON 76 E per acre in the amount of water needed for uniform application. If the weeds are young and growing actively 1 1/2 to 2 pints per acre will provide control of many species. Deep rooted perennial weeds may require repeated treatments in the same year or in subsequent years.

Chaparral Brush Species: To control chamise, manzanita, buckbrush, coastal sage and certain other chaparral species, use 1 1/2 to 2 quarts per acre in 8 to 10 gallons of water. A gallon of oil per acre may be included in the spray mixture for added effectiveness. For effective control, the brush must be fully leafed out and growing actively when sprayed. Retreatment may be needed.

Big Sagebrush: Use 1 1/2 to 2 quarts per acre in 2 to 3 gallons of oil or in 3 to 5 gallons of oil-water emulsion spray. For effective control the sagebrush should be in full foliage and growing actively when sprayed.

WEED CONTROL IN NON-CROP AREAS SUCH AS LAWNS, AIRFIELDS, ROADSIDES, VACANT LOTS, DRAINAGE DITCH BANKS: Apply 1 to 2 quarts of ESTERON 76 E per acre in the amount of water needed for uniform application. Usually 1 1/2 quarts per acre provides good weed control under average conditions. Treat when weeds are young and growing well. Do not use on gall greens nor on dichandra or other broadleaf herbaceous ground covers. Do not use on creeping grasses such as bent and St. Augustine except for spot treating, nor on newly seeded turf until grass is well established. Legumes are usually damaged or killed so do not treat areas where the legumes are desired. Deep rooted perennial weeds may require repeated treatments in the same season or in subsequent years.

WOODY PLANT CONTROL IN NON-CROP AREAS: To control species susceptible to 2,4-D in right of ways, fence rows, roadsides, and along drainage ditchbanks, spray brush up to 5 to 8 feet tall after spring foliage is well developed, using 2 to 3 quarts of ESTERON 76 E in 100 gallons of water and wetting all parts of the brush including foliage, stems and bark. This may require up to 400 gallons of spray per acre for adequate coverage of solid stands of brush. Make application in such a way as to prevent drift of the spray off the area being treated. Spraying can be effective at any time up to 3 weeks before frost as long as soil moisture is sufficient for active growth of the brush. Less effective control may be obtained during hot dry weather when soil moisture is deficient and plants are not actively growing. Oil and wetting agent may be added to the spray, if needed for increased effectiveness. For more resistant species and for general control of mixed brush use ESTERON Brush Killer or ESTERON 245 herbicide.

SPOT TREATMENT: To control broadleaf weeds in small non-cropland areas with a hand sprayer, use 1/2 pint of ESTERON 76 E in 3 gallons of water and spray to thoroughly wet all weed foliage. Keep spray mixture agitated to prevent separation.

USE PRECAUTIONS

Do not apply ESTERON 76 E herbicide directly to, or otherwise permit it to come into contact with cotton, grapes, fruit trees, vegetables, flowers or other desirable crops or ornamental plants which are sensitive to 2,4-D herbicide. Do not permit spray mix to drift onto them since even very small quantities of the spray, which may not be visible, can cause severe injury during both growing and dormant periods. Use coarse sprays to minimize drift. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible. By applying 20 gallons or more of spray per acre, by using no more than 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips, by spraying when wind velocity is low, and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone type insecticide or other nozzles that produce a fine droplet spray. With aircraft application, drift can be reduced by applying no less than 5 gallons of spray per acre, by using no more than 20 pounds spray pressure at the nozzles, by using nozzles which produce a coarse spray pattern, and by spraying only when the wind velocity is less than 5 miles per hour.

Applications by aircraft, ground rig and hand dispenser should be carried out only when there is no hazard from spray drift. Do not apply in the vicinity of cotton, grapes, tomatoes or other desirable 2,4-D susceptible crop or ornamental vegetation. Do not spray when the wind is blowing toward susceptible crops or ornamental plants.

This is a high-volatile 2,4-D ester formulation. Vapors from this product may injure susceptible plants growing nearby. Do not use in or near a greenhouse. Excessive amount of this herbicide in the soil may temporarily inhibit seed germination or plant growth.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where it is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. Do not contaminate irrigation ditches or water used for irrigation or domestic purposes.

To avoid injury to desirable plants, do not handle or apply other agricultural chemicals with the same equipment used for ESTERON 76 E, except as specified on this label. The product can be stored in an unheated building but do not store near herbicides, weed insecticides, or fungicides. If exposed to a freezing temperature, it should be warmed to at least 40 F and mixed thoroughly before using. Do not reuse containers. Dispose of empty containers by punching holes in them and burying with waste in a non-cropland area. Do not water supplies or follow official local recommendations for safe disposal.

Local conditions may affect the use of herbicide. Contact your State Agriculture Experiment Station or Extension Service weed specialists for advice in making treatments that are suited to best fit local conditions. Be sure that use of this product conforms to all applicable regulations. Apply this product only as specified on this label.

CAUTION
KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
MAY CAUSE IRRITATION
Avoid Contact with Eyes, Skin and Clothing

NOTICE: While the use of this product is restricted to certain uses, it is not intended for use in the control of weeds in the vicinity of cotton, grapes, fruit trees, vegetables, flowers or other desirable crops or ornamental plants which are sensitive to 2,4-D herbicide. Do not permit spray mix to drift onto them since even very small quantities of the spray, which may not be visible, can cause severe injury during both growing and dormant periods. Use coarse sprays to minimize drift. With ground equipment, spray drift can be lessened by keeping the spray boom as low as possible. By applying 20 gallons or more of spray per acre, by using no more than 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips, by spraying when wind velocity is low, and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone type insecticide or other nozzles that produce a fine droplet spray. With aircraft application, drift can be reduced by applying no less than 5 gallons of spray per acre, by using no more than 20 pounds spray pressure at the nozzles, by using nozzles which produce a coarse spray pattern, and by spraying only when the wind velocity is less than 5 miles per hour.