

ACCEPTED  
1-24-72  
464-272

# TIPPON T6

## BRUSH AND WEED KILLER

**SPECIMEN LABEL**

Containing 9.2 Pounds per Gallon of Powerful, Low Volatility Esters of 2,4,5-T • 2,4,5-T Acid Equivalent 6 Pounds per Gallon  
FOR THE CONTROL OF TREES, BRUSH AND BROAD-LEAVED WEEDS

**Active Ingredient:**

2,4,5-Trichlorophenoxyacetic Acid, Propylene Glycol Butyl Ether Esters ----- 91.6%  
2,4,5-Trichlorophenoxyacetic Acid Equivalent 59.7%

Inert Ingredients ----- 8.4%

E.P.A. Registration No. 464-272

**DIRECTIONS**

Tippon T6 is recommended for use in controlling herbaceous and woody plants growing in rangelands, pastures, fence rows, farmyards and right-of-ways, including certain 2,4-D resistant plants such as ash, brambles, ground cherry, hawthorn, horse nettle, maple, mesquite, oak, osage orange, palmetto, poison ivy, prickly pear cactus, salmonberry, wild blackberry, wild rose, and certain species of Ribes.

**PREPARING THE SPRAY:** Use only diesel oil, No. 1 or No. 2 fuel oil, or kerosene. Tippon T6 herbicide is a non-emulsifiable formulation to be used in oil only. Add Tippon T6 to the required amount of oil in the spray tank or mixing tank and mix thoroughly.

**BASAL BARK TREATMENT:** Brush and small trees can be controlled by spraying the basal parts of brush stems and tree trunks to a height of 12 to 15 inches from the ground line. Use a solution of 2 gallons of Tippon T6 in 148 gallons (1 pint in 9 gallons) of oil. With certain species, 4 gallons of Tippon T6 in 145 gallons of oil (1 pint in 4½ gallons) of oil is effective. Knapsack or power equipment may be used, but complete wetting of the indicated area is necessary, particularly at the ground line. This means spraying until run-down or run-off to the ground line is noticeable. Old or rough bark requires more spray volume than young or smooth bark. Low pressures are desirable. Apply at any time, including the winter months, except when snow or water prevents spraying to the ground line. Often delayed response and killing can be expected.

**DORMANT CANE BROADCAST:** Treat any time after brush is dormant and most of the foliage has dropped. Sprays should be concentrated at the base of stems and in addition, the upper parts of the stems should be broadcast sprayed enough to wet them. Under root suckering species such as sumac, persimmon, sassafras and locust, also spray

the group area to control small root suckers that may not be visible. Mix 1 gallon of Tippon T6 in 100 gallons of fuel oil. Brush of average density and 4-6 feet high may take up to 150 gallons of spray mixture per acre.

**STUMP TREATMENT:** Where growth is more than 6 to 8 feet tall, cut it close to the ground and spray the freshly cut stumps and stubs with 2 gallons of Tippon T6 in 148 gallons (or 1 pint in 9 gallons) of oil, mixed thoroughly. For stumps and stubs of resistant species 4 gallons of Tippon T6 in 146 gallons (1 pint in 4½ gallons) of oil has been effective. Wet thoroughly all exposed bark, as well as cut surfaces. This means spraying until run-down or run-off to the ground line is noticeable. Old or rough bark requires more spray volume than young or smooth bark. Apply at any time, including the winter months, except when snow or water prevents spraying to the ground line. Best results are obtained on freshly cut stumps two inches across or larger.

**"FRILL" TREATMENT:** For large trees, make a single-hack girdle or "frill" of overlapping axe cuts completely around the tree as close to the ground as possible. Treat the injured area with a mixture of 2 gallons of Tippon T6 in 148 gallons (½ pint in 4½ gallons) of oil.

**AIRPLANE APPLICATION:** To control mesquite, use ½ to ¾ pint of Tippon T6 in 4 gallons of oil per acre. Apply 50 to 80 days after first leaves appear. Do not treat if drought has prevented good foliage growth. For post oak and black oak, use 2½ pints of Tippon T6 in 4 to 6 gallons of oil per acre. Read "Preparing the Spray" before mixing. Apply after foliage is fully developed. Repeat treatment should be applied in accordance with local recommendations. NOTE: Soil moisture should be adequate for normal growth.

**USE PRECAUTIONS**

Do not apply Tippon T6 directly to, or otherwise permit it to come into direct contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, black locust or other desirable plants which are sensitive to 2,4,5-T and do not permit spray mists containing it to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. (Coarse sprays are less likely to drift.) Accordingly, applications

by airplane, ground rigs, and hand dispensers should be carried out only when there is no hazard from drift. Do not apply by airplane in the vicinity of cotton, grapes, or other desirable 2,4,5-T susceptible vegetation. At higher temperatures vaporization may cause injury to susceptible plants growing nearby.

Do not use on creeping grasses, such as bent, except for spot spraying, nor on freshly seeded turf until grass has become well established. (Most legumes are usually damaged or killed.)

Do not contaminate irrigation ditches or water used for domestic purposes. Do not store near fertilizers, seeds, insecticides, or fungicides. To avoid injury to desirable plants, do not store, handle, or apply other agricultural chemicals with the same containers or equipment used with Tippon T6.

Do not graze dairy animals on treated areas within 6 weeks after application. Do not slaughter meat animals grazing on treated areas within 2 weeks after application.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Apply only as specified on this label. Be sure that use of this product conforms to all applicable regulations.

Rinse equipment and containers and dispose of waste by burying in non-croplands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not use around the home or recreation areas, or similar sites.

Local conditions may affect the use of herbicides. State experimental station or extension weed specialists in many states issue recommendations to fit local conditions.

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN  
MAY BE HARMFUL IF SWALLOWED  
MAY CAUSE SKIN IRRITATION  
Avoid Contact with Eyes, Skin and Clothing**

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonable fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**THE DOW CHEMICAL COMPANY  
MIDLAND, MICHIGAN 48640**