



ESTERON* 76 BE

HERBICIDE

Contains Butyl Esters of 2,4-D

*For the Control of Many Broadleaf Weeds, Herbaceous Perennials and Woody Plants
Susceptible to 2,4-D in Grass Pastures, Certain Crops and Non-Crop Areas.*

ACTIVE INGREDIENT:
 2,4-Dichlorophenoxyacetic Acid Butyl Esters† 75.2%
 INERT INGREDIENTS 24.8%
 2,4-Dichlorophenoxyacetic Acid Equivalent†; 90.0% → 5.64 lb/gal
 †Heater Specific by AOAC Method No. 8.D01-8
 EPA Reg. No. 484-279
 EPA Est. 484-MI-1

PRECAUCION AL USUARIO: Si usted no lee Ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
 TRANSLATION: (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

HARMFUL IF SWALLOWED • MAY CAUSE IRRITATION
 Avoid Contact with Eyes, Skin and Clothing
 Do Not Cut or Weld Container

In case of an emergency endangering life or property involving this product, call collect 817-638-4400.

AGRICULTURAL CHEMICAL
 Do Not Ship or Store with Food, Feeds, Drugs or Clothing

18.93 L / 5 GAL



TERON 76 BE

HERBICIDE

Contains Butyl Esters of 2,4-D

Effective on Many Broadleaf Weeds, Herbaceous Perennials and Woody Plants
Equivalent to 2,4-D in Grass Pastures, Certain Crops and Non-Crop Areas.

Acetic Acid Butyl Esters†75.2%
Acetic Acid24.8%
Acetic Acid Equivalent†: 60.0% — 5.84 lb/gal
EPA Method No. 8.D01-5

EPA Est. 464-MI-1

NOTE: If used on lawns, no use este producto
ya sido explicado anteriormente.
USER: If you cannot read English, do not use this
product (has been fully explained to you.)

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

HARMFUL IF SWALLOWED • MAY CAUSE IRRITATION
Avoid Contact with Eyes, Skin and Clothing
Do Not Cut or Weld Container

In case of an emergency endangering life or
property involving this product, call collect
517-636-4400

AGRICULTURAL CHEMICAL
Do Not Store or Share with Food, Feed,
Drugs or Clothing

18.93 L / 5 GAL



ESTERON 76BE

HERBICIDE

SP

USE DIRECTIONS

ESTERON 76BE is a selective herbicide for the control of weeds in corn, sorghum, and small grain crops. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants. The herbicide is applied as a spray or granules, depending on the crop and weed type. It is important to follow the application rates and timing instructions carefully to ensure effective weed control while minimizing damage to the crop.

Do not apply ESTERON 76BE where spray drift may contact nearby susceptible crops or other desirable plants. Do not apply in the vicinity of 2,4-D sensitive crops or ornamental plants since vapors from this product may cause injury to such crops or plants. Read and follow all Use Precautions given on this label.

NOTE: Do not apply ESTERON 76BE where spray drift may contact nearby susceptible crops or other desirable plants. Do not apply in the vicinity of 2,4-D sensitive crops or ornamental plants since vapors from this product may cause injury to such crops or plants. Read and follow all Use Precautions given on this label.

TO PREPARE THE SPRAY: Add the required amount of ESTERON 76BE to the spray tank. Add water to the tank to the required volume. Mix thoroughly. Apply immediately.

WEED CONTROL IN SMALL GRAINS NOT UNDERSEEDED WITH A LEGUME: NOTE: Do not apply ESTERON 76BE to small grains not underseeded with a legume. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Spring Wheat and Barley: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the crop is in the early stages of growth.

Winter Wheat and Barley: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the crop is in the early stages of growth.

Spring Seeded Oats: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the crop is in the early stages of growth.

Preharvest Treatment: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the crop is in the early stages of growth.

WEED CONTROL IN CORN

Preemergence: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply before the corn is planted. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Postemergence: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the corn is in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

WEED CONTROL IN SORGHUM: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the sorghum is in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

WEED CONTROL IN GRASS SEED CROPS: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the grass seed crop is in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

WEED AND BRUSH CONTROL IN PASTELAND AND GRASS PASTURES BY AIR OR GROUND EQUIPMENT: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the pasture is in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Broodleaf Weeds: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the broodleaf weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Chaparral Brush Species: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the chaparral brush species are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Big Sagebrush: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the big sagebrush is in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

WEED CONTROL IN NON-CROP AREAS SUCH AS LAWNS, AIRFIELDS, ROADSIDES, YACANT LOTS, DRAINAGE DITCH BANKS: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the non-crop areas are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

WOODY PLANT CONTROL IN NON-CROP AREAS: Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the woody plants are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

ESTERON 76BE is a selective herbicide for the control of weeds in corn, sorghum, and small grain crops. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants. The herbicide is applied as a spray or granules, depending on the crop and weed type. It is important to follow the application rates and timing instructions carefully to ensure effective weed control while minimizing damage to the crop.

Do not apply ESTERON 76BE where spray drift may contact nearby susceptible crops or other desirable plants. Do not apply in the vicinity of 2,4-D sensitive crops or ornamental plants since vapors from this product may cause injury to such crops or plants. Read and follow all Use Precautions given on this label.

SPOT TREATMENT: To control weeds in small areas, apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Application by aircraft: ground rig. there is no hazard from spray drift. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

This is a high volatile 2,4-D ester: susceptible plants growing nearby. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

The product is toxic to fish: Keep out of cleaning of equipment or disposal. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Toxic to fish: susceptible plants. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Use with caution: Do not apply ESTERON 76BE to small grains not underseeded with a legume. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Do not apply: Do not apply ESTERON 76BE to small grains not underseeded with a legume. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Do not apply: Do not apply ESTERON 76BE to small grains not underseeded with a legume. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Do not apply: Do not apply ESTERON 76BE to small grains not underseeded with a legume. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Do not apply: Do not apply ESTERON 76BE to small grains not underseeded with a legume. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

Do not apply: Do not apply ESTERON 76BE to small grains not underseeded with a legume. Apply ESTERON 76BE at the rate of 1.0 to 1.5 lb. a.i. per acre. Apply when the weeds are in the early stages of growth. It is effective against a wide range of weeds, including broadleaf weeds, grasses, and woody plants.

THE DOW CHEMICAL COMPANY
AND SUBSIDIARIES
MIDLAND, MICHIGAN 48640, USA * BIRKENHEAD, ENGLAND * HONG KONG
CORAL GABLES, FLORIDA 33134, USA * SARASOTA, ONTARIO, CANADA
* Trademark of THE DOW CHEMICAL COMPANY

1984



IRON 76 BE

HERBICIDE

SPECIMEN LABEL
(BACK)
REDUCED TO 71%

WEED CONTROL IN CORN

Preemergence: Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the corn plants. Apply before the corn plants are 4 inches high. **Postemergence:** Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the corn plants. Apply before the corn plants are 4 inches high. **Emergence:** Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the corn plants. Apply before the corn plants are 4 inches high.

WEED CONTROL IN SORGHUM MILO

Preemergence: Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the sorghum milo plants. Apply before the sorghum milo plants are 4 inches high. **Postemergence:** Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the sorghum milo plants. Apply before the sorghum milo plants are 4 inches high. **Emergence:** Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the sorghum milo plants. Apply before the sorghum milo plants are 4 inches high.

WEED CONTROL IN GRASS SEED CROPS

Preemergence: Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the grass seed crops. Apply before the grass seed crops are 4 inches high. **Postemergence:** Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the grass seed crops. Apply before the grass seed crops are 4 inches high. **Emergence:** Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the grass seed crops. Apply before the grass seed crops are 4 inches high.

WEED AND BRUSH CONTROL IN RANGELAND AND GRASS PASTURES BY AIR OR GROUND EQUIPMENT

NOTE: Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the rangeland and grass pastures. Apply before the rangeland and grass pastures are 4 inches high.

Broadleaf Weeds

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the broadleaf weeds. Apply before the broadleaf weeds are 4 inches high.

Chopwood Brush Species

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the chopwood brush species. Apply before the chopwood brush species are 4 inches high.

Big Sagebrush

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the big sagebrush. Apply before the big sagebrush are 4 inches high.

WEED CONTROL IN NON CROP AREAS SUCH AS LAWNS, AIRFIELDS, ROADSIDES, VACANT LOTS, DRAINAGE DITCH BANKS

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the non crop areas. Apply before the non crop areas are 4 inches high.

WOODY PLANT CONTROL IN NON CROP AREAS

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the woody plants. Apply before the woody plants are 4 inches high.

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the weeds. Apply before the weeds are 4 inches high.

SPOT TREATMENT

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the weeds. Apply before the weeds are 4 inches high.

USE PRECAUTIONS

Do not apply IRON 76 BE to any crop or other desirable vegetation. Do not apply IRON 76 BE to any crop or other desirable vegetation. Do not apply IRON 76 BE to any crop or other desirable vegetation.

Application by aircraft, ground rig and hand dispenser should be carried out only when there is no hazard from spray drift. Do not apply in the vicinity of cotton, grapes, tomatoes or other desirable 2,4-D susceptible crop or ornamental vegetation. Do not spray when the wind is blowing toward susceptible crops or ornamental plants.

This is a high-volatile 2,4-D ester formulation. Vapors from this product may injure susceptible plants growing nearby. Do not use in a greenhouse. Exercise caution in the use of this product in the vicinity of susceptible plants.

This product is toxic to fish, keep out of lakes, streams and ponds. Do not contaminate water by cleaning equipment and spillage. Do not contaminate irrigation ditches or water used for irrigation purposes.

This product is highly flammable. Do not use near open flames, heat, or other sources of ignition. Do not use in the vicinity of electrical equipment. Do not use in the vicinity of electrical equipment.

Do not use in the vicinity of electrical equipment. Do not use in the vicinity of electrical equipment. Do not use in the vicinity of electrical equipment.

Be sure that use of this product conforms to all applicable regulations. Apply this product only as specified on this label.

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the weeds. Apply before the weeds are 4 inches high.

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the weeds. Apply before the weeds are 4 inches high.

Apply 1.5 to 2.0 lb. of IRON 76 BE per acre to the soil in the row between the weeds. Apply before the weeds are 4 inches high.

THE DOW CHEMICAL COMPANY

AND SUBSIDIARIES

MIDLAND MICHIGAN 48640 USA • MORGEN SWITZERLAND • HONG KONG
CORAL GABLES, FLORIDA 33134, USA • SARNIA ONTARIO CANADA

Trademark of THE DOW CHEMICAL COMPANY