



SPECIMEN LABEL

KORLAN* 5G

INSECTICIDE GRANULES

CONTROLLED BY THE U.S. DEPARTMENT OF AGRICULTURE
* CHICKEN BODY LICE ON POULTRY

USE PRECAUTIONS

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, or Clothing

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.

LIVESTOCK REMEDY - NOT FOR HUMAN USE

DIRECTIONS FOR USE

CAUTION

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED**

Avoid contamination of food, feedstuffs, and water supplies.
Avoid skin contact • Wash thoroughly with soap and water after handling

NOTE TO PHYSICIAN OR VETERINARIAN

THE DOW CHEMICAL COMPANY

LB.

86-1257 PRINTED IN U.S.A. IN SEPTEMBER, 1975.

REPLACES SPECIMEN LABEL 86-1257 PRINTED IN NOVEMBER, 1974.
DISCARD PREVIOUS SPECIMEN LABELS.
REVISIONS INCLUDE: (1) REVISED USE PRECAUTIONS SECTION.

ACCEPTED
NOV 14 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 404-316



SPECIMEN LABEL

KORLAN* 5G

INSECTICIDE GRANULES

- CONTROLS:**
- HOG LICE on all ages of swine - Newborn too.
 - HOUSEFLY MAGGOTS (Larvae).
 - CHICKEN BODY LICE on POULTRY

Active Ingredient:

Ronnel (0,0-dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate)

5%
95%

Inert Ingredients:

Trademark of The Dow Chemical Company abroad
From technical grade - not suitable for internal use in animals

E.P.A. Registration No. 464-316-AA

E.P.A. Est. 464-MI-1

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, or Clothing

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION: (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

LIVESTOCK REMEDY - NOT FOR HUMAN USE

DIRECTIONS FOR USE

Distribute KORLAN 5G Insecticide Granules with a mechanical spreader, scoop or can

Control of Hog Lice: Apply KORLAN 5G Insecticide Granules evenly over the bedding for swine at the rate of 1/2 lb /100 ft² of area. No mixing with the bedding is necessary. One treatment will eliminate the hog lice infestation when the animals use the bedding regularly. Retreat if reinfestation occurs.

Control of Chicken Body Lice on Poultry: Scatter KORLAN 5G Insecticide Granules evenly over the litter at the rate of 1 lb./100 ft². KORLAN 5G may be used in broiler, layer, and replacement pullet houses. Retreat if reinfestation occurs.

Control of Housefly Maggots (Larvae): Caged-Layer Poultry Houses: Apply KORLAN 5G Insecticide Granules at the rate of 1 lb /100 ft² of area. Apply evenly over manure piles, under layer cages, and over the floor area where flies breed. Retreat as needed.

NOTE: If first applied when flies are numerous, it may take 2 weeks or more for a noticeable reduction in the number of adult flies. Quick kill of these flies may be accomplished by the use of a surface treatment with KORLAN* 24E Insecticide. (See KORLAN 24E product label for directions.)

USE PRECAUTIONS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this product.

CAUTION

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED**

**Avoid contamination of food, feedstuffs, and water supplies
Avoid skin contact • Wash thoroughly with
soap and water after handling**

In case of eye contact, flush eye with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN OR VETERINARIAN

This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

B775

THE DOW CHEMICAL COMPANY

AND SUBSIDIARIES

MIDLAND, MICHIGAN 48640, USA ZURICH, SWITZERLAND HONG KONG, B.C.C.
CORAL GABLES, FLORIDA 33134, USA SARNIA, ONTARIO, CANADA

* Trademark of THE DOW CHEMICAL COMPANY





KORLAN* 5G

INSECTICIDE GRANULES

- CONTROLS:**
- HOG LICE on all ages of swine - Newborn too.
 - HOUSEFLY MAGGOTS (Larvae).
 - CHICKEN BODY LICE on POULTRY

Active Ingredient:

Ronnel (0,0-dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate)

5%

Inert Ingredients

95%

Trademark of The Dow Chemical Company abroad
From technical grade, not suitable for internal use in animals

E.P.A. Registration No. 464-316-AA

E.P.A. Est. 464-MI-1

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, or Clothing

PRECAUCION AL USUARIO: Si usted no lee Inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION: (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

LIVESTOCK REMEDY — NOT FOR HUMAN USE

DIRECTIONS FOR USE

Distribute KORLAN 5G Insecticide Granules with a mechanical spreader, scoop or can

Control of Hog Lice: Apply KORLAN 5G Insecticide Granules evenly over the bedding for swine at the rate of 1/2 lb./100 ft² of area. No mixing with the bedding is necessary. One treatment will eliminate the hog lice infestation when the animals use the bedding regularly. Retreat if reinfestation occurs.

Control of Chicken Body Lice on Poultry: Scatter KORLAN 5G Insecticide Granules evenly over the litter at the rate of 1 lb./100 ft². KORLAN 5G may be used in broiler, layer, and replacement pullet houses. Retreat if reinfestation occurs.

Control of Housefly Maggots (Larvae): Caged-Layer Poultry Houses: Apply KORLAN 5G Insecticide Granules at the rate of 1 lb./100 ft² of area. Apply evenly over manure piles, under layer cages, and over the floor area where flies breed. Retreat as needed.

NOTE: If first applied when flies are numerous, it may take 2 weeks or more for a noticeable reduction in the number of adult flies. Quick kill of these flies may be accomplished by the use of a surface treatment with KORLAN* 24E Insecticide. (See KORLAN 24E product label for directions.)

USE PRECAUTIONS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this product.

CAUTION

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED**

**Avoid contamination of food, feedstuffs, and water supplies
Avoid skin contact • Wash thoroughly with
soap and water after handling**

In case of eye contact, flush eye with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN OR VETERINARIAN

This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

B775


THE DOW CHEMICAL COMPANY

AND SUBSIDIARIES

MIDLAND, MICHIGAN 48640, USA ZURICH, SWITZERLAND HONG KONG, BCC

CORAL GABLES, FLORIDA 33134, USA SARNIA, ONTARIO, CANADA

*Trademark of THE DOW CHEMICAL COMPANY





SPECIMEN LABEL

KORLAN* 5G

INSECTICIDE GRANULES

CONTROLS: • HOG LICE on all ages of swine - Newborn too.
• HOUSEFLY MAGGOTS (Larvae).
• CHICKEN BODY LICE on POULTRY

Active Ingredient

Ronnel (0,0-dimethyl 0-(2,4,5-trichlorophenyl) phosphorothioate)

5%

Inert Ingredients

95%

E.P.A. Registration No. 464-316-AA

E.P.A. Est. 464-MI-1

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, or Clothing

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION: (TO THE USER. If you cannot read English, do not use this product until the label has been fully explained to you.)

LIVESTOCK REMEDY — NOT FOR HUMAN USE

DIRECTIONS FOR USE

Control of Hog Lice Apply KORLAN 5G insecticide granules evenly over the body of the swine at the rate of 1 lb. per 100 lb. of area. No mixing with the bedding is necessary. Use the product weekly to maintain the hog free of lice. Retest when the animals use the product regularly. Retreat if infestation recurs.

Control of Chicken Body Lice on Poultry. Scatter KORLAN 5G insecticide granules evenly over the floor at the rate of 1 lb. per 100 lb. KORLAN 5G may be used in poultry houses, and in other types of houses. Retreat if infestation recurs.

Control of Housefly Maggots (Larvae) Caged-Layer Poultry Houses. Apply KORLAN 5G insecticide granules at the rate of 1 lb. per 100 lb. of area. Apply evenly over manure piles, under layer cages, and over the floor up to where flies breed. Retreat as needed.

NOTE: If first applied when flies are numerous, it may take 2 weeks to obtain a noticeable reduction in the number of adults. If a week of this treatment is not sufficient, repeat application. For complete treatment with KORLAN 5G, see the separate label directions.

USE PRECAUTIONS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where it may be a key to food. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this product.

CAUTION

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED**

**Avoid contamination of food, feedstuffs, and water supplies—
Avoid skin contact • Wash thoroughly with
soap and water after handling**

In case of eye contact, flush eye with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN OR VETERINARIAN

This material is a salicylate ester. Treat symptomatically. Atropine is an antidote.

NOTICE: Some warnings that the product is resistant to some insects and is was many times the purpose of this label. When used in accordance with the directions under this label, the user assumes the risk of the use of this product. **WARRANTY:** THE WARRANTY OF MERCHANTABILITY OF THIS PRODUCT IS LIMITED TO THE USE OF THIS PRODUCT IN ACCORDANCE WITH THE DIRECTIONS ON THE LABEL. THE USER ASSUMES THE RISK OF THE USE OF THIS PRODUCT.

B775

THE DOW CHEMICAL COMPANY

AND SUBSIDIARIES

MIDLAND MICHIGAN 48640 USA ZURICH SWITZERLAND HONG KONG BCC

CORAL GABLES FLORIDA 33134 USA SARNIA ONTARIO CANADA

*Trademark of THE DOW CHEMICAL COMPANY



86-1257 PRINTED IN U.S.A. IN SEPTEMBER, 1975.

REPLACES SPECIMEN LABEL 86-1257 PRINTED IN NOVEMBER, 1974.
DISCARD PREVIOUS SPECIMEN LABELS.
REVISIONS INCLUDE: (1) REVISED USE PRECAUTIONS SECTION.

