

PRODUCT ADVANTAGES

Makes a highly stable livestock dip in hard or soft water • Makes both spray and dip • Excellent wetting as spray or dip • Readily mixes with oil for back-rubber use • For livestock of all ages, newborn too! • This one really works

LIVESTOCK REMEDY, NOT FOR HUMAN USE

USE PRECAUTIONS

- Do not dip or spray sheep and goats within 28 days of slaughter.
- Do not spray beef cattle with 3/4% KORLAN 4E within 14 days of slaughter.
- Do not dip or spray beef cattle with 1/2% KORLAN 4E within 7 days of slaughter. Beef cattle may be dipped or sprayed with 1/8% or 1/4% any time prior to slaughter. Backrubber may be used any time on beef or dairy cattle.
- Do not use spray or dip on animals producing milk for human consumption. Dairy cows should be treated early in their dry period.
- Do not spray dairy animals with 3/4% of KORLAN 4E for 14 days prior to freshening. If dairy cows or heifers freshen within 14 days of the last spray treatment, milk must not be used for food for the required 14 days after the last spray treatment with 3/4% KORLAN 4E.
- Do not spray dairy animals with 1/4% or 1/2% KORLAN 4E for 7 days prior to freshening. If dairy cows or heifers freshen within 7 days of the last spray treatment, milk must not be used for food for the required 7 days after the last spray treatment with 1/4% or 1/2% KORLAN 4E.
- Remove lactating dairy animals from buildings while applying residual fly sprays.
- Do not allow insecticide to come in contact with food, feed or water supplies. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.
- Warning:** Ronnel is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides or chemicals.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

LOT MM

5 GAL

86-1193 PRINTED IN U.S.A. IN NOVEMBER, 1971.

REPLACES SPECIMEN LABEL 86-1193 PRINTED IN AUGUST, 1968.

REVISIONS INCLUDE: (1) TABLES REVISED TO INCLUDE USES FOR CONTROL OF SCREWORMS, COCKROACHES, AND MISC. INSECTS. (2) USE PRECAUTIONS REVISED TO REFLECT NEW TOLERANCES FOR RONNEL. (3) PRECAUTIONARY LABELING REVISED. (4) PRODUCT ADVANTAGES SECTION REVISED. (4) U.S.D.A. CHANGED TO E.P.A. (5) WORLD-WIDE COMPANY NAME USED.

| Problem, ear ticks and screwworms | 1/4% | 2 qt | 5 gal | 9 qt |
|---|------|-------|--------|-----------|
| Sheep, Goats, Lambs, and Kids: Lice, sheep keds, and hornflies | 1/4% | 2 qt | 5 gal | 9 qt |
| Wool maggots and screwworms | 1/2% | 1 gal | 10 gal | 4 1/2 gal |
| Hogs: Lice | 1/4% | 2 qt | 5 gal | 9 qt |

2. To maintain proper concentration recharge vat when 300 gallons of dip solution have been removed by animals.
3. To replace evaporation loss add water only to previous level of dip solution.
4. If flooding has occurred add KORLAN 4E at the recharge rate.
5. Retreat as necessary but not more often than every 2 weeks.

BACKRUBBER RECOMMENDATIONS

Cattle Backrubber — For hornflies. Aids in the control of lice if used continuously. Aids in the control of face flies if backrubber is placed so animals can rub face. Place where animals regularly travel or congregate.

Mix 1 gallon KORLAN 4E with 55 gallons diesel or mineral oil. Saturate fibre. Recharge as necessary. Do not mix with used motor oil.

SPRAY RECOMMENDATIONS

| PROBLEM | Ronnel (%) | Amount KORLAN 4E for | | | REMARKS | |
|--|------------------|----------------------------|--------------------|----------------------|---|--|
| | | 1 gal | 25 gal | 100 gal | | |
| LIVESTOCK INSECTS | | | | | | |
| Cattle and Calves: | | | | | | |
| Hornflies, ear ticks and screwworms | 1/2% | 1 1/3 oz | 2 pt | 1 gal | Dilute with water. Spray entire animal to wet thoroughly. Retreat as necessary but not more often than every 2 weeks. | |
| Lice | 1/4% | 2/3 oz | 1 pt | 2 qt | | |
| Lone Star and Winter Ticks | 3/4% | 2 oz | 3 pt | 1 1/2 gal | | |
| Sheep, Goats, Lambs and Kids: | | | | | | |
| Hornflies, screwworms and wool maggots | 1/2% | 1 1/3 oz | 2 pt | 1 gal | | |
| Lice and sheep keds | 1/4% | 2/3 oz | 1 pt | 2 qt | | |
| Hogs: Lice | 1/4% | 2/3 oz | 1 pt | 2 qt | | |
| RESIDUAL FLY SPRAY | | | | | | Spray walls, ceilings, partitions to obtain complete coverage with no runoff. Use 1 gallon of diluted spray per 500-1000 sq. ft. |
| A 1/2% spray will give about 2 weeks control. A 1% spray will give 4-6 weeks control. | | | | | | |
| Farm Buildings: poultry houses, livestock shelters, milk room, dairy and hog barns. | 1/2% or 1% | 1 1/3 oz or 2 2/3 oz | 1 qt or 2 qt | 1 gal or 2 gal | | Add 3 ounces of sugar per gallon for added effectiveness. To prevent contamination keep milk and milking equipment covered while spraying. |
| Meat, poultry, and other food processing plants, refuse areas and containers | 1% | 2 2/3 oz | 2 qt | 2 gal | Do not apply in the edible products areas of food processing plants. | |
| Restaurants and supermarkets, Outside area. | 1/2% or 1% | 1 1/3 oz or 2 2/3 oz | 1 qt or 2 qt | 1 gal or 2 gal | Spray outside walls, receiving docks, doorways, and refuse areas. | |
| Dwellings—Outside areas, including porches, patios, and garages. | 1/2% or 1% | 1 1/3 oz or 2 2/3 oz | 1 qt or 2 qt | 1 gal or 2 gal | Spray to point of runoff. | |
| Caged layers—on wire | 2% | 5 1/3 oz | 1 gal | 4 gal | Spray poultry droppings beneath cages | |

ACCEPTED
1-28-72
464-330
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDE ACT OF
1947 (7 U.S.C. 136) REGISTERED
EPA REG. NO. 100-100000000000000000
TO ATTACHED COMMENTS.

SPECIMEN

KORLAN* 4E INSECTICIDE CONCENTRATE

LIVESTOCK SPRAY, DIP AND BACKRUBBER



DIRECTIONS

KORLAN 4E insecticide concentrate mixes readily with water for spray and dip and with oil for backrubber use. Follow directions given in the tables.

DIP RECOMMENDATIONS

| PROBLEM | Ronnal (%) | Amount KORLAN 4E for | | | REMARKS |
|---------------------------------------|------------|-----------------------|----------|---------------|---|
| | | Initial charge of vat | | Recharge rate | |
| | | 100 gal | 1000 gal | 300 gal | |
| LIVESTOCK INSECTS | | | | | 1. Dipping solution should be thoroughly stirred prior to use. Run 25-30 head through the vat as "stirrers" and then return them to the main herd to be redipped. 2. To maintain proper concentration recharge vat when 300 gallons of dip solution have been removed by animals. 3. To replace evaporation loss add water only to previous level of dip solution. 4. If flooding has occurred add KORLAN 4E at the recharge rate. 5. Retreat as necessary but not more often than every 2 weeks. |
| Cattle and Calves: | | | | | |
| Lice | ¼% | 1 qt | 2½ gal | 9 pt | |
| Hornflies, ear ticks and screwworms | ¼% | 2 qt | 5 gal | 9 qt | |
| Sheep, Goats, Lambs, and Kids: | | | | | |
| Lice, sheep keds, and hornflies | ¼% | 2 qt | 5 gal | 9 qt | |
| Wool maggots and screwworms | ½% | 1 gal | 10 gal | 4½ gal | |
| Hogs: Lice | ¼% | 2 qt | 5 gal | 9 qt | |

BACKRUBBER RECOMMENDATIONS

Cattle Backrubber — For hornflies. Aids in the control of lice if used continuously. Aids in the control of face flies if backrubber is placed so animals can rub face. Place where animals regularly travel or congregate.

Mix 1 gallon KORLAN 4E with 55 gallons diesel or mineral oil. Saturate fibre. Recharge as necessary. Do not mix with used motor oil.

SPRAY RECOMMENDATIONS

| PROBLEM | Ronnal (%) | Amount KORLAN 4E for | | | REMARKS |
|---|------------|----------------------|--------|---------|--|
| | | 1 gal | 25 gal | 100 gal | |
| LIVESTOCK INSECTS | | | | | Dilute with water only. Spray entire animal to wet thoroughly. Retreat as necessary but not more often than every 2 weeks. |
| Cattle and Calves: | | | | | |
| Hornflies, ear ticks and screwworms | ½% | 1½ oz | 2 pt | 1 gal | |
| Lice | ¼% | ¾ oz | 1 pt | 2 qt | |
| Lone Star and Winter Ticks | ¾% | 2 oz | 3 pt | 1½ gal | |
| Sheep, Goats, Lambs and Kids: | | | | | |
| Hornflies, screwworms and wool maggots | ½% | 1½ oz | 2 pt | 1 gal | |
| Lice and sheep keds | ¼% | ¾ oz | 1 pt | 2 qt | |
| Hogs: Lice | ¼% | ¾ oz | 1 pt | 2 qt | |
| RESIDUAL FLY SPRAY | | | | | Spray walls, ceilings, partitions to obtain complete coverage with no runoff. Use 1 gallon of diluted spray per 500-1000 sq. ft. |
| A ½% spray will give about 2 weeks control. A 1% spray will give 4-6 weeks control. | | | | | |

SPRAY RECOMMENDATIONS Continued

| PROBLEM | Ronnal (%) | Amount KORLAN 4E for | | | REMARKS |
|--|----------------|----------------------|---------------------|----------------------|--|
| | | 1 gal | 25 gal | 100 gal | |
| FLY BAIT Where flies congregate: floors, windows, and doorways. | 2% | 5½ oz | 1 gal | 4 gal | Add 3 ounces of sugar per gallon diluted spray. Spray or sprinkle at 1 to 2 day intervals until control is obtained. |
| COCKROACH CONTROL Use 1% spray for German roaches, 2% for others. | | | | | Apply as a spot spray to all areas frequented by cockroaches. |
| Buildings: dwellings, institutions and food handling establishments | 1% or 2% | 2¾ oz or 5½ oz | 2 qt or 1 gal | 2 gal or 4 gal | Apply as a spot spray to all areas frequented by cockroaches. Do not use in edible product areas or in serving areas while food is exposed. |
| Food Processing Plants | 1% or 2% | 2¾ oz or 5½ oz | 2 qt or 1 gal | 2 gal or 4 gal | Do not apply in the edible products areas of food processing plants. |
| Meat Packing and Poultry Processing Plants | 2% | 5½ oz | 1 gal | 4 gal | |
| MISCELLANEOUS INSECTS Fleas and spiders in buildings, laundry trucks, and dog kennels. | 1% | 2¾ oz | 2 qt | 2 gal | Apply 1 gallon of dilute spray per 1000 sq. ft. Spray floors, walls, cracks, crevices, sleeping quarters of pets, etc. where insects breed and harbor. |
| Bedbugs in buildings and sleeping quarters, on beds and bedding. | 1% | 2¾ oz | 2 qt | 2 gal | Spray mattress and bedding lightly, including seams and folds, cracks and crevices in bed frames, walls, baseboards, and floors. Allow bedding and mattress to dry before using. |
| Fleas in lawns and kennel runs. | 1% | 2¾ oz | 2 qt | 2 gal | Spray infested areas. Keep children off treated area until spray has dried. |
| Ants, crickets, mosquitoes and silverfish. | 1% | 2¾ oz | 2 qt | 2 gal | Spray areas where insects frequent. |
| Chinch Bugs in lawns (Florida only). Retreat as necessary. Thin Turf or Thick Turf (thatch and organic matter 1½ inch thick or more) | 15% | 4 oz | 10 oz | 40 oz | Apply following a good rain or a drench with ½ to 1 inch of water. Dilute with water only. Use 25 gallons diluted spray per 1250 sq. ft. For thick turf, repeat in two weeks if necessary. |

CAUTION

**KEEP OUT OF REACH OF CHILDREN
INJURIOUS TO THE EYES • IRRITATING TO THE SKIN
HARMFUL IF SWALLOWED • COMBUSTIBLE**

Avoid Eye Contact • Avoid Contact with the Skin • Avoid Prolonged Breathing of Spray Mists • Avoid Contamination of Food, Feedstuffs, and Water Supplies • Keep Away from Heat and Open Flame • Keep Container

| | | | | |
|---|----|-------|--------|--------|
| Sheep, Goats, Lambs, and Kids: Lice, sheep keds, and hornflies | ¼% | 2 qt | 5 gal | 9 qt |
| Wool maggots and screwworms | ½% | 1 gal | 10 gal | 4½ gal |
| Hogs: Lice | ¼% | 2 qt | 5 gal | 9 qt |

vat when 300 gallons of dip solution have been removed by animals.
 3. To replace evaporation loss add water only to previous level of dip solution.
 4. If flooding has occurred add KORLAN 4E at the recharge rate.
 5. Retreat as necessary but not more often than every 2 weeks.

BACKRUBBER RECOMMENDATIONS

Cattle Backrubber — For hornflies. Aids in the control of lice if used continuously. Aids in the control of face flies if backrubber is placed so animals can rub face. Place where animals regularly travel or congregate.

Mix 1 gallon KORLAN 4E with 55 gallons diesel or mineral oil. Saturate fibre. Recharge as necessary. Do not mix with used motor oil.

SPRAY RECOMMENDATIONS

| PROBLEM | Rennel (%) | Amount KORLAN 4E for | | | REMARKS | |
|--|----------------|----------------------|--------------------|----------------------|--|--|
| | | 1 gal | 25 gal | 100 gal | | |
| LIVESTOCK INSECTS | | | | | | |
| Cattle and Calves: | | | | | | |
| Hornflies, ear ticks and screwworms | ½% | 1½ oz | 2 pt | 1 gal | Dilute with water only. Spray entire animal to wet thoroughly. Retreat as necessary but not more often than every 2 weeks. | |
| Lice | ¼% | ¾ oz | 1 pt | 2 qt | | |
| Lone Star and Winter Ticks | ¾% | 2 oz | 3 pt | 1½ gal | | |
| Sheep, Goats, Lambs and Kids: | | | | | | |
| Hornflies, screwworms and wool maggots | ½% | 1½ oz | 2 pt | 1 gal | | |
| Lice and sheep keds | ¼% | ¾ oz | 1 pt | 2 qt | | |
| Hogs: Lice | ¼% | ¾ oz | 1 pt | 2 qt | | |
| RESIDUAL FLY SPRAY | | | | | | |
| A ½% spray will give about 2 weeks control. A 1% spray will give 4-6 weeks control. | | | | | | Spray walls, ceilings, partitions to obtain complete coverage with no runoff. Use 1 gallon of diluted spray per 500-1000 sq. ft. |
| Farm Buildings: poultry houses, livestock shelters, milk room, dairy and hog barns. | ½% or 1% | 1½ oz or 2¾ oz | 1 qt or 2 qt | 1 gal or 2 gal | | Add 3 ounces of sugar per gallon for added effectiveness. To prevent contamination keep milk and milking equipment covered while spraying. |
| Meat, poultry, and other food processing plants, refuse areas and containers | 1% | 2¾ oz | 2 qt | 2 gal | Do not apply in the edible products areas of food processing plants. | |
| Restaurants and supermarkets, Outside area. | ½% or 1% | 1½ oz or 2¾ oz | 1 qt or 2 qt | 1 gal or 2 gal | Spray outside walls, receiving docks, doorways, and refuse areas. | |
| Dwellings — Outside areas, including porches, patios, and garages. | ½% or 1% | 1½ oz or 2¾ oz | 1 qt or 2 qt | 1 gal or 2 gal | Spray to point of runoff. | |
| Caged layers — on wire | 2% | 5½ oz | 1 gal | 4 gal | Spray poultry droppings beneath cages. | |

| | or 2% | or 5½ oz | or 1 gal | or 4 gal | |
|--|-------|----------|----------|----------|--|
| Meat Packing and Poultry Processing Plants | 2% | 5½ oz | 1 gal | 4 gal | food processing plants |
| MISCELLANEOUS INSECTS | | | | | |
| Fleas and spiders in buildings, laundry trucks, and dog kennels. | 1% | 2¾ oz | 2 qt | 2 gal | Apply 1 gallon of dilute spray per 1000 sq ft. Spray floors, walls, cracks, crevices, sleeping quarters of pets etc. where insects breed and harbor. |
| Bedbugs in buildings and sleeping quarters, on beds and bedding. | 1% | 2¾ oz | 2 qt | 2 gal | Spray mattress and bedding lightly, including seams and folds, cracks and crevices in bed frames, walls, baseboards, and floors. Allow bedding and mattress to dry before using. |
| Fleas in lawns and kennel runs | 1% | 2¾ oz | 2 qt | 2 gal | Spray infested areas. Keep children off treated area until spray has dried. |
| Ants, crickets, mosquitoes and silverfish. | 1% | 2¾ oz | 2 qt | 2 gal | Spray areas where insects frequent. |
| Chinch Bugs in lawns (Florida only). Retreat as necessary. | | | | | Apply following a good rain or a drench with ½ to 1 inch of water. |
| Thin Turf or Thick Turf (thatch and organic matter 1½ inch thick or more). | 15% | 4 oz | 10 oz | 40 oz | Dilute with water only. Use 25 gallons diluted spray per 1250 sq ft. For thick turf, repeat in two weeks if necessary. |

CAUTION

KEEP OUT OF REACH OF CHILDREN
INJURIOUS TO THE EYES • IRRITATING TO THE SKIN
HARMFUL IF SWALLOWED • COMBUSTIBLE

Avoid Eye Contact • Avoid Contact with the Skin • Avoid Prolonged Breathing of Spray Mists • Avoid Contamination of Food, Feedstuffs, and Water Supplies • Keep Away from Heat and Open Flame • Keep Container Closed • Wash thoroughly with Soap and Water after Handling
In case of eye contact, flush them with plenty of water and get medical attention.

NOTE TO PHYSICIAN OR VETERINARIAN: This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

THE DOW CHEMICAL COMPANY

AND SUBSIDIARIES

MIDLAND, MICHIGAN 48640, USA ZURICH, SWITZERLAND HONG KONG, BCC
 CORAL GABLES, FLORIDA 33134, USA SARNIA, ONTARIO, CANADA

* Trademark of THE DOW CHEMICAL COMPANY

A1071

IST, 1968.

DUDE USES FOR CONTROL
 (2) USE PRECAUTIONS
 (3) PRECAUTIONARY
 ION REVISED. (4)
 IPANY NAME USED.

ACCEPTED
 1-28-72
 464-330
 UNDER THE FEDERAL INSECTICIDE
 FOR ECN...
 ED... SUBJECT
 TO ATTACHED COMMENTS.

SPECIMEN LABEL



KORLAN* 4E INSECTICIDE CONCENTRATE

LIVESTOCK SPRAY, DIP AND BACKRUBBER



KILLS

ACTIVE INGREDIENTS

| | |
|---|-------|
| Ro-nel (O,O-dimethyl O(2,4,6-trichlorophenyl) phosphorothioate) | 41.2% |
| Aromatic petroleum derivative solvents | 29.3% |
| Inert ingredients | 29.5% |

E.P.A. Registration No. 464-330-AA

PRODUCT ADVANTAGES

Makes a highly stable livestock dip in hard or soft water • Makes both spray and dip • Excellent wetting as spray or dip • Readily mixes with oil for back-rubber use • For livestock of all ages, newborn too! • This one really works

LIVESTOCK REMEDY NOT FOR HUMAN USE

USE PRECAUTIONS

Keep out of reach of children. Avoid contact with skin, eyes, nose, mouth, and clothing. Avoid breathing spray mists. Avoid contact with food, feedstuffs, and water supplies. Keep away from heat and open flame. Keep container closed. Wash thoroughly with soap and water after handling. In case of eye contact, flush them with plenty of water and get medical attention.

NOTE TO PHYSICIAN OR VETERINARIAN: This product contains Ro-nel, which is a phosphorothioate insecticide. It is highly effective against a wide range of insects and mites. It is also highly effective against ticks and lice. It is safe for use on all livestock, including newborn animals.

Warning

Notice

LOT MM

5 GAL

KORLAN 4E

DIRECTIONS

DIP RECOMMENDATIONS

Amount KORLAN 4E per gallon of water

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

BACKRUBBER RECOMMENDATIONS

SPRAY RECOMMENDATIONS

Amount KORLAN 4E per gallon of water

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

SPRAY RECOMMENDATIONS

DIP RECOMMENDATIONS

Amount KORLAN 4E per gallon of water

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

PROBLEMS

STICK INSECTS

LIVESTOCK

THE SPRAY RATE

100 lb 1000 lb 3000 lb

CAUTION

KEEP OUT OF REACH OF CHILDREN
INJURIOUS TO THE EYES • IRRITATING TO THE SKIN
HARMFUL IF SWALLOWED • COMBUSTIBLE

Avoid Eye Contact • Avoid Contact with the Skin • Avoid Prolonged Breathing of Spray Mists • Avoid Contamination of Food, Feedstuffs, and Water Supplies • Keep Away from Heat and Open Flame • Keep Container Closed • Wash thoroughly with Soap and Water after Handling. In case of eye contact, flush them with plenty of water and get medical attention.

NOTE TO PHYSICIAN OR VETERINARIAN

THE DOW CHEMICAL COMPANY

MIDLAND, MICHIGAN 48640, U.S.A. ZÜRICH, SWITZERLAND. TOKYO, JAPAN. BRISBANE, AUSTRALIA. SINGAPORE. MALAYA. SINGAPORE. HONG KONG. CANADA. MEXICO. GUATEMALA. COSTA RICA. PANAMA. VENEZUELA. ARGENTINA. BRAZIL. CHILE. COLOMBIA. CUBA. DOMINICAN REPUBLIC. ECUADOR. EL SALVADOR. GUYANA. HAITI. HONDURAS. ICELAND. INDIA. INDONESIA. JAMAICA. JAPAN. KOREA. KUWAIT. LAOS. LEBANON. LIBERIA. MALAYSIA. MALTA. MEXICO. MOROCCO. NEPAL. NICARAGUA. NORWAY. OMAN. PAKISTAN. PANAMA. PARAGUAY. PERU. PHILIPPINES. POLAND. PORTUGAL. QATAR. ROMANIA. SAUDI ARABIA. SINGAPORE. SLOVAKIA. SOUTH AFRICA. SPAIN. SWEDEN. SWITZERLAND. THAILAND. TRINIDAD AND TOBAGO. TUNISIA. U.S.A. U.K. U.R.S.S. VENEZUELA. YUGOSLAVIA.

1-28-72
464-330

SPECIMEN LABEL