

SPECIMEN LABEL
(BACK)
REDUCED TO 85%



DOWPON* C

**GRASS
KILLER**

DIRECTIONS

GENERAL INFORMATION: DOWPON C is a highly effective, low-cost herbicide for control of established annual and perennial grasses in noncrop areas such as drainage ditch and drainage canal banks, roadsides, fencerows, headlands, around buildings and other farm and industrial structures. Dissolve 15 pounds of DOWPON C in 100 gallons of water and add 2 quarts of DYNAWET* wetting agent included in this package. Apply in sufficient volume to thoroughly wet all grass foliage. This will usually require at least 200 gallons of spray per acre treated.

TO CONTROL JOHNSONGRASS, BERMUDAGRASS, OTHER PEST GRASSES AND CATTAILS: Apply at any time when the grass is growing actively. When used during the spring or summer wait until the first seed heads appear — usually when Johnsongrass is 20 to 40 inches tall or when Bermudagrass is 6 to 12 inches tall. Follow up with another treatment, as required in late summer or fall. Allow the grass to regrow to the growth stage described above before re-treating. When the first treatment is made in late summer or fall allow the grass to reach the first seed-head stage, but apply at least 2 weeks before frost. A follow-up application may be needed the following spring or summer. Two to four well timed and thoroughly applied treatments during two successive growing seasons should

TO SUPPRESS BERMUDAGRASS: 15 pounds of DOWPON C dissolved in 200 gallons of water and sprayed to wet the Bermudagrass at a rate of 200 gallons per acre should provide good growth suppression on drainage ditch banks and similar places where growth is desired to hold the soil in place. Apply any time Bermudagrass growth constitutes a problem. Re-treat as needed. Note: This may cause considerable kill of Bermudagrass under some conditions such as vigorous growth at time of treatment. Do not reuse container. Destroy when empty.

USE PRECAUTIONS

DO NOT USE ON LAND TO BE PLANTED TO FOOD CROPS.

To avoid injury to desirable plants, do not allow spray or spray drift containing DOWPON C to contact foliage or roots of such plants, including lawns and other desirable turf. Do not let spray get into water used for irrigation or other domestic purposes. Flush spraying equipment with plenty of water immediately after use to avoid contamination of later sprays or possible corrosion.

WARNING

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

DIRECTIONS

GENERAL INFORMATION: DOWPON C is a highly effective, low-cost herbicide for control of established annual and perennial grasses in noncrop areas such as drainage ditch and drainage canal banks, roadsides, fencerows, headlands, around buildings and other farm and industrial structures. Dissolve 15 pounds of DOWPON C in 100 gallons of water and add 2 quarts of DYNAWET* wetting agent included in this package. Apply in sufficient volume to thoroughly wet all grass foliage. This will usually require at least 200 gallons of spray per acre treated.

TO CONTROL JOHNSONGRASS, BERMUDAGRASS, OTHER PEST GRASSES AND CATTAILS: Apply at any time when the grass is growing actively. When used during the spring or summer wait until the first seed heads appear—usually when Johnsongrass is 20 to 40 inches tall or when Bermudagrass is 6 to 12 inches tall. Follow up with another treatment, as required, in late summer or fall. Allow the grass to regrow to the growth stage described above before re-treating. When the first treatment is made in late summer or fall allow the grass to reach the first seed-head stage, but apply at least 2 weeks before frost. A follow-up application may be needed the following spring or summer. Two to four well timed and thoroughly applied treatments during two successive growing seasons should provide excellent control of Johnsongrass and Bermudagrass. Dead grass may be burned one month after treatment. On cattails apply when new growth is about 3 feet high up to first seed-head stage. Re-treat as necessary.

TO CONTROL MIXED SPECIES INCLUDING GRASSES, BROADLEAVED WEEDS AND WOODY PLANTS: Apply a tank mixture consisting of 15 pounds DOWPON C, 2 quarts DYNAWET and 2 to 4 quarts KURON* herbicide in 100 gallons of water. Spray to thoroughly wet all vegetation. Follow use precautions for both DOWPON C and KURON when tank mixtures are employed.

TO SUPPRESS BERMUDAGRASS: 15 pounds of DOWPON C dissolved in 200 gallons of water and sprayed to wet the Bermudagrass at a rate of 200 gallons per acre should provide good growth suppression on drainage ditch banks and similar places where growth is desired to hold the soil in place. Apply any time Bermudagrass growth constitutes a problem. Re-treat as needed. Note: This may cause considerable kill of Bermudagrass under some conditions such as vigorous growth at time of treatment. Do not reuse container. Destroy when empty.

USE PRECAUTIONS

DO NOT USE ON LAND TO BE PLANTED TO FOOD CROPS.

To avoid injury to desirable plants, do not allow spray or spray drift containing DOWPON C to contact foliage or roots of such plants, including lawns and other desirable turf. Do not let spray get into water used for irrigation or other domestic purposes. Flush spraying equipment with plenty of water immediately after use to avoid contamination of later sprays or possible corrosion.

WARNING

**KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED
MAY CAUSE BURNS**

**Do Not Get in Eyes, on Skin, or on Clothing or Shoes
Avoid Breathing Spray Mist**

In case of contact, immediately flush skin or eyes with plenty of water for at least 15 minutes; for eyes, get medical attention. Remove contaminated clothing and wash before re-use.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

H1074

THE DOW CHEMICAL COMPANY

AND SUBSIDIARIES

MIDLAND, MICHIGAN 48640, USA ZURICH, SWITZERLAND HONG KONG, BCC
CORAL GABLES, FLORIDA 33134, USA SARNIA, ONTARIO, CANADA

* Trademark of THE DOW CHEMICAL COMPANY

ACCEPTED
March 5, 1975
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 481-23

SPECIMEN LABEL
REDUCED TO 85%



86-1173 PRINTED IN U.S.A. IN DECEMBER, 1974

REPLACES SPECIMEN LABEL 86-1173 PRINTED IN AUGUST, 1972

REVISIONS INCLUDE: (1) EPA ESTABLISHMENT NUMBER ADDED
(2) AGRICULTURAL CHEMICAL STATEMENT ADDED