

ACCEPTED
 APR 2 1968
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND REPELLENT ACT
 FOR ECONOMIC INSECTICIDES RE-
 GISTERED UNDER NO. 464-361



SPECIMEN LABEL

TORDON[®] 212 MIXTURE

HERBICIDE

for the control of annual and deep rooted perennial weeds

ACTIVE INGREDIENTS:
 4-amino-3,5,6-trichloropicolinic acid, as the
 triisopropanolamine salt.....18.1%
 2,4-dichlorophenoxyacetic acid, as the
 triisopropanolamine salt.....37.7%
INERT INGREDIENTS.....44.2%

Acid Equivalents:
 4-amino-3,5,6-trichloropicolinic acid* 10.1%-1 lb. gal.
 2,4-dichlorophenoxyacetic acid.....20.2%-2 lb. gal.

*Known under the trademark TORDON U.S.D.A. Registration No. 464-361

CAUTION
KEEP OUT OF REACH
OF CHILDREN

Read Complete Precautions
on Back Panel

ONE-HALF GALLON

GENERAL INFORMATION

Use TORDON 212 Mixture herbicide as a treatment to control patches of annual and deep rooted perennial weeds including Canada thistle, field bindweed (perennial morning glory), leafy spurge, bur ragweed, Russian knapweed, milkweed, dock, sowthistle, larkspur, toadflax, musk thistle, pigweed, wild sunflower, American wormwood, povertyweed and white horse nettle growing in non-cropland areas such as fence rows, around farm buildings and outer banks of ditches.

TORDON 212 Mixture herbicide is highly potent, persistent and water soluble. To prevent damage to desirable crops and plants follow all directions and precautions.

USE DIRECTIONS

Mix with water and apply as a coarse, low pressure spray (20 to 50 lb. per sq. in.) Apply anytime when fully developed green leaves are present.

For General Use: The rate of TORDON 212 Mixture required varies according to weed species and geographical location. The following table shows the amount of TORDON 212 Mixture that should be mixed in water to make 100 gallons of spray. Apply uniformly to wet the weeds without run-off. This will usually require about 100 gallons per acre.

Some of the weeds to be controlled	TORDON 212 Mixture to use in 100 gal. spray	Remarks
Dock, Larkspur, Pigweed Sowthistle Sunflower (wild) Thistle (Canada) Thistle (Musk) Toadflax (Dalmation) Wormwood (American)	1/2 to 2 gal.	Use lower rates in low rainfall areas in the northern states such as Idaho, Montana, North Dakota, Oregon, South Dakota, Washington and Wyoming. Higher rates should be used in southern states or where rainfall is greater such as Arizona, Arkansas, Kansas, Missouri, New Mexico, Oklahoma and Texas.
Bindweed (field) Horsenettle (white) Knapweed (Russian) Milkweed Ragweed (bur) Spurge (leafy) Toadflax (yellow)	1 to 3 gal.	

For Use on Round Patches of Weeds: Apply the required spray mixture at the amount indicated in the following table.

Feet across Round Patch to be treated (weed area plus 10 foot border)	Gallons of spray mixture to apply
25	1.0
50	4.5
75	10.0
100	18.0
235 or (1 acre)	100.0

NOTE: For small amounts of spray use TORDON 212 Mixture at rate of 1 1/4 to 2 1/2 fluid ounces in each gallon of water.

TORDON 212 Mixture herbicide is effective against a wide range of weeds. For more information contact your State Agricultural Experiment Station or Extension Service for recommendations. Be sure that the use of this product conforms to all applicable laws and regulations.

Be Sure You Follow All Use Precautions Given on This Label and Remember The Following:

1. Use only the recommended amounts. 2. TORDON 212 Mixture should be applied to the soil surface and not mixed over in the soil. Treated soil should not be moved out of the area. Treated soil should not be mixed with soil. Treated soil should not be mixed with soil. Treated soil should not be mixed with soil. Treated soil should not be mixed with soil.

USE PRECAUTIONS

Do Not Allow Spray Drift. TORDON and 2,4-D herbicides may cause damage to plants if applied during either growing or dormant periods. Do not apply or otherwise permit spray to contact desirable plants such as vegetables, ornamentals, cotton, tobacco, tomatoes, potatoes, beans and other valuable broadleaved plants, nor soil containing seeds of such plants. Apply TORDON 212 Mixture only when there is little or no wind. Do not apply by air, as this may cause drift. Do not contaminate water. To avoid crop or other plant damage, do not allow spray to fall onto inner banks or bottom of irrigation canals and do not irrigate through treated areas. Do not use for other domestic purposes.

Do Not Move Treated Soil. Do not go over treated areas or harvesting equipment, or move soil from treated areas. Do not treat areas intended to be used for desks. It usually requires up to 3 years for TORDON herbicide to be removed from the soil. Do not mix in the sprayer with other weedkillers. Other Precautions: Do not store near food, feedstuff, fertilizers or other pesticides. To avoid injury to desirable plants, do not use TORDON 212 Mixture should not be reused to contain equipment and containers thoroughly with water and dispose of crop lands away from water supplies. Containers should be cleaned and buried with waste.

CAUTION
KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. MAY CAUSE SKIN IRRITATION. Avoid Contact with Clothing. Avoid Breathing Spray Mists. Wash Well After Use. Container Closed When Not Using. In case of contact with water; and get medical attention. Remove grossly contaminated clothing and wash before reuse.

NOTICE: Seller warrants that the product conforms to the specifications stated on the label when used under normal conditions of use, but neither this warranty nor the use of this product contrary to label instructions or under conditions not reasonably foreseeable to seller, shall constitute an admission of liability on the part of the seller.

U. S. Patent No. 3,285,922

THE DOW CHEMICAL COMPANY
MIDLAND, MICHIGAN

86-1186 PRINTED IN U.S.A. IN OCTOBER 1967.

THIS IS THE INITIAL PRINTING OF THIS SPECIMEN LABEL.

ACCEPTED
 APR 2 1968
 THE FEDERAL INSECTICIDE
 ACT AND THE FERTILIZER ACT
 ECONOMIC INSECTICIDE REGISTRATION
 NO. 464361

SPECIMEN LABEL



TORDON[®] 212 MIXTURE

HERBICIDE

annual and deep rooted perennial weeds

18.1%
 37.7%
 44.2%

CAUTION
 KEEP OUT OF REACH
 OF CHILDREN

Read Complete Precautions
 on Back Panel

U.S.D.A. Registration No. 464-361

ONE-HALF GALLON

GENERAL INFORMATION

Use TORDON 212 Mixture herbicide as a treatment to control patches of annual and deep rooted perennial weeds including Canada thistle, field bindweed (perennial morning glory), leafy spurge, bur ragweed, Russian knapweed, milkweed, dock, sowthistle, larkspur, toadflax, musk thistle, pigweed, wild sunflower, American wormwood, povertyweed and white horenettle growing in non-cropland areas such as fence rows, around farm buildings and outer banks of ditches.

TORDON 212 Mixture herbicide is highly potent, persistent and water soluble. To prevent damage to desirable crops and plants follow all directions and precautions.

USE DIRECTIONS

Mix with water and apply as a coarse, low pressure spray (20 to 50 lb. per sq. in.) Apply anytime when fully developed green leaves are present.

For General Use: The rate of TORDON 212 Mixture required varies according to weed species and geographical location. The following table shows the amount of TORDON 212 Mixture that should be mixed in water to make 100 gallons of spray. Apply uniformly to wet the weeds without run-off. This will usually require about 100 gallons per acre.

Some of the weeds to be controlled	TORDON 212 Mixture to use in 100 gal. spray	Remarks
Dock, Larkspur, Pigweed Sowthistle Sunflower (wild) Thistle (Canada) Thistle (Musk) Toadflax (Dalmation) Wormwood (American)	1/2 to 2 gal.	Use lower rates in low rainfall areas in the northern states such as Idaho, Montana, North Dakota, Oregon, South Dakota, Washington and Wyoming. Higher rates should be used in southern states or where rainfall is greater such as Arizona, Arkansas, Kansas, Missouri, New Mexico, Oklahoma and Texas.
Bindweed (field) Horenettle (white) Knapweed (Russian) Milkweed Ragweed (bur) Spurge (leafy) Toadflax (yellow)	1 to 3 gal.	

For Use on Round Patches of Weeds: Apply the required spray mixture at the amount indicated in the following table.

Feet across Round Patch to be treated (weed area plus 10 foot border)	Gallons of spray mixture to apply
25	1.0
50	4.5
75	10.0
100	18.0
235 or (1 acre)	100.0

NOTE: For small amounts of spray use TORDON 212 Mixture at rate of 1/4 to 2/2 fluid ounces in each gallon of water.

TORDON 212 Mixture herbicide is effective against a wide range of weeds. Consult your State Agricultural Experiment Station or Extension Service weed specialists for local recommendations. Be sure that the use of this product conforms to all applicable regulations.

Be Sure You Follow All Use Precautions

Given on This Label and Remember These Key Points:
 1. Use only the recommended amounts. 2. TORDON 212 Mixture is persistent. It will carry over in the soil. Treated soil should not be moved out of the treated area. 3. TORDON 212 Mixture is water soluble. It can move with water in irrigation or drainage ditches. 4. Spray drift can damage crops.

USE PRECAUTIONS

Do Not Allow Spray Drift. TORDON and 2,4-D herbicides are highly potent. Tiny amounts may cause damage to plants if applied during either growing or dormant periods. Do not use high pressure sprays. Do not apply or otherwise permit TORDON 212 Mixture or sprays containing it to contact desirable plants such as vegetables, flowers, grapes, fruit trees, ornamentals, cotton, tobacco, tomatoes, potatoes, beans of all types including soybeans, and other valuable broadleaved plants, nor soil containing roots of nearby valuable plants. Apply TORDON 212 Mixture only when there is little or no hazard from spray drift. Coarse sprays are least likely to drift. Do not apply by air, as this increases drift hazard.

Do Not Contaminate Water. To avoid crop or other plant injury, do not treat or allow spray drift to fall onto inner banks or bottom of irrigation and drainage ditches. Dike around and do not irrigate through treated areas. Do not contaminate water used for drinking or other domestic purposes.

Do Not Move Treated Soil. Do not go over treated areas with land levelers, cultivation or harvesting equipment, or move soil from treated areas by any other means.

Do Not Treat Areas Intended To Be Used For Desirable Plants or Food Crops. It usually requires up to 3 years for TORDON herbicide to be deactivated by the soil.

Do Not Mix in the Sprayer With Other Weedkillers or Other Pesticides.

Other Precautions: Do not store near food, feedstuff, fertilizer, seeds, insecticides, fungicides or other pesticides. To avoid injury to desirable plants, containers and sprays for TORDON 212 Mixture should not be reused to contain or apply other materials. Wash equipment and containers thoroughly with water and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with waste.

CAUTION

KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. CAUSES EYE INJURY. MAY CAUSE SKIN IRRITATION. Avoid Contact with Eyes, Skin and Clothing. Avoid Breathing Spray Mists. Wash Well After Handling or Use. Keep Container Closed When Not Using. In case of contact, flush eyes with plenty of water; and get medical attention. Remove grossly contaminated clothing and wash before reuse.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

U. S. Patent No. 3,285,925

967

THE DOW CHEMICAL COMPANY
 MIDLAND, MICHIGAN

REGISTERED IN U.S.A. IN OCTOBER 1967.

INITIAL PRINTING OF THIS SPECIMEN LABEL.