



2,4-D SODIUM SALT

WEED KILLER

Active Ingredients:

2,4-Dichlorophenoxyacetic Acid, Sodium Salt, Monohydrate ... 95%
(2,4-Dichlorophenoxyacetic Acid Equivalent 80.5%)

Inert Ingredients 5%

E.P.A. Registration No. 464-395-AA

DIRECTIONS

WEED CONTROL IN ASPARAGUS: To control wild morningglory and certain other broad' af weeds in asparagus plantings, use 2 to 2½ pounds of 2,4-D Sodium Salt per acre in about 60 gallons of water. In airplane application, use 2 pounds per acre in about 12 gallons of water. A wetting agent can be added if desired in both ground and airplane sprays. Apply on actively growing weeds, usually in April or May. If spears are present, treat immediately after cutting. Make no more than 2 applications and these should be spaced at least 1 month apart during the harvest season. Spears contacted by the spray may be malformed; these spears should be cut immediately and discarded. Post-harvest spraying should be only by ground rig, using drop nozzles to avoid spraying the fern.

USE PRECAUTIONS

Do not apply 2,4-D Sodium Salt, directly to, or otherwise permit it to come into contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants which are sensitive to 2,4-D Sodium Salt and similar materials, and do not permit spray mist containing it to drift onto them, since even minute quantities of the spray may cause severe injury during both growing and dormant periods. (Coarse sprays are less likely to drift.) Accordingly, applications by airplane, ground rigs and hand dispensers should be carried out

86-1222 PRINTED IN U.S.A. IN FEBRUARY, 1972.

REPLACES SPECIMEN LABEL 86-1222 PRINTED IN MARCH,

REVISIONS INCLUDE: (1) ASPARAGUS RECOMMENDATION
CHANGED TO "E.P.A."

2/2

SPECIMEN LABEL ACCEPTED
MAR 9 1970

only when there is no hazard from drift. Do not apply by airplane in the vicinity of cotton, grapes or other desirable 2,4-D Sodium Salt susceptible vegetation. Do not contaminate irrigation ditches or water used for domestic purposes.

Do not store near fertilizers, seeds, insecticides fungicides. Excessive amounts of 2,4-D Sodium Salt in the soil may temporarily inhibit seed germination or plant growth.

To avoid injury to desirable plants, do not store, handle or apply other agricultural chemicals with the same containers or equipment used for 2,4-D Sodium Salt.

Local conditions may effect the use of herbicides. State Agricultural Experiment Station and State Extension Service weed specialists in many states issue recommendations for local conditions.

Be sure that use of this product conforms to all applicable regulations.

Rinse equipment and containers and dispose of wastes by burying in noncrop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

CAUTION
KEEP OUT OF REACH OF CHILDREN
IRRITATING DUST
HARMFUL IF SWALLOWED
Avoid Inhaling Dust
Avoid Contact with Skin, Eyes, or Clothing

**DO NOT STORE NEAR SEEDS, INSECTICIDES,
FUNGICIDES OR FERTILIZERS**

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

B1171

THE DOW CHEMICAL COMPANY
MIDLAND, MICHIGAN 48640

1971.

S REVISED. (2) "USDA"