

ACCEPTED


FEB 7 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR EUGENE W. HINSON REGISTER-
ED UNDER NO. 447177 SUBJECT
TO APPLICABLE REGULATIONS.

TAKE TIME



OBSERVE LABEL
DIRECTIONS



KORLAN

(KORLAN) (D)

PRESSURIZED AEROSOL SPRAY

WOUND TREATMENT

FOR THE TREATMENT OF WOUNDS

IN LIVESTOCK

(KORLAN) (D)

NEVER USE ON CHILDREN

KEEP OUT OF REACH OF CHILDREN

1 POUND

THE DOW CHEMICAL COMPANY

MIDLAND, MICHIGAN 48640

- Kills fast • Saves time
- Worms leave wound quickly

Blue color added for marking treated areas.

CAUTION

KEEP OUT OF REACH OF CHILDREN
CONTENTS UNDER PRESSURE
INJURIOUS TO EYES
HARMFUL IF SWALLOWED

- Do Not Get In Eyes • Avoid Breathing Spray Mists • Harmful If Swallowed
- Wash Hands Thoroughly After Using
- Avoid Contamination of Food and Feedstuffs • Do Not Puncture • Keep Away from Heat; Store At Room Temperature • Never Throw Container Into Fire or Incinerator • Destroy Container When Empty.

In case of human eye contact, flush eyes immediately with plenty of water and get prompt medical attention.

WARNING. KORLAN is a cholinesterase inhibitor. Do not use this product on animal's simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Atropine is an antidote. Consult a veterinarian at the first sign of an adverse reaction.

LIVESTOCK REMEDY —
NOT FOR HUMAN USE
FOR EXTERNAL USE ONLY

DIRECTIONS FOR USE

For use on cattle, sheep, and horses.

Hold container about 3 inches from wound or infested area. Direct spout of container toward the area to be treated. Press valve down to release spray. Wet thoroughly by spraying directly into and about 1 inch around wound.

Screwworm and Blowfly infestations: Use particular care to get spray into deep pockets caused by screwworms. Retreat at 5 to 7 day intervals as needed.

To prevent Screwworm and Blowfly infestations: Apply directly to all open or fresh wounds resulting from dehorning, castration, shearing cuts, etc. Treat navels of newborn animals. Early treatment of wounds may require several treatments. Retreat at 5 to 7 day intervals as needed.

Fleeco-worm infestations: Spray infested area liberally, working toward the center of the infestation. Shearing or removing the wool from the infested area is not necessary. Retreat at 5 to 7 day intervals as needed.

NOTICE: Seller warrants that the product is free from adulteration and contains the amount of active ingredient stated on the label. Seller does not warrant that the product will be effective against all insects and diseases mentioned on the label. Seller does not warrant that the product will be effective against all insects and diseases mentioned on the label. Seller does not warrant that the product will be effective against all insects and diseases mentioned on the label.

Cameras Not
For Copy

ast • Saves time
s leave wound quickly
or added for marking treated

CAUTION

**OUT OF REACH OF CHILDREN
CONTENTS UNDER PRESSURE
INJURIOUS TO EYES
HARMFUL IF SWALLOWED**

**Get In Eyes • Avoid Breathing
Mists • Harmful If Swallowed
Wash Hands Thoroughly After Using
Prevent Contamination of Food and
Fur • Do Not Puncture • Keep
Away from Heat; Store At Room Tem-
perature • Never Throw Container Into
Incinerator • Destroy Container
When Empty.**

In case of human eye contact, flush eyes
thoroughly with plenty of water and get
prompt medical attention.

KORLAN is a cholinesterase in-
hibitor. Do not use this product on animals
recently treated with or exposed to cho-
linesterase inhibiting drugs, pesticides, or
antidotes. Atropine is an antidote. Consult
a veterinarian at the first sign of an adverse

G172

KORLAN

INSECTICIDE

PRESSURIZED LIVESTOCK SPRAY

WOUND PROTECTANT - KILLS
SCREW-WORMS - BLOWFLY MAGGOTS
FOR LIVESTOCK OF ALL AGES
NEWBORN TOO

Active ingredients:
Korlan 50% cholinesterase inhibitor 5.00%
Dihydroxyacetophenone 2.00%
Xylene 2.00%
Total ingredients 9.00%
Residual 1.00%
Dilution of the above ingredients completely standard
with Federal and State standards.

U.S. Patent 2,812,000

CAUTION

KEEP OUT OF REACH OF CHILDREN
Read Complete Precautions on Side Panel

1 POUND

THE DOW CHEMICAL COMPANY

MIDLAND, MICHIGAN 48640

**LIVESTOCK REMEDY —
NOT FOR HUMAN USE
FOR EXTERNAL USE ONLY**

DIRECTIONS FOR USE

For use on cattle, sheep, and horses.

Hold container about 3 inches from wound
or infested area. Direct spout of container
toward the area to be treated. Press valve
down to release spray. Wet thoroughly by
spraying directly into and about 1 inch around
wound.

Screwworm and Blowfly infestations: Use
particular care to get spray into deep pockets
caused by screwworms. Re-treat at 5 to 7
day intervals as needed.

**To prevent Screwworm and Blowfly infesta-
tions:** Apply directly to all open or fresh
wounds such as from dehorning, castration,
shear cuts, etc. Treat navels of newborn ani-
mals. Freely bleeding wounds may require
several treatments. Re-treat at 5 to 7 day
intervals as needed.

Fleece-worm infestations: Spray infested
area liberally, working toward the center of
the infestation. Shearing or removing the
wool from the infested area is not necessary.
Re-treat at 5 to 7 day intervals as needed.

NOTICE: Seller warrants that the product conforms to its
chemical description and is reasonably fit for the purposes
stated on the label when used in accordance with directions
under normal conditions of use, but neither this warranty nor
any other warranty of MERCHANTABILITY or FITNESS FOR
A PARTICULAR PURPOSE, express or implied, extends to the
use of this product contrary to label instructions, or under
abnormal conditions, or under conditions not reasonably con-
siderable to seller, and buyer assumes the risk of any such use.

**Cameraman's Note
Poor Copy**

ACCEPTED

FEB 7 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 44-411 SUBJECT
TO ATTACHED COMMENTS.

TAKE TIME



OBSERVE LABEL
DIRECTIONS



- Kills fast • Saves time
- Worms leave wound quickly

Blue color added for marking treated areas.

CAUTION

**KEEP OUT OF REACH OF CHILDREN
CONTENTS UNDER PRESSURE
INJURIOUS TO EYES
HARMFUL IF SWALLOWED**

**Do Not Get In Eyes • Avoid Breathing
Spray Mists • Harmful If Swallowed
• Wash Hands Thoroughly After Using
• Avoid Contamination of Food and
Feedstuffs • Do Not Puncture • Keep
Away from Heat; Store At Room Tem-
perature • Never Throw Container Into
Fire or Incinerator • Destroy Container
When Empty.**

In case of human eye contact, flush eyes immediately with plenty of water and get prompt medical attention.

WARNING. KORLAN is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. Atropine is an antidote. Consult a veterinarian at the first sign of an adverse reaction.

KORLAN

INSECTICIDE

PRESSURIZED LIVESTOCK SPRAY

WOUND PROTECTANT - KILLS

SCREW-WORMS BLOWFLY MAGGOTS

FOR LIVESTOCK OF ALL AGES

NEWBORN TO 6

Trade Name: Korlan
Registered U.S. Patent Office
DOW CHEMICAL COMPANY
MIDLAND, MICHIGAN 48640
© 1971 DOW CHEMICAL COMPANY
All rights reserved.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Read Complete Directions on Side Panel

1 POUND

THE DOW CHEMICAL COMPANY

MIDLAND, MICHIGAN 48640

**LIVESTOCK REMEDY —
NOT FOR HUMAN USE
FOR EXTERNAL USE ONLY**

DIRECTIONS FOR USE

For use on cattle, sheep, and horses.

Hold container about 3 inches from wound or infested area. Direct spout of container toward the area to be treated. Press valve down to release spray. Wet thoroughly by spraying directly into and about 1 inch around wound.

Screwworm and Blowfly infestations: Use particular care to get spray into deep pockets caused by screwworms. Re-treat at 5 to 7 day intervals as needed.

To prevent Screwworm and Blowfly infestations: Apply directly to all open or fresh wounds such as from dehorning, castration, shear cuts, etc. Treat navels of newborn animals. Freely bleeding wounds may require several treatments. Re-treat at 5 to 7 day intervals as needed.

Flies-worm infestations: Spray infested area liberally, working toward the center of the infestation. Shearing or removing the wool from the infested area is not necessary. Re-treat at 5 to 7 day intervals as needed.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.