

SPECIMEN LABEL



DIRECTIONS
FOR USE



DIRECTIONS
FOR USE



KORLAN 2

Kills Lice and Horn Flies on Cattle by Pour-On Treatment
Kills Keds on Sheep by Spot-On Treatment

KORLAN^{*} 2

INSECTICIDE

**Kills Lice and Horn Flies on Cattle by Pour-On Treatment
Kills Keds on Sheep by Spot-On Treatment**

ADVANTAGES OF POUR-ON TREATMENT

- No costly equipment necessary.
- No unusual handling required.
- Not necessary to wet entire animal.
- Treat young calves too.

ADVANTAGES OF SPOT-ON TREATMENT

- Ready-To-Use
- No Mixing Necessary
- Economical
- Small Dose Per Head
- No Costly Equipment Needed
- No Unusual Handling Required

Active Ingredient:
 Ronnel (O,O-dimethyl O-(2,4,5-trichlorophenyl)
 phosphorothioate) 24.5%
 Inert Ingredients: 75.5%
 E.P.A. Registration No. 484-416-AA E.P.A. Est. No. 464-MI-2

*Trademark of The Dow Chemical Company abroad
 From technical grade. Not suitable for internal use in
 animals

WARNING
KEEP OUT OF REACH
OF CHILDREN

Read Complete Precautions
 on Rear Panel

3.79 L / 1 GAL

THE DOW CHEMICAL COMPANY

ACCEPTED
JAN 14 1976
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, this pesticide
 registered under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 EPA Reg. No. 464-416

SPECIMEN LABEL
 REDUCED TO 75%



DIRECTIONS

CONTROL OF KEDS ON SHEEP • SPOT-ON

A. READY-TO-USE • NO MIXING NECESSARY

B. APPLICATION:

Spot the ready-to-use KORLAN 2 Insecticide along the center line of the back of the animal at the rate of 1 cc. per 50 lbs. body weight. An automatic syringe set at 30 cc. is convenient method of applying KORLAN 2. Used to be...

A single treatment with KORLAN 2 insecticide is effective for control of keds on sheep; however, to prevent reinfestation, it is essential to treat all animals in a flock, and any new animals which subsequently may be added.

KORLAN 2 insecticide should be used in short-haired animals only, no more than 6 weeks after shearing of wooled animals. Treat as often as necessary, but no more often than every 2 weeks.

Do not use on heavily wooded animals because of difficulty in reaching the treated area.

Note: Plastic syringes should not be used to apply KORLAN 2 for control of sheep keds.

THIS PACKAGE CONTAINS ENOUGH KORLAN 2 INSECTICIDE FOR KEDS TO TREAT:

- 1261-50 lb. lambs • 841-75 lb. lambs
- 630-100 lb. lambs or ewes
- 504-125 lb. sheep • 420-150 lb. sheep

NOTICE: This material is a restricted use pesticide. It is to be used only by persons who are registered with the EPA under the Federal Insecticide, Fungicide, and Rodenticide Act. It is to be used only in accordance with the label directions. It is to be stored in its original container and is to be disposed of in accordance with the label directions. It is to be used only in accordance with the label directions. It is to be used only in accordance with the label directions.



KORLAN 2

INSECTICIDE

**Kills Lice and Horn Flies on Cattle by Pour-On Treatment
 Kills Keds on Sheep by Spot-On Treatment**

ADVANTAGES OF POUR-ON TREATMENT

- No costly equipment necessary
- No unusual handling required
- Not necessary to wet entire animal
- Treat young calves too

ADVANTAGES OF SPOT-ON TREATMENT

- Ready-To-Use
- No Mixing Necessary
- Economical
- Small Dose Per Head
- No Costly Equipment Needed
- No Unusual Handling Required

WARNING
 KEEP OUT OF REACH OF CHILDREN
 Read Complete Precautions on Rear Panel

3.79 L / 1 GAL



DIRECTIONS

CONTROL OF LICE AND HORN FLIES ON CATTLE • POUR-ON

A. MIXING: Please read carefully.

CAUTION: ALWAYS ADD WATER TO KORLAN 2. NEVER ADD KORLAN 2 TO WATER.

The treatment mixture will contain **4 parts water** and **1 part KORLAN 2**. This quantity is given by proportion of 4:1 on product.

1. With vigorous stirring, slowly add 4 parts of water to 1 part of KORLAN 2. Continue stirring for a minimum of 2 minutes. Then, when stirring has stopped, slowly add the remaining 3 parts of water to the mixture until a thick, creamy emulsion is formed.

Remember: Always add the water to the KORLAN 2 with vigorous stirring.

2. In cold weather, use warm water (20° R.) in making KORLAN 2 mixtures. Do not use tap water which is as well as cold water.

3. Use a clean, closed bucket for mixing. Do not use a mixture. NEVER add KORLAN 2 to water. Do not use a mixture of water and KORLAN 2. Do not use a mixture of water and KORLAN 2.

NOTE: KORLAN 2 is a restricted use pesticide. It is to be used only by persons who are registered with the EPA under the Federal Insecticide, Fungicide, and Rodenticide Act. It is to be used only in accordance with the label directions. It is to be stored in its original container and is to be disposed of in accordance with the label directions.

B. APPLICATION:

Apply the treatment mixture along the center line of the back of the animal, spreading it over the entire length of the animal. Do not allow the mixture to run down the sides of the animal. However, if the animal is heavily infested with lice or horn flies, it may be necessary to apply the mixture to the sides of the animal. A minimum of 1 cup of mixture is required for application and spreading the mixture.

A single treatment with KORLAN 2 Insecticide is usually highly effective for control of lice and horn flies on cattle heavily infested with ticks and other parasites. It is to be treated after 2 or 3 weeks. Horn flies will be controlled in 2 weeks, however, insects with a reproductive cycle of heavy rains will require a second application.

KORLAN 2 may be used any time of the year when horn flies are a problem. Treatment should be repeated as needed.

THIS PACKAGE CONTAINS ENOUGH KORLAN 2 INSECTICIDE FOR LICE AND HORN FLIES TO TREAT:

- 320-200 lb. cattle • 210-300 lb. cattle
- 180-400 lb. cattle • 125-500 lb. cattle
- 105-600 lb. cattle • 90-700 lb. cattle
- 80-900 lb. or heavier cattle



KORLAN 2

**Kills Lice and Horn Flies on Cattle by Pour-On
 Kills Keds on Sheep by Spot-On Treatment**

USE PRECAUTIONS

Do not use KORLAN 2 on animals producing milk for human consumption. Dairy cows should be treated early in the dry period. Do not treat within 7 days of freshening. If dairy cows or heifers freshen within 7 days of treatment, milk must not be used for food within 7 days of treatment. Under some conditions, KORLAN 2 may cause slight scurfiness in cattle and sheep.

KORLAN 2 Insecticide is a cholinesterase inhibitor. Do not use on animals producing milk for human consumption. Do not use on animals producing milk for human consumption. Do not use on animals producing milk for human consumption.

Treat healthy animals only.

Do not apply to sheep within 28 days of slaughter.

Rinse equipment and containers and dispose of wastes by dropping lands away from water supplies. Containers should be punctured holes in them and buried with wastes.

Avoid contamination of food, feedstuffs, and water supplies.

Keep material out of reach of animals.

WARNING

**KEEP OUT OF REACH OF CHILDREN
 IRRITATING TO THE SKIN
 MAY BE HARMFUL IF SWALLOWED
 MAY BE HARMFUL OR FATAL
 IF ABSORBED THROUGH SKIN
 COMBUSTIBLE**

In case of eye contact, flush with plenty of water • Do skin. In case of skin contact wash thoroughly with soap. Remove contaminated clothing and wash before re- thoroughly with soap and water after handling • Keep heat or open flame.

NOTE TO PHYSICIAN OR VETERINARIAN: This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

12102-016-3

SEE DIRECTIONS ON SIDE PANEL

THE DOW CHEMICAL COMPANY
 AND SUBSIDIARIES

MIDLAND MICHIGAN 48640 USA ZÜRICH SWITZERLAND HONG KONG
 CORAL GABLES FLORIDA 33134 USA SARNIA ONTARIO CANADA
 *Trademark of THE DOW CHEMICAL COMPANY

86-1245 PRINTED IN U.S.A. IN DECEMBER, 1975.

REPLACES SPECIMEN LABEL 86-1245 PRINTED JULY, 1975.
 DISCARD PREVIOUS SPECIMEN LABELS
 REVISIONS INCLUDE: (1) ADDITIONAL USE AND DIRECTIONS FOR KEDS ON SHEEP.

ACCEPTED
 JAN 14 1976
 EPA



SPECIMEN LABEL
 REDUCED TO 75%



KORLAN 2

**Kills Lice and Horn Flies on Cattle by Pour-On Treatment
 Kills Keds on Sheep by Spot-On Treatment**

USE PRECAUTIONS

Do not use Korlan 2 on animals producing milk for human consumption. Dairy animals should be treated during the dry period. Do not treat dairy cows. A 24-hour withdrawal period is required before the milk from treated animals is used for human consumption. Do not use Korlan 2 on animals that are to be used for food.

For sheep and lambs, KORLAN 2 is a spot-on treatment for the control of lice and keds.

KORLAN 2 is a pour-on treatment for the control of lice and horn flies on cattle. It is a pesticide which is highly effective against these pests. It is a contact insecticide which is highly effective against these pests. It is a contact insecticide which is highly effective against these pests.

Treat healthy animals only.

Do not apply to sheep with a wet coat.

Rinse equipment and containers and dispose of contents in a safe place. Do not use on crop lands, away from water, and do not use on animals that are to be used for food.

Avoid contamination of feed, feed troughs, and water troughs.

Keep animals out of reach of children.

WARNING

**KEEP OUT OF REACH OF CHILDREN
 IRRITATING TO THE SKIN
 MAY BE HARMFUL IF SWALLOWED
 MAY BE HARMFUL OR FATAL
 IF ABSORBED THROUGH SKIN
 COMBUSTIBLE**

In case of eye contact, flush with plenty of water • Do not get on the skin. In case of skin contact wash thoroughly with soap and water • Remove contaminated clothing and wash before reuse • Wash thoroughly with soap and water after handling • Keep away from heat or open flame.

NOTE TO PHYSICIAN OR VETERINARIAN: Korlan 2 is a pesticide which is highly effective against these pests. It is a contact insecticide which is highly effective against these pests.

SEE DIRECTIONS ON SIDE PANEL

THE DOW CHEMICAL COMPANY

THE DOW CHEMICAL COMPANY
 1801 Alexander Drive
 Freeport, New York 11520

KORLAN 2^{*}

INSECTICIDE

**Kills Lice and Horn Flies on Cattle by Pour-On Treatment
 Kills Keds on Sheep by Spot-On Treatment**

ADVANTAGES OF POUR-ON TREATMENT

- No costly equipment necessary.
- No unusual handling required.
- Not necessary to wet entire animal.
- Treat young calves too.

ADVANTAGES OF SPOT-ON TREATMENT

- Ready-To-Use
- No Mixing Necessary
- Economical
- Small Dose Per Head
- No Costly Equipment Needed
- No Unusual Handling Required

Active Ingredient
 Ronnel (O,O-dimethyl O-(2,4,5-trichlorophenyl)
 phosphorothioate 24.5%
 Inert Ingredients 75.5%
 EPA Registration No. 464-416-AA EPA Est. No. 464-MI-2

Trademark of The Dow Chemical Company abroad
 From technical grade. Not suitable for internal use in
 animals.

WARNING
**KEEP OUT OF REACH
 OF CHILDREN**

Read Complete Precautions
 on Rear Panel

3.79 L / 1 GAL

DIRECTIONS CONTROL OF LICE AND HORN FLIES ON CATTLE • POUR-ON

A MIXING: Please read carefully

CAUTION: Do not use on animals producing milk for human consumption.

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

1 part KORLAN 2 4 parts water

**THIS PACKAGE CONTAINS ENOUGH
 KORLAN 2 INSECTICIDE FOR LICE AND
 HORN FLIES TO TREAT**

320 200 lb. cattle • 210 300 lb. cattle
 160 400 lb. cattle • 125 500 lb. cattle
 105 600 lb. cattle • 90 700 lb. cattle
 80 800 lb. or heavier cattle

U.S.A. IN DECEMBER, 1975.

LABEL AND PRINTING DATE, 1975.

FORM NO. 100-100

© 1975 THE DOW CHEMICAL COMPANY



KORLAN^{*} 2

INSECTICIDE

Kills Lice and Horn Flies on Cattle by Pour-On Treatment
Kills Keds on Sheep by Spot-On Treatment

ADVANTAGES OF POUR-ON TREATMENT

- No costly equipment necessary.
- No unusual handling required.
- Not necessary to wet entire animal.
- Treat young calves too.

ADVANTAGES OF SPOT-ON TREATMENT

- Ready-To-Use
- No Mixing Necessary
- Economical
- Small Dose Per Head
- No Costly Equipment Needed
- No Unusual Handling Required

Active Ingredient:
 Ronnel† (O,O-dimethyl O-
 (2,4,5-trichlorophenyl)
 phosphorothioate††) 24.5%
 Inert Ingredients: 75.5%
 E.P.A. Registration No. 464-416-AA
 E.P.A. Est. No. 464-MI-2

†Trademark of The Dow Chemical Company abroad
 ††From technical grade.
 Not suitable for internal use in animals

WARNING
KEEP OUT OF REACH
OF CHILDREN

Read Complete Precautions
 on Rear Panel

1.89 L / 64 FL OZ (2 QT)

FOR CATTLE LABEL



SPECIMEN LABEL



KORLAN

**Kills Lice and Horn Flies on Cattle by Pour-On
Kills Keds on Sheep by Spot-On**

USE PRECAUTIONS

Do not use KORLAN 2 on animals producing milk. Dairy cows should be treated early in their dry period within 7 days of freshening. If dairy cows or heifers are treated, milk must not be used for food within 7 days.

Under some conditions, KORLAN 2 may cause slight irritation to the skin of cattle and sheep.

KORLAN 2 Insecticide is a cholinesterase inhibiting pesticide chemical having cholinesterase inhibiting activity. Do not use on animals until 24 hours after treatment.

Treat healthy animals only.

Do not apply to sheep within 28 days of slaughter.

Rinse equipment and containers and dispose of crop lands away from water supplies. Contain and bury or incinerate empty containers and their contents. Do not use for anything other than intended purpose.

Avoid contamination of food, feedstuffs, and water.

Keep material out of reach of animals.

WARNING

**KEEP OUT OF REACH OF CHILDREN
IRRITATING TO THE SKIN
MAY BE HARMFUL IF SWALLOWED
MAY BE HARMFUL IF ABSORBED THROUGH SKIN
COMBUSTIBLE**

In case of eye contact, flush with plenty of water. In case of skin contact wash thoroughly with soap and water after handling. Remove contaminated clothing and wash thoroughly with soap and water after handling.

NOTE TO PHYSICIAN OR VETERINARIAN: This is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote. 12142-003-2

SEE DIRECTIONS ON SIDE OF CONTAINER

THE DOW CHEMICAL COMPANY

AGRICULTURAL DIVISION, 1801 NORTH AVENUE, WILMINGTON, DELAWARE 19880

DIRECTIONS
CONTROL OF KEDS ON SHEEP • SPOT-ON

A. READY-TO-USE • NO MIXING NECESSARY

B. APPLICATION:
Spot the ready-to-use KORLAN 2 Insecticide along the center line of the back of the animal at the rate of 3 cc/ml per 50 lb. body weight. An automatic syringe set at 3 cc is a convenient method of applying KORLAN 2 Insecticide.

A single treatment with KORLAN 2 Insecticide is effective for control of keds on sheep, however, to prevent reinfestation, it is essential to treat all animals in a flock and any new animals which subsequently may be added.

KORLAN 2 Insecticide should be used on short wooled animals only no more than 6 weeks after shearing. If reinfestation occurs, retreat as necessary, but not more often than every 2 weeks.

Do not use on long woolled animals because of potential wool damage in treated area.

Note: Plastic syringes should not be used to apply KORLAN 2 for control of sheep keds.

THIS PACKAGE CONTAINS ENOUGH KORLAN 2 INSECTICIDE FOR KEDS TO TREAT:

- 630-50 lb. lambs • 120-75 lb. lambs
- 315-100 lb. lambs or ewes
- 252-125 lb. sheep • 210-150 lb. sheep

NOTICE: This is a cholinesterase inhibiting pesticide chemical having cholinesterase inhibiting activity. Do not use on animals until 24 hours after treatment. Under some conditions, KORLAN 2 may cause slight irritation to the skin of cattle and sheep. Treat healthy animals only. Do not apply to sheep within 28 days of slaughter. Rinse equipment and containers and dispose of crop lands away from water supplies. Contain and bury or incinerate empty containers and their contents. Do not use for anything other than intended purpose. Avoid contamination of food, feedstuffs, and water. Keep material out of reach of animals.

KORLAN 2
INSECTICIDE

**Kills Lice and Horn Flies on Cattle by Pour-On Treatment
Kills Keds on Sheep by Spot-On Treatment**

ADVANTAGES OF POUR-ON TREATMENT

- No costly equipment necessary.
- No unusual handling required.
- Not necessary to wet entire animal.
- Treat young calves too.

ADVANTAGES OF SPOT-ON TREATMENT

- Ready-To-Use
- No Mixing Necessary
- Economical
- Small Dose Per Head
- No Costly Equipment Needed
- No Unusual Handling Required

WARNING

KEEP OUT OF REACH OF CHILDREN

Read Complete Precautions on Rear Panel

Active Ingredient
Ronnell, O,O-dimethyl O (2,4,5-trichlorophenyl) phosphorothioate 24.5
Inert Ingredients 75.5
EPA Registration No. 461-115-AA
EPA Est. No. 161-MI-2
Trade Name: The DOW Chemical Company
Formulation: emulsion
Net Content: 2 QT (64 FL OZ)

1.89 L 64 FL OZ (2 QT)

DIRECTIONS
CONTROL OF LICE AND HORN FLIES ON CATTLE • POUR-ON

A. MIXING: Please read carefully.

CAUTION: ALWAYS ADD WATER TO KORLAN 2. NEVER ADD KORLAN 2 TO WATER.

The treatment mixture will contain 4 parts water and 1 part KORLAN 2. These 2 quarts will make 2 1/2 gallons of mixed pour-on product.

1. With 1 quart of water, slowly add 1/2 of the total amount of water to be used to the total amount of KORLAN 2 to be used, and continue stirring for a minimum of 2 minutes. Then, when stirring vigorously, slowly add the remaining 3 parts of water. Continue stirring until a thick creamy emulsion is formed.

Remember: Always add the water to the KORLAN 2 with vigorous mixing.

2. If you have water, use warm water to help in making the KORLAN 2 mixture. Store the mixture in a clean container as well as add water.

3. Use a clean 1/2 gallon bucket for each batch of mixture. NEVER add KORLAN 2 to a partially full batch of mixture. It should be used as soon as it is mixed properly.

NOTE: KORLAN 2 should be stored in a cool, dry place. KORLAN 2 should be stored in a clean, dry container. Do not store KORLAN 2 in hot water, or in a place where it will be exposed to direct sunlight.

B. APPLICATION:

Apply the thick mixture along the center line of the back of the animal, spreading it over the entire length of the animal. Use 1 quart of mixture for each animal per treatment. However, if the animal is a dairy cow, use 1/2 quart of mixture. A mixing and dosing procedure is given on the label for applying and spreading the mixture.

A single treatment with KORLAN 2 Insecticide is usually highly effective for control of lice and horn flies on cattle. However, if the treated animal is heavily infested with sucking lice, it may need to be retreated after 2 or 3 weeks. Horn flies will be controlled for 3 weeks, however, as long as they do not lay eggs. The heavy fly will be controlled immediately.

KORLAN 2 Insecticide will kill the eggs of the pest when they hatch. This is a property of the formulation and is not treated as such.

THIS PACKAGE CONTAINS ENOUGH KORLAN 2 INSECTICIDE FOR LICE AND HORN FLIES TO TREAT:

- 160-200 lb. cattle • 106-300 lb. cattle
- 80-400 lb. cattle • 64-500 lb. cattle
- 54-600 lb. cattle • 45-700 lb. cattle
- 40-800 lb. or heavier cattle



KORLAN 2

INSECTICIDE

**Kills Lice and Horn Flies on Cattle by Pour-On Treatment
Kills Keds on Sheep by Spot-On Treatment**

ADVANTAGES OF POUR-ON TREATMENT

- No costly equipment necessary
- No unusual handling required
- Not necessary to wet entire animal
- Treat young calves too

ADVANTAGES OF SPOT-ON TREATMENT

- Ready-To-Use
- No Mixing Necessary
- Economical
- Small Dose Per Head
- No Costly Equipment Needed
- No Unusual Handling Required

WARNING

KEEP OUT OF REACH OF CHILDREN

Read Complete Precautions on Rear Panel

Active Ingredient

Ronnel (O,O-dimethyl O (2,4,5-trichlorophenyl) phosphorothioate)

Inert Ingredients

E.P.A. Registration No. 164-116-AA
E.P.A. Est. No. 164-M-2

Trade Name of The Dow Chemical Company
DOW CHEMICAL COMPANY

1.89 L 64 FL OZ (2 QT)



DIRECTIONS

CONTROL OF LICE AND HORN FLIES ON CATTLE • POUR-ON

A. MIXING: Please read carefully.

CAUTION: ALWAYS ADD WATER TO KORLAN 2. NEVER ADD KORLAN 2 TO WATER.

The treatment mixture will contain 4 parts water and 1 part KORLAN 2. These 2 quarts will make 2 1/2 gallons of mixed pour-on product.

1. With vigorous stirring, slowly add 1 1/2 quarts of water to be used to the total amount of KORLAN 2 to be used, and continue stirring for a minimum of 2 minutes. Then, while stirring, generously slowly add the remaining 3 parts of water. Continue mixing until a thick, creamy emulsion is formed.

Remember, always add the water slowly to KORLAN 2 with vigorous mixing.

2. In cold weather, use warm water (70-80 F.) in making KORLAN mixtures because this product will not mix as well in cold water.

3. Use a clean lined bucket for each batch of pour-on mixture. NEVER add KORLAN 2 to a batch of 4 parts of water or pour-on mixture because it will fail to thicken properly.

NOTE: KORLAN 2 insecticide should be stored above 32 F. KORLAN 2 starts crystalizing at 20 F. Crystals can be redissolved by warming the container of KORLAN 2 in hot water or by putting it in a heated room for 24 hours or more.

B. APPLICATION:

Apply the thick mixture along the center line of the back of the animal. Spread it over the entire length of the animal, using 1 1/2 quarts of the mixture per 1,000 lb. body weight. 8 1/2 quarts is adequate for an animal weighing 800 lbs. or more. A long handled dipper is convenient for applying and spreading the mixture.

A single treatment with KORLAN 2 insecticide is usually highly effective for control of lice on cattle. However, cattle heavily infested with sucking lice may need to be re-treated after 2 or 3 weeks. Horn flies will be controlled for 3 weeks, however, as is true with any topical insecticide, heavy rains will decrease residual activity.

KORLAN 2 may be used any time of the year when horn or horn flies are a problem. If reinfestation occurs, re-treat as needed.

THIS PACKAGE CONTAINS ENOUGH KORLAN 2 INSECTICIDE FOR LICE AND HORN FLIES TO TREAT:

- 180-200 lb. cattle • 106-300 lb. cattle
- 80-400 lb. cattle • 64-500 lb. cattle
- 54-600 lb. cattle • 45-700 lb. cattle
- 40-800 lb. or heavier cattle

SPECIMEN LABEL



KORLAN 2

**Kills Lice and Horn Flies on Cattle by Pour-On Treatment
Kills Keds on Sheep by Spot-On Treatment**

USE PRECAUTIONS

Do not use KORLAN 2 on animals producing milk for human consumption. Dairy cows should be treated early in their dry period. Do not treat dairy cows within 7 days of freshening. If dairy cows or heifers freshen within 7 days of treatment, milk must not be used for food within 7 days of treatment.

Under some conditions, KORLAN 2 may cause slight scurfiness to the backs of cattle and sheep.

KORLAN 2 insecticide is a cholinesterase inhibitor. Do not use any drug or pesticide chemical having cholinesterase inhibiting activity, either simultaneously or within a few days before or after treatment with KORLAN 2.

Treat healthy animals only.

Do not apply to sheep within 28 days of slaughter.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

Avoid contamination of food, feedstuffs, and water supplies.

Keep material out of reach of animals.

WARNING

**KEEP OUT OF REACH OF CHILDREN
IRRITATING TO THE SKIN**

**MAY BE HARMFUL IF SWALLOWED
MAY BE HARMFUL OR FATAL
IF ABSORBED THROUGH SKIN**

COMBUSTIBLE

In case of eye contact, flush with plenty of water • Do not get on the skin. In case of skin contact wash thoroughly with soap and water • Remove contaminated clothing and wash before reuse • Wash thoroughly with soap and water after handling • Keep away from heat or open flame.

NOTE TO PHYSICIAN OR VETERINARIAN: This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

12142-003-2

131-175

SEE DIRECTIONS ON SIDE PANEL

THE DOW CHEMICAL COMPANY

MIDLAND, MICHIGAN 48661, U.S.A. • ZÜRICH, SWITZERLAND • HONG KONG, CHINA
LORAIN, BARCELONA, BRISBANE, U.S.A. • SARNIA, ONTARIO, CANADA
• TOKYO, JAPAN • SYDNEY, AUSTRALIA