

464-448

07/10/1978

1/7



LORSBAN* 4E

INSECTICIDE

*For Control of Certain Insects Infesting
Cotton and Peaches*

Complete Directions for Use, Use Precautions and Restrictions,
Hazards and Other Important Information
For the Proper Use, Handling, Storage and Disposal of this Product

EPA Reg. No. 464-448

A678

Trademark of THE DOW CHEMICAL COMPANY

**READ ENTIRE LABEL
BEFORE USING THIS PRODUCT**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION: (TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

In case of an emergency endangering life or property involving this product, call collect 517-636-4400.

ACTIVE INGREDIENTS:

Chlorpyrifos (O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate) 40.7%
Aromatic petroleum derivative solvent 22.8%

INERT INGREDIENTS: 36.5%

Contains 4 pounds of chlorpyrifos per gallon

CONTENTS

Hazards	Page
Human and Domestic Animal	3
Physical and Chemical	3
Environmental	4
Directions for Use	
General Information	5
Mixing Instructions	5
Use and Dosage Recommendations	5
Cotton	5
Peaches	8
Storage and Disposal	9

3/7

HAZARDS**KEEP OUT OF REACH OF CHILDREN****WARNING****PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS****MAY BE FATAL IF SWALLOWED
MAY BE ABSORBED THROUGH
SKIN • MAY BE INJURIOUS
TO EYES AND SKIN ••****Do Not Take Internally • Do Not Get
in Eyes, on Skin or on Clothing
Avoid Breathing Vapors and
Spray Mist •••••****Wash Thoroughly After Handling****In case of contact, immediately flush con-
taminated eyes with plenty of water and
get prompt medical attention. Promptly
wash contaminated skin with soap and
plenty of water. Wash contaminated
clothing before reuse.****NOTE TO PHYSICIAN:** Chlorpyrifos is a
cholinesterase inhibitor. Treat symp-
tomatically. Atropine is an antidote.**Physical and Chemical hazards****COMBUSTIBLE****Do Not Use or Store Near Heat
or Open Flame • Do Not Cut or
Weld Container****In case of an emergency endangering
life or property involving this
product, call collect****517-636-4400****AGRICULTURAL CHEMICAL****Do Not Ship or Store with Food, Feeds,
Drugs or Clothing****Environmental**

This product is highly toxic to bees ex-
posed to direct treatment or residues on
crops. Avoid use when bees are actively
foraging. Protective information may be
obtained from your Cooperative Agri-
cultural Extension Service.

LORSBAN 4E insecticide is toxic to fish,
birds and other wildlife. Fish and crusta-
ceans may be killed if their waters are
contaminated with this product. Keep out
of lakes, streams, ponds, tidal marshes
and estuaries. Do not apply where runoff is
likely to occur. Do not apply when weather
conditions favor drift from treated areas.
Do not contaminate water by cleaning of
equipment or disposal of wastes. Use this
product only as specified on this label.

DIRECTIONS FOR USE

**It is a violation of federal law to use this
product in a manner inconsistent with its
labeling.**

General Information

LORSBAN 4E insecticide forms an emulsion when diluted with water and is suitable for use in all conventional hand- and power-operated ground spray equipment. Aerial spray equipment also may be used where specified.

Mixing Instructions

To prepare the spray, start filling the spray tank with the required amount of water, add the recommended amount of LORSBAN 4E, and complete filling the tank with the balance of the water needed. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture.

Use and Dosage Recommendations

COTTON: Use LORSBAN 4E for control of the following pests at the dosages indicated:

Plant bugs

(*Lygus*, *Mirids*) 1 pint per acre;

Pink bollworm, bollworms,

(*Heliothis* spp.), **boll weevil,**

spider mites 2 pints per acre;

Mix the required dosage with sufficient water to ensure thorough coverage of plants and apply using aerial or power-operated ground spray equipment. For aerial application, use at least 1 gallon of water per acre. Treat when field counts indicate damaging insect populations are developing or present. Re-treat as necessary to maintain control.

PAGE 6

For effective control of spider mites when large numbers of eggs are present, apply a second spray within 5 to 7 days after initial treatment to control newly hatched larvae.

For best results on bollworms, it is suggested that fields be scouted twice per week and treatments made when worms are 1/4 inch or less in length. The following table illustrates the size of worms in relation to age and stage of development (instar) as a guide to timing of treatments for best control. From the table it can be seen that a scouting schedule of only once per week will not be satisfactory since the worms may be too big to control effectively by the seventh or eighth day.

Timing for the best worm control

	Age (days)	Size	Instar
Get the worms at this stage	Hatch	1/16"	Hatch
	3	3/32"	I
	5	9/32"	II
	6	7/16"	III
	8	11/16"	IV
1/16"			
3/32"			
9/32"			
7/16"			
11/16"			



Proper application techniques help to ensure thorough spray coverage and correct dosage and are thus important in obtaining good control of pests. Consider these suggestions when applying LORSBAN 4E Insecticide on cotton:

Aerial Application: Shorten boom length to reduce spray from entering the vortices at the wing tips. Swath width should be reduced when wind direction is the same as direction of spraying.

The proper nozzle arrangement and swath width to avoid skips and excessive vortices effect can be checked out by flying over a paper tape (adding machine paper) using water with or without methylene blue dye. (The dye gives a permanent record.)

Flying at a height of 5-15 feet above the target results in the best coverage.

Nozzle orientation of the boom is important. More break-up occurs when nozzles are pointed straight down versus the straight back position. Desired droplet size (100-200 microns) can be obtained by angling the nozzles somewhere in this range.

Marking of swath by flagging or permanent markers is essential.

Ground Application: Orient the boom and nozzles so that uniform coverage is obtained. The swath width should not be wider than the boom; drift spray is wasted spray so do not depend on it. Use flat spray or disc-core hollow cone nozzles with maximum spacing of 20 inches and a spray pressure of 40-60 psi with a droplet size of 100-200 microns.

Restrictions: Do not apply within 14 days before harvest. Do not allow livestock to graze in treated areas. Do not feed gin trash or treated forage to livestock.

PEACHES: Use LORSBAN 4E Insecticide for the control of peach tree borers by application as a trunk spray before newly hatched borers enter the trees. Mix 3 quarts of LORSBAN 4E with 100 gallons of water and apply as a coarse, low-pressure spray to give uniform coverage of tree trunks. Thoroughly wet all bark areas from ground level to scaffold limbs. Do not allow spray to contact fruit. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat in your area.

Restrictions: Make only one application per season. Do not apply within 14 days before harvest. Do not allow meat or dairy animals to graze in treated orchards.

5/7

STORAGE AND DISPOSAL

Storage: Keep away from food, feedstuffs, and domestic water supplies. Handle concentrate in a ventilated area. Keep container closed.

Disposal: Do not reuse empty container for any purpose. Promptly crush or perforate and bury with wastes in a location away from water supplies. Follow official local pesticide container disposal regulations where applicable.

ATTENTION:

Do Not Cut or Weld Container

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

NOTES

