

SPECIMEN LABEL



GARLON* 4

HERBICIDE

For the Control of Woody Plants and Broadleaf Weeds on Rights-of-Way,
Industrial Sites and Non-crop Areas, and for Use in Forest Site Preparation

ACTIVE INGREDIENT

Triclopyr (3,3,4-trichloro-2-pyridinyloxyacetic acid), butoxyethyl ester

44.3%

INERT INGREDIENTS

Acid Equivalent: Triclopyr
Contains Petroleum Distillates
EPA Registration No. 464-554

44.3% - 1 lb/gal
EPA Est. 164-213

PRECAUCION AL USUARIO: Si usted no lee Ingles, no use este producto hasta que se allquela le haya sido explicada ampliamente.

TRANSLATION TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Precautionary Statements
Hazardous to Humans and Domestic Animals

HARMFUL IF SWALLOWED

Avoid Contact with Skin, Eyes or Clothing

Avoid Contamination of Food

Wash Thoroughly After Handling

Statement of Practical Treatment

In case of contact, flush skin with plenty of water. Call medical attention if irritation persists. Remove and wash contaminated clothing before reuse.

If swallowed, do not induce vomiting. Call a physician.

Physical or Chemical Hazards

COMBUSTIBLE

Do Not Use or Store Near Heat or Open Flame

Do Not Cut or Weld Container

Environmental Hazards

This pesticide is toxic to fish. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur.

See Back Panel for Important Use Precautions.

In case of an emergency and/or burning fire or property involving this product, call collect 817-838-4400

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, Drugs or Clothing

18.93 L / 5 gal

GENERAL INFORMATION

Alder	Choke Cherry	Elm	Sassafras
Aspen	Cottonwood	Maple	Sumac
Birch	Crateagus	Mulberry	Sweetgum
Buckeye	Hawthorn	Oak	Sycamore
Butternut	Dogwood	Persimmon	Tanbarkberry
Cherry	Elderberry	Pine	Tulip Poplar
Coconut	Hamamelis	Poplar	Willow
Cornus	Hickory	Sambucus	Winged Elm
Cypress	Juniper	Spicebush	
Deciduous	Dandelion	Plantain	Yarrow
Field Bindweed	Lambquarters	Smartweed	Wid. Lovage

DIRECTIONS FOR USE

FOLIAR APPLICATIONS

Apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs.

HIGH VOLUME SPRAYS WITH GROUND EQUIPMENT

Apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs.

Use Precautions should be used to reduce spray drift.

LOW VOLUME SPRAYS WITH GROUND EQUIPMENT

Apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs.

Woody Plant Control

Apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs.

Grassweed Control

Apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs.

AERIAL APPLICATION (Helicopter Only)

Apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs. For control of insects and diseases, apply to foliage of trees and shrubs. For control of weeds, apply to foliage of trees and shrubs.

BASAL BARK AND DORMANT BRUSH TREATMENTS

Apply to basal bark and dormant brush. For control of insects and diseases, apply to basal bark and dormant brush. For control of weeds, apply to basal bark and dormant brush. For control of insects and diseases, apply to basal bark and dormant brush. For control of weeds, apply to basal bark and dormant brush.

STORAGE AND DISPOSAL

Store in a cool, dry place. For disposal, follow local regulations. Do not use in areas where it is prohibited. For disposal, follow local regulations. Do not use in areas where it is prohibited.

USE PRECAUTIONS

Do not use in areas where it is prohibited. For disposal, follow local regulations. Do not use in areas where it is prohibited. For disposal, follow local regulations. Do not use in areas where it is prohibited. For disposal, follow local regulations.

NOTICE: This product is a restricted use pesticide. It is not to be used by anyone who is not a certified applicator or under their direct supervision. For more information, see the label.

SPECIMEN LAB
IBAC 1
REDUCED TO 25%

INSERT "B"

INSERT "C"

THE BOW CHEMICAL COMPANY
AND SUBSIDIARIES
MIDLAND, MICHIGAN 48640, USA HORGEN, SWITZERLAND HONG KONG
CORAL GABLES, FLORIDA 33134, USA SARNIA, ONTARIO, CANADA
* Trademark of THE BOW CHEMICAL COMPANY

GARLON* 4 PROPOSED LABEL

"INSERT A"

Black Medic	Goldenrod	Oxalis
Bull Thistle	Ground Ivy	Sowthistle
Canada Thistle	Lespedeza	Spiderwort
Catnip	Matchweed	Sweet Clover
Cinquefoil	Milkweed	Teasel
Clover	Milky Spurge	Wild Carrot (Queen Annes Lace)
Creeping Beggarweed	Mustard	Wild Parsnip
		Yarrow

"INSERT B"

Use GARLON 4 Herbicide at rates of 1 to 4 quarts in a total volume of 20 to 100 gallons per acre as a water spray mixture. Apply at any time weeds are actively growing. GARLON 4 Herbicide at 1/2 to 6 pints may be tank mixed with 1 to 2 quarts of DMA-4, FORMULA* 40, ESTERON* 99 Concentrate, TORDON* K, Or TORDON* 101 Mixture Herbicides to improve the spectrum of activity.

"INSERT C"

NOTE:

Conifer planted sooner than one month after treatment with GARLON 4 at less than 1 gallon per acre or sooner than two months after treatment at 1 to 2 gallons per acre may be injured. When tank mixtures of herbicides are used for forest site preparation, labels for all products in the mixture should be consulted and the longest recommended waiting period observed.

SPOT TREATMENT TO CONTROL CLUMPS OF RESPROUTING HARDWOODS SUCH AS BIG LEAF MAPLE USING A HOVERING HELICOPTER IN FORESTS.

STEM TREATMENT BEFORE LEAF-OUT. Mix 1 to 2 gallons of GARLON 4 Herbicide with about 20 gallons diesel oil and

"INSERT C" (cont'd)

enough water to make 100 gallons of solution. Apply as an invert emulsion by means of a hovering helicopter equipped with a nozzle system to direct sufficient spray to cover the stems to the ground line of the sprouted trees, usually 3/4 to 1 1/2 gallon per clump.

NOTE:

Conifers contacted by this spray may be seriously injured; in existing plantations, drift control systems, such as invert emulsions, should be used to minimize injury to adjacent conifers. A dye or other marking system to designate treated trees may be used.

"INSERT D"

Thinline Basal Bark Treatment: To control susceptible woody plants with stems less than 6 inches in diameter, apply undiluted GARLON 4 in a thin stream to all sides of the stems about 6 inches above the base of the plants. The stream should be directed horizontally to apply a narrow band of GARLON 4 around each stem or clump. From 2 to 15 ml. of chemical will be required for treatment of single stems and from 25 to 100 ml. to treat clumps of stems. Use an applicator metered or calibrated to deliver the small amounts required.

"INSERT E"

Avoid Injurious Spray Drift. Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible may seriously injure susceptible plants. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured. It is suggested that a continuous smoke

"INSERT E" (cont'd)

column at or near the spray site or a smoke generator on the spray equipment be used to detect air movement, lapse conditions, or temperature inversions (stable air). If the smoke layers or indicates a potential of hazardous spray drift, do not spray.