

**ACCEPTED**  
 FEB 12 1981  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act  
 EPA Reg. 2

**LOT**



# Dursban<sup>\*</sup> WB05

## insecticide

**Controls Numerous Pests in and Around Households**

**ACTIVE INGREDIENTS:**

Chlorpyrifos [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl)phosphorothioate]..... 0.50%  
 Xylene range aromatic solvent..... 0.33%

**INERT INGREDIENTS:**

99.17%

EPA Registration No 464-560

EPA Est. 464-MI-1

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

**Precautionary Statements  
 Hazards to Humans**

Avoid contact with eyes, skin and clothing • Avoid breathing vapors or spray mist • Keep away from food, feedstuffs and domestic water supplies  
 Wash thoroughly after handling

**Statement of Practical Treatment**

**If Swallowed:** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. **If on Skin:** In case of contact, remove contaminated clothing and immediately wash skin with plenty of soap and water. **If in Eyes:** Flush eyes with plenty of water.

**NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

**Physical or Chemical Hazards**

**Do Not Use or Store Near Heat or Open Flame  
 Do Not Store Below 32 Degrees F**

**Environmental Hazards**

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DURSBAN WB05 is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

In case of an emergency endangering life or property involving this product, call collect  
**517-638-4400**

**AGRICULTURAL CHEMICAL**  
 Do Not Ship or Store with Food,  
 Feeds, Drugs or Clothing

**PRECAUCION AL US**  
 ducto hasta que la et  
**TRANSLATION: (TO**  
 use this product untl

It is a violation of Fedr  
 with its labeling.

**General information**  
 pests indicated in the  
 undiluted spray (0.5%

Do not apply sprays o  
 and switch boxes or o  
 hazard. Do not use in pe  
 spray to contact food,  
 dishes and food handi  
 taminated by applicat  
 contact treated surfac  
 bowls (tanks) before se

**Application Within**  
 Private Residences in  
 Served) Including, Bu  
 Bakeries, Bottling Pl  
 treatment in and aro  
 cockroaches (including  
 crickets, firebrats, sil  
 confused flour beet  
 moths, rice weevils,  
 beetles. Apply as a co  
 areas. Do not introduce  
 food or food-contactin  
 rooms and closets; alo  
 stoves, refrigeration un  
 floor drains; and aroun  
 py to ant trails and whe  
 may encompass crack  
 material directly into op  
 equipment legs and ba  
 elements of constructi  
 capable of delivering a p  
 ment as needed but not  
 similar food service esta  
 other types of food ha  
 back), DURSBAN WB0  
 but emergency use sh  
**APPLICATION OF THE**  
**MENTS OTHER THAN**  
**TREATMENT ARE NOT**

86-1587 PRINTED IN U.S.A. IN JANUARY, 1981.  
 THIS IS THE INITIAL PRINTING.

**SPECIMEN LABEL**  
REDUCED TO 95%

**PRECAUCION AL USUARIO.** Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.  
**TRANSLATION: (TO THE USER)** If you cannot read English, do not use this product until the label has been fully explained to you.

**USE DIRECTIONS**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**PESTS INDOORS**

**General Information:** Use DURSBAN WB05 insecticide to control the pests indicated in the areas listed below by application of the product as an undiluted spray (0.5% chlorpyrifos). Repeat the treatment as needed.

Do not apply sprays of this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

**Application Within Food Handling Establishments (Places Other Than Private Residences in Which Food is Held, Processed, Prepared or Served) Including, But Not Limited to: Restaurants, Grocery Stores, Bakeries, Bottling Plants, Canned and Grain Mills:** Use as a spot treatment in and around both food and nonfood areas to control cockroaches (including strains resistant to certain other insecticides), ants, crickets, firebrats, silverfish, spiders, and stored product insects such as confused flour beetles, saw-toothed grain beetles, Indian meal moths, rice weevils, Mediterranean flour moths, and red flour beetles. Apply as a coarse, low pressure (20 psi or less) spray to localized areas. Do not introduce the spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners or rooms and closets, along and behind baseboards; beneath and behind stoves, refrigeration units and similar food preparator, areas and equipment; floor drains, and around plumbing and other utility installations. For ants apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and bases or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a pin stream of insecticide should be used. Repeat treatment as needed but not more often than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (callback), DURSBAN WB05 may be applied after 2 days from the last treatment, but emergency use should be limited to once per month.

**APPLICATION OF THIS PRODUCT IN FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A SPOT AND/OR CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.**

**Application Within Residential Buildings Including Homes and Apartment Buildings and Within Nonfood Areas of Industrial, Institutional, and Commercial Buildings Including Hospitals, Stores, Manufacturing Plants and Warehouses: For Spot Treatment Only -** Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets; cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators and cabinets, around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. **Note:** A period of 4 to 7 days is normally required for maximum effect on cockroaches.

**For control of carpet beetles,** thoroughly apply as a spray for spot treatment along baseboards and edges of carpeting, under carpeting, rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

**For control of fleas and brown dog ticks,** thoroughly apply as a spray for spot treatment to infested areas such as pet beds and resting quarters, nearby cracks and crevices; along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment of pet area. **DO NOT TREAT PETS WITH THIS PRODUCT.** To control the source of flea infestation, pets inhabiting the treated premises should be treated with a product registered for application to animals.

**PESTS OUT-OF-DOORS**

**Fleas, mosquitoes:** For outdoor use only as an aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects might or congregate. Repeat as necessary.

**Ants, brown dog ticks, crickets, fleas and earwigs:** For treatment of localized infestation of these insects in outdoor areas where there are weeds or brushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat applications as infestations warrant and as reinfestation occur.

**DISPOSAL**

Do not reuse empty container. Wrap container and put in trash collection.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

U.S. Patent No. 3,244,586

86-1581

181

**WB05**

de

**Around Households**

phosphorothioate]..... 0.50%  
..... 0.33%  
..... 99.17%

EPA Est. 464-MI-1

**CHILDREN**

**Physical or Chemical Hazards**

Use or Store Near Heat or Open Flame  
Do Not Store Below 32 Degrees F

**Environmental Hazards**

It is highly toxic to bees exposed to direct residues on plants. Protective information obtained from your Cooperative Agricultural Extension Service.

WB05 is toxic to fish, birds, and other animals out of lakes, streams, ponds, tidal marshes, etc. Shrimp and crabs may be killed at applications recommended on this label. Do not apply where important resources. Do not apply where runoff may occur. Do not apply when weather conditions are such that runoff may occur. Do not contaminate water with equipment or disposal of wastes.

**AGRICULTURAL CHEMICAL**

(Do Not Ship or Store with Food, Feeds, Drugs or Clothing)

**THE DOW CHEMICAL COMPANY**

AND SUBSIDIARIES

MIDLAND, MICHIGAN 48640, USA HORGEN, SWITZERLAND HONG KONG  
CORAL GABLES, FLORIDA 33134, USA SARNIA, ONTARIO, CANADA

\*Trademark of THE DOW CHEMICAL COMPANY

**ROSES, FLOWERS, ORNAMENTALS**

SECURITY Sevin-Maneb Spray will control the insects listed below on ornamental plants, roses, flowers, shrubs and shade trees and also will control diseases on the specific plants listed. Apply SECURITY Sevin-Maneb Spray at the rate of 2 1/2 tablespoons per gallon of water (1 gallon per 100 gallons of water) and spray thoroughly covering both upper and lower leaf surfaces. For insects, begin application when insects or their damage first appear and repeat at 7-10 day intervals or as necessary to maintain control. For disease follow suggested timing of application for each plant listed. The addition of 1/2 teaspoon of SECURITY Spreader-Sticker per gallon of water (2 oz. per 100 gallons) may improve residual insect control.

**INSECTS CONTROLLED**

- Ahfs
- Apple Aphid
- Armyworm
- Azalea Leafminer
- Bagworms
- Ch Leafminer
- Water Beetle
- Boxelder Bug
- Boxwood Leafminer
- Brownail Moth
- Cankerworms
- Catalpa Sphinx
- Chiggers
- Coolley Spruce Gail Aphid
- Cutworms
- Cypress Tip Moth
- Douglas Fir Tussock Moth
- Eastern Spruce Gail Aphid
- Elm Leaf Aphid
- Elm Leaf Beetle
- Elm Spanworm
- Firaphid Mites
- Green Pine Shoot Moth
- Fall Armyworm
- Flea Beetles
- Fuller Rose Beetle
- Gail Midges
- Gail Wasps
- Green Striped Mapleworm
- Grasshoppers
- Gypsy Moth
- Hackberry Nipple-Gale Maker
- Holly Bud Moth
- Holly Leafminer
- Jackpine Budworm
- Japanese Beetle
- Jeffery Pine Needleminer
- June Beetles
- Lace Bugs
- Leafhoppers
- Leafrollers
- Locust Borer
- Maple Leafcutter
- Mealy Bugs
- Mimosa Webworm
- Nantucket Pine Tip Moth
- Oak Leafminers
- Oak Leaf Skeletonizer
- Oakworm Complex
- Oleander Caterpillar
- Olive Ash Borer
- Orange Striped Oakworm
- Orange Tortrix
- Periodical Cicada
- Pine Sawfly
- Pine Spittlebug
- Pitch Pine Tip Moth
- Plant Bugs
- Poinsetta Hornworm
- Psyllids
- Puss Caterpillar
- Redhumped Oakworm
- Rose Aphid
- Rose Chafer
- Rose Slug
- Saddled Prominent Sawflies (exposed)
- Scale Insects
- Sowbugs
- Spiny Elm Caterpillar
- Springtails
- Spruce Budworm
- Spruce Needleminer
- Subtropical Pine Tip Moth
- Tent Caterpillars
- Thorn Bug

- Thrips (exposed)
- Ticks
- Walnut Caterpillar
- Webworms
- Western Hemlock Looper
- Western Spruce Budworm
- Willow Leaf Beetles
- Yellow Poplar weevil

**DISEASES CONTROLLED**

**ASTERS** To control RUST and STEM ROT, begin spraying before or when disease first appears and repeat at 7 to 10 day intervals.

**AZALEAS, CAMELLIAS** To control PETAL BLIGHTS, apply to litter and soil beneath plants, also to flowers of azaleas. Begin when flower buds swell. Repeat at 3 to 5 day intervals.

**CARNATIONS, PANSIES** To control ALTERNARIA BLIGHT or LEAF SPOT, ANTHRACNOSE and BOTRYTIS BLIGHT, begin spraying when new growth starts and repeat weekly.

**DAHLIAS, IRIS, ZINNIAS** To control ALTERNARIA BLIGHT or LEAF SPOT, and BOTRYTIS BLIGHT, spray when new growth starts and repeat weekly.

**DOGWOOD** To control ANTHRACNOSE Spray tree thoroughly. Begin when buds open. Repeat when bracts fall, 4 weeks later and in late summer.

**GERANIUMS, HYDRANGEAS, LILACS** To control BOTRYTIS BLIGHT, begin sprays when new growth starts, before disease appears, or at first sign of disease. Repeat at 7 to 10 day intervals.

**GLADIOLUS** To control BOTRYTIS BLIGHT, CURVULARIA and STEMPHYLIUM LEAF SPOTS, begin spraying when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.

**PEONIES** To control ALTERNARIA LEAF BLIGHT or SPOT, BOTRYTIS BLIGHT and PHYTOPHTHOA BLIGHT, apply spray to foliage and soil in early spring and early fall and at 7 to 14 day intervals during growing season.

**ROSES** To control BLACK SPOT, CERCOSPORA LEAF SPOT (Texas) and RUST (California), begin spraying when first leaves unfold and repeat on 7 to 10 day intervals through season.

**SNAPDRAGONS** To control BOTRYTIS BLIGHT, DOWNY MILDEW and RUST, apply sprays at time of emergence and repeat at 7 to 10 day intervals.

**DISPOSAL**

Do not reuse empty container. Wrap container and put in trash collection.

**NOTICE:** The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label. 086121

**MANUFACTURERS**  
**WOOLFOLK CHEMICAL WORKS, INC.**  
**Fort Valley, Georgia 31030**

**SECURITY**  
**Sevin-Maneb Spray**

**Controls Insects & Diseases on Vegetables, Lawns, Roses, Flowers and Ornamentals.**

**ACTIVE INGREDIENTS:**

- Carbaryl (1-naphthyl methylcarbamate) . . . . . 11.10%
- Maneb (Manganese ethylene bisdithiocarbamate) . . . . . 13.00%
- (Total Manganese as metallic . . . . . 2.68%)

**INERT INGREDIENTS: . . . . . 75.90%**

EPA Reg. No. 769-499 EPA Est. 769-GA-1 Total 100.00%  
**SEVIN** is a registered trademark of Union Carbide Corporation for carbaryl insecticide.

This product contains 1 pound Carbaryl and 1 1/2 pounds Maneb per gallon.

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

**IF INHALED:** Remove victim to fresh air.

**IF ON SKIN:** Wash with soap and water. Remove any contaminated clothing.

**IF IN EYES:** Flush eyes with water.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS: HARMFUL IF SWALLOWED OR INHALED.** May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention. Wear regular long-sleeved work clothing while spraying. Wash hands and face before eating. Wash thoroughly after handling. **NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. **DO NOT USE 2-PAM, OPIATES OR CHOLINESTERASE INHIBITING DRUGS.**

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

**PROTECT BEES:** This product is highly toxic to bees exposed to direct treatment or to residues remaining on treated areas. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

It is a violation of the FIFRA Act to apply this product in a manner inconsistent with the label.

To insure proper use, please read the label carefully. Do not pour any low pressure hose spray. For HOSE application, be sure to dilute to the label strength. Be sure to use the recommended filling procedure during prevention.

Apply in the recommended amount. Do not exceed the recommended interval. Do not apply more than the recommended number of times. Do not apply if insects are present.

Apply to the following plants: Alliums, Armyworm, Asparagus, Aster, Beans, Blister Beetles, China, Corn, Corn, Cow, Cucumbers, Eggplant, Fall, Flea, Grass, Greenhouse.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**MIXING INSTRUCTIONS**

To insure a uniform product, gently shake contents before opening. SECURITY Sevin-Maneb Spray may be applied with any low pressure sprayer such as knap-sack type, compressed air tank-type, hose-on or other similar lawn or garden spray equipment.

For HOSE-ON sprayers, it is recommended that this product be diluted 1 part product to 1 part water before filling applicator bottle. This will require setting the hose sprayer to apply twice the recommended label amount. (Example: If the label states apply 3 1/2 tablespoons per gallon, the sprayer should be set to apply 7 tablespoons.) Check sprayer to see that it delivers the proper amount of material before using. Be sure the sprayer tank and spray tip or screens are clean before use. For sprayers other than hose-on, fill the spray tank with 1/2 to 3/4 the desired amount of clean water. Pour the recommended amount of product into the tank and complete filling to desired level with water. Before and frequently during use, shake or agitate the spray mixture in the tank to prevent settling. It is recommended that spray equipment be thoroughly cleaned after each day's use.

**CONVERSION TABLE**

1 ounce = 2 tablespoons = 6 teaspoons

**VEGETABLES**

Apply in sufficient volume of water to provide good coverage. In the garden, approximately one gallon of diluted spray will be required per 100 feet of row. This is equivalent to 150 gallons per acre. Direct spray to cover both upper and lower leaf surfaces. Apply this product at the rate of 2 gallons per acre or 3 1/2 tablespoons per 1 gallon of water. Follow the time interval between last application and harvest indicated by the number in ( ) following each crop. A (0) indicates this product may be applied up to and including the day of harvest. For insect control, begin application when insects or their damage first appear and re-treat as necessary (except post-harvest on asparagus only). For disease control, follow the specific directions for each crop. Do not feed forage or hay from treated crops to livestock.

**INSECT CONTROL**

- |                        |                       |
|------------------------|-----------------------|
| Alfalfa Caterpillar    | Harlequin Bug         |
| Armyworm               | Imported Cabbageworm  |
| Asparagus Beetle       | Japanese Beetle       |
| Aster Leafhopper       | Lace Bug              |
| Bean Leaf Beetle       | Leathopper            |
| Blister Beetle         | Limabean Pod Borer    |
| Chinch Bug             | Lygus Bug             |
| Colorado Potato Beetle | Melonworm             |
| Corn Earworm           | Mexican Bean Beetle   |
| Corn Rootworm          | Pea Leaf Weevil       |
| Cowpea Curculio        | Pea Weevil            |
| Cucumber Beetle        | Red-Necked Peanutworm |
| Cutworms               | Sap Sucker            |
| European Corn Borer    | Southern Corn Borer   |
| Fall Armyworm          | Spittlebug            |
| Flea Beetle            | Squash Bug            |
| Grasshopper            | Stink Bug             |
| Green Cloverworm       | Tarnished Plant Bug   |

- Three Corned Alfalfa Hopper
- Thrips
- Tomato Fruitworm
- Tomato Hornworm

- Tomato Pinworm
- Velveteen Caterpillar
- Western Bean Cutworm
- Yellow Striped Armyworms

**DISEASE CONTROL**

**ASPARAGUS** (Post Harvest) To control RUST, apply immediately after harvest. Repeat at 10 day intervals as long as necessary. In young fields that will not be harvested, begin when disease appears and repeat at 10 day intervals.

**BEANS** (Lima, Snap or String) (4) To control ANTHRACNOSE, DOWNY MILDEW and RUST, begin spraying when plants begin to bloom or when disease first appears. Repeat applications at 4 to 7 day intervals.

**BROCCOLI** (3) **BRUSSELS SPROUTS** (3) **CABBAGE** (7) **CAULIFLOWER** (3) **KOHLRABI** (3) To control ALTERNARIA LEAF SPOT and DOWNY MILDEW, apply to transplants in the garden or field as soon as disease threatens and repeat weekly as needed (every 3 to 5 days under severe conditions). Add 1/2 teaspoon SECURITY Spreader-Sticker to each gallon of spray mixture. Remove residues at harvest by washing or trimming.

**CARROTS** (0) To control ALTERNARIA and CERCOSPORA LEAF SPOTS, begin spraying when plants are 6 weeks old and repeat at 7 to 10 day intervals.

**COLLARDS** (14) **KALE** (14) **MUSTARD GREENS** (14) **TURNIPS** (14) To control ALTERNARIA LEAF SPOT and DOWNY MILDEW, begin application when disease first appears in area and repeat at 7 to 10 day intervals. Remove excess residues at harvest by washing.

**CORN, SWEET** (0) To control HELMINTHOSPORIUM BLIGHT, begin applications when disease first threatens and repeat at 4 to 7 day intervals until harvest.

**CUCUMBERS** (5) **SQUASH, SUMMER and WINTER** (5) To control ANTHRACNOSE, DOWNY MILDEW, ALTERNARIA (Macrosporium), LEAF SPOT and ANGULAR LEAF SPOT on cucumbers, and PYTHIUM FRUIT ROT, begin applications when runners form and repeat weekly as needed.

**EGGPLANT** (0) **PEPPERS** (0) To control ANTHRACNOSE, EARLY BLIGHT (Alternaria) on eggplant, CERCOSPORA LEAF SPOT (frog-eye spot) on peppers and FRUIT ROT, apply as fruit begins to form and repeat at weekly intervals through harvest.

**LETTUCE** (Head-10, Leaf-14), **ENDIVE** (14) To control DOWNY MILDEW, apply when disease first appears and repeat at 7 to 10 day intervals (3 to 5 days under severe conditions). Remove residues at harvest from head lettuce by stripping and trimming and from leaf lettuce and endive by washing or other effective means.

**MELONS** (Cantaloupe, Cassabas, Crenshaws, Honeydews, Honey Balls, Muskmelons, Persian Melons and Watermelons) (5) To control ANTHRACNOSE, CERCOSPORA LEAF SPOT, DOWNY MILDEW and GUMMY STEM BLIGHT, begin applications when runners begin to form or when disease threatens. Repeat every 7 to 10 days as needed. Under severe conditions, shorten intervals to 3 to 5 days.

**PEAS** (Black-eyed peas, Cowpeas) (4) To control DOWNY MILDEW, begin at first sign of disease and repeat at 7 to 10 day intervals.

**POTATOES** (0) To control EARLY and LATE BLIGHT, apply every 7 to 10 days throughout the season.

**PUMPKINS** (5) To control ANGULAR LEAF SPOT and DOWNY MILDEW, begin applications when disease threatens and repeat at 7 to 10 day intervals as needed.

**SPINACH** (14) To control DOWNY MILDEW (blue mold) and WHITE RUST, begin applications when disease first appears in area and repeat at 7 to 10 day intervals. Remove residues at harvest by washing or other effective means.

**TOMATOES** (5) To control ANTHRACNOSE, EARLY and LATE BLIGHT, GRAY LEAF SPOT (Stemphylium) and SEPTORIA LEAF SPOT, begin applications when first cluster fruit is well formed and repeat at 7 to 10 day intervals. More frequent applications may be necessary starting when plants are set in fields if weather conditions favor disease. To avoid injury, do not use on young seedlings in greenhouses.

**LAWNS**

Home Lawns, Recreational and Commercial (non-grazed) turf Areas. SECURITY Sevin-Maneb Spray will control the insects and diseases listed. Use a sufficient amount of water for thorough coverage. For best results, mow lawn before application. Do not graze treated areas. Do not feed clippings to livestock.

**INSECTS CONTROLLED:** Apply at first sign of insects or their damage and repeat as necessary.

- |                   |                         |
|-------------------|-------------------------|
| Ants              | Grasshoppers            |
| Armyworms         | June Beetles            |
| Bluegrass Billbug | Leafhoppers             |
| Centipedes        | Lucerne Moth            |
| Chiggers          | Milipedes               |
| Cutworms          | Mosquitoes              |
| Earwigs           | Sod Webworm             |
| Essex Skipper     | (lawn moths)            |
| European Chater   | Sowbugs                 |
| Fall Armyworm     | Springtails             |
| Fiery Skipper     | Ticks                   |
| Fleas             | Yellow Striped Armyworm |

**DISEASE CONTROLLED:** To control BROWN PATCH (Helminthosporium Blight and Melting Out), begin at first sign of disease and repeat at 7 to 10 day intervals. To control LEAF, STEM and STRIPE RUSTS, begin when rust pustules are first seen and repeat at 7 to 14 day intervals.

**AMOUNT TO USE**

AREA TO TREAT	AMOUNT TO USE
500 sq. ft.	13 oz.
1,000 sq. ft.	26 Oz.
2,500 sq. ft.	to 3 quarts
5,000 sq. ft.	1 1/2 gallons

ADDITIONAL DIRECTIONS FOR USE ON THE OTHER SIDE PANEL. FEB 12 1981

NET CONTENTS  
ONE GALLON

769-494

32

es, Lawns, Roses,

11.10%  
13.00%

75.90%  
Total 100.00%

Corporation for carbaryl

12 pounds Maneb per

**TREATMENT**

er and induce vomiting by  
not induce vomiting or give  
let medical attention.

Remove any contaminated

**CHILDREN.**

**N**

**STATEMENTS**

**ANIMALS: HARMFUL IF**  
injection of eyes, nose, throat and  
contact with skin, eyes or  
enty of water. For eyes, get  
sweated work clothing while  
ating. Wash thoroughly after  
is a moderate, reversible  
dotal. DO NOT USE 2-PAM,

**DRUGS.**

**HAZARDS**

akes, streams or ponds. Do not  
st apply when weather conditions  
contaminate water by cleaning of  
this product only as specified on

y toxic to bees exposed to direct  
reated areas. Do not apply when  
crop, or weeds blooming in the  
e timed to provide the maximum  
ne next period of bee activity.

31