

PM 25
464-576

Specimen Label

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Access*

Herbicide

ACCEPTED
JUL 21 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 464-576

*
ACCESS

Active Ingredient(s):
Picloram (4-amino-3,5,6-trichloropicolinic acid), isooctyl esters 17.1%
Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid), Butoxyethyl Ester 32.5%
Inert Ingredients 50.4%
Acid Equivalents:
Picloram - 11.7%, 1 lb/gal
Triclopyr - 23.4%, 2 lb/gal
Contains petroleum distillates.
E.P.A. Registration No. 464-576
E.P.A. Est. 464-MI-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

AVISO:

PRECAUCION AL USUARIO:
Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals
HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN • CAUSES EYE IRRITATION

Avoid Contact With Skin, Eyes, Or Clothing
• Wash Thoroughly With Soap And Water After Handling And Before Eating Or Smoking

STATEMENT OF PRACTICAL TREATMENT: In case of contact, flush skin or eyes with plenty of water. Get medical attention if irritation persists. Remove and wash contaminated clothing before reuse. If swallowed, do not induce vomiting. Call a physician.

Physical or Chemical Hazards

COMBUSTIBLE • Do Not Use or Store Near Heat or Open Flame. • Do Not Cut or Weld Container.

Environmental Hazards

This pesticide is toxic to fish • Do not apply directly to any body of water • Do not contaminate water used for irrigation or domestic purposes by cleaning of equipment or disposal of wastes • Do not allow run-off or spray to contaminate wells, irrigation ditches or any body of water used for irrigation or domestic purposes • Do not make application when circumstances favor

movement from treatment site •

Picloram is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation and drinking purposes. Users are advised not to apply picloram where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

NOTICE

Read the entire label. Use only according to label directions. Before buying or using this product, read "WARRANTY LIMITATIONS AND DISCLAIMER" elsewhere on this label. If terms are not acceptable, return unopened package at once to seller for full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under "WARRANTY LIMITATIONS AND DISCLAIMER."

IN CASE OF AN EMERGENCY

endangering life or property involving this product, call collect 517-636-4400
AGRICULTURAL CHEMICAL
Do Not Ship or Store with Food, Feeds, Drugs, or Clothing

BEST AVAILABLE COPY

2 of 5

GENERAL INFORMATION

ACCESS Herbicide is recommended for the control of unwanted woody plants in forests, rights-of-way such as electrical power lines, communication lines, pipelines, roadsides, and railroads and on non-crop areas including industrial manufacturing and storage sites and fence rows.

Woody plant species controlled include:

Ash	Elm	Maple	Poplar
Aspen	Hackberry	Oak	Oceanspray
Birch	Hickory	Pine	Tanoak
Cherry	Locust	Sassafras	Multiflora rose

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

~~BASAL BARK TREATMENT~~

In rights-of-way, other non-crop areas and forests, use 1 to 2 gallons of ACCESS Herbicide in enough diesel oil, No. 1 or No. 2 fuel oil or kerosene to make 100 gallons of spray mixture. Substitute other diluents or oils only in accordance with the manufacturer's recommendations. To obtain the best control of hard-to-control species, such as ash and elm, or trees with diameters larger than 4 inches, use the higher recommended rate. Basal treatments are less effective on trees with diameters larger than 6 to 8 inches. Apply with knapsack sprayer or power spraying equipment using low pressure (20-40 psi). Spray the basal parts of brush and tree trunks to a height of 12-15 inches from the ground. Thorough wetting of the indicated area including the root collar is necessary for good control. Spray until run-off at the ground line is noticeable. For root suckering species such as aspen, sassafras and locust, also spray the ground under the plants to cover small root suckers which may not be visible above the soil surface. Rough bark requires more spray than smooth bark. Apply at any time, including the winter months, except when snow or water prevent spraying to the ground line. Best control will be obtained when applications are made during the late dormant or active growing season.

For oil-water mixture application, mix 2 to 3 gallons ACCESS Herbicide/22 to 23 gallons of oil and 1 gallon of Sponto 712, or equivalent, and add to 74 gallons water as directed below. Use the application procedures specified above. For best results with oil-water mixtures, treat only stems 2 inches or less in diameter.

Insert A → ~~Mixing Oil Sprays: Use diesel oil, No. 1 or No. 2 fuel oil or kerosene. Use other oils or diluents only as recommended by their manufacturer. Add ACCESS Herbicide to the required amount of oil in the spray tank or mixing tank and mix thor-~~

oughly. If this mixture stands over 4 hours, reagitiation is required.

~~Mixing Oil-Water Sprays: First, premix the ACCESS Herbicide, oil and surfactant in a separate container. Do not allow any mixtures containing water to get into the ACCESS Herbicide or the premix. Fill the spray tank about half full with water, then slowly add the premix with continuous agitation and complete filling the tank with water. Continue moderate agitation.~~

Note: If the premix is put in the tank without any water, the first water added may form a thick "invert" (water in oil) emulsion which will be hard to break.

USE PRECAUTIONS

Apply this product only as specified on this label.

Do Not Make Application When Circumstances Favor Movement From Treatment Site.

Do not apply or otherwise permit ACCESS Herbicide or its spray mist to contact soil used to grow desirable, susceptible plants. Do not allow ACCESS Herbicide to contact susceptible plants such as grapes, tobacco, vegetable crops, flowers or other desirable broadleaf plants. Coarse sprays are less likely to drift. Do not apply with hollow cone-type insecticide nozzle or other nozzles that normally produce a fine-droplet spray.

Do Not Contaminate Water Intended for Irrigation or Domestic Purposes. To avoid injury to crops or other desirable plants, do not treat or allow spray drift or run-off to fall onto banks or bottoms of irrigation ditches, either dry or containing water, or other channels that carry water that may be used for irrigation or domestic purposes. Do not apply to snow or frozen ground.

Water contaminated with residues of picloram from drainage ditch bank uses shall not be used to irrigate crops which are not registered for use with this chemical.

Do not rotate food or feed crops on treated land if they are not registered for use with picloram until an adequately sensitive bioassay or chemical test shows that no detectable picloram is present in the soil.

Do not move treated soil to other areas or use it to grow plants if they are not registered for use with picloram until an adequate sensitive bioassay or chemical test shows that no detectable picloram is present in the soil.

Untreated trees occasionally are affected by root uptake of herbicide leached into the soil or excreted by the roots of nearby treated trees. If such injury cannot be tolerated, do not use ACCESS Herbicide close to desirable trees.

Insert C

BEST AVAILABLE COPY

3 of 5

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

STORAGE: Store above 28 F or until before use

DISPOSAL:

Prohibitions: Open dumping is prohibited

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Do not reuse container. Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General: Consult federal, state or local disposal authorities for approved alternative procedures.

Be sure that use of this product conforms to all applicable regulations.

WARRANTY LIMITATIONS AND DISCLAIMER

The Dow Chemical Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions therein under normal conditions of use. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify in writing The Dow Chemical Company of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of The Dow Chemical Company or any other seller will be one of the following, at the election of The Dow Chemical Company:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations And Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations And Disclaimer in any manner.

01780-L1

187



THE DOW CHEMICAL COMPANY
Midland, Michigan 48674 U.S.A.

*Trademark of THE DOW CHEMICAL COMPANY

BEST AVAILABLE COPY

4 of 5
ACCESS Herbicide
EPA Reg. No. 464-576
Proposed Label Revision
February 10, 1987

Insert A:

Mixing Instructions:

General Information: In forests, rights-of-way, and other non-crop areas, use ACCESS Herbicide in oil or oil-water spray mixtures prepared as described below. When preparing mixtures, use as oils either diesel fuel, No. 1 or No. 2 fuel oil, or kerosene. Substitute other oils or diluents only as recommended by the oil or diluent's manufacturer.

Insert B:

Mixing Oil-Water Sprays: Premix 2 to 3 gallons of ACCESS Herbicide, 22 to 23 gallons of oil, and 1 gallon of Sponto 712, or equivalent. Do not allow any mixtures containing water to get into the ACCESS Herbicides or the premix. Add the premix to 74 gallons of water as follows. Fill the spray tank with half the amount of required water, then slowly add the premix with continuous agitation and add the remainder of the required water. Continue moderate agitation.

BEST AVAILABLE COPY

5 of 5
Insert C:

Application Instructions:

Basal Bark Treatment: Use 1 to 2 gallons of ACCESS Herbicide in enough oil to make 100 gallons of spray mixture. Herbicide concentration should vary with size and susceptibility of the species treated and season of application. Basal treatments are less effective on trees with basal diameters larger than 6 inches. Apply with knapsack or backpack sprayer or power spraying equipment using low pressure at the spray nozzle (20 to 40 psi). Spray the basal parts of brush and tree stems thoroughly wetting the indicated area, including the root collar, for good control. Spray until run-off at the ground line is noticeable. For root suckering species, such as aspen, sassafras, and locust, also spray the ground under the plants to cover small root suckers which may not be visible above the soil surface. Rough bark requires more spray than smooth bark. Apply at any time, including the winter months, except when snow or water prevent spraying to the ground line. Best control will be obtained when applications are made during the late dormant or active growing season.

Alternatively, the oil-water mixture described above may be applied using the application procedures including winter month treatment, specified in the preceding paragraph. For best results with oil-water mixtures, treat only stems less than 2 inches in basal diameter.

Low Volume Basal Bark Treatment: To control susceptible woody plants with stems less than 6 inches in basal diameter, mix 20 to 30 gallons of ACCESS Herbicide in enough oil to make 100 gallons of spray mixture. (This is equivalent to 25.6 to 38.4 fluid ounces of ACCESS Herbicide per gallon of spray mix.) Apply with a backpack or knapsack sprayer using low pressure and a solid cone or flat fan nozzle. Spray the basal parts of brush and tree stems in a manner which thoroughly wets the lower stem, including the root collar area, but not to the point of runoff. Herbicide concentration should vary with size and susceptibility of the species treated and season of application. Apply at any time, including the winter months, except when snow or ice prevent spraying to the ground line.

Thinline Basal Bark Treatment: To control susceptible woody plants with stems less than 6 inches in basal diameter, apply undiluted ACCESS Herbicide in a thin stream to all sides of the lower stems. The stream should be directed horizontally to apply a narrow band of ACCESS Herbicide around each stem or clump. From 2 to 15 ml of chemical will be required for treatment of single stems. Use application equipment metered or calibrated to deliver the small amounts required.

BEST AVAILABLE COPY