

PLEASE NOTE, WHERE THIS LABEL HAS DURSBAN 2E, THE LABEL FOR THE PRODUCT SUBMITTED FOR REGISTRATION SHOULD HAVE DURSBAN 2EC



SPECIMEN LABEL
REDUCED TO 70%

Dursban 2EC insecticide

ACCEPTED
with COMMENTS
by EPA

DEC 1 1982

1107-576

Xylene range aromatic solvent

Controls Numerous Pests of Households, Turf, and Ornamental Plants. Also Recommended for Mosquito Control and Area Control of Ticks and Chiggers.

To Be Applied Only By or Under the Supervision of Pest Control Operators, Mosquito Abatement Districts, and Other Trained Personnel Responsible for Mosquito Control.

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0 diethyl O (3,5,6 trichloro 2 pyridyl) phosphorothioate]
Aromatic petroleum germicide solvent

INERT INGREDIENTS

Contains 2 pounds of chlorpyrifos per gallon
E P A Registration No 464 343

E P A Est 464 MI 1

PRECAUCION AL USUARIO: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
TRANSLATION (TO THE USER): If you cannot read English, do not use this product until the label has been fully explained to you.

24.1%
59.4
15.6

..... **WAR**

MAY BE FATAL IF SWALLOWED • MAY BE INJURIOUS IF INHALED • SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

In case of an emergency endangering life or property involving this product, call collect 517-636-4400

18.93 L / 5 gal

86-1176 PRINTED IN U.S.A. IN MARCH, 1982.
REPLACES SPECIMEN LABEL 86-1176 PRINTED IN AUGUST, 1981
DISCARD PREVIOUS SPECIMEN LABELS.
REVISIONS INCLUDE: (1) ADDED CONTROL OF MOSQUITO LARVAE IN WASTE WATER, (2) ADDED CONTROL OF SOUTHERN MASKED CHAFER LARVAE, (3) CHANGED "KEEP OUT OF LAKES, STREAMS, AND PONDS" TO "DO NOT APPLY DIRECTLY TO WATER".

an*

cide

for
iggers.

Public Health Organizations,
for Insect Control Programs.

ACH OF CHILDREN

WARNING
MAY BE ABSORBED THROUGH SKIN
TO EYES AND SKIN
ADDITIONAL PRECAUTIONS

AGRICULTURAL CHEMICAL
Do Not Ship or Store with Food, Feeds,
Drugs or Clothing

Triple rinse (or equivalent) and offer for recycling or reconditioning, or dispose of in a sanitary landfill or by incineration if permitted by State and local authorities.

KEEP OUT OF REACH OF CHILDREN

WARNING

PRECAUTIONARY STATEMENTS

Hazards to Humans

**MAY BE FATAL IF SWALLOWED
MAY BE ABSORBED THROUGH SKIN
MAY BE INJURIOUS TO EYES AND SKIN**

Do Not Take Internally • Do Not Get in Eyes or on Skin • Wash Thoroughly After Handling • Wash Contaminated Clothing Before Reuse • Flush Contaminated Eyes with Plenty of Water and Get Prompt Medical Attention • Avoid Breathing Vapors and Spray Mist • Avoid Spraying Food Crops • Keep Away from Food, Feedstuffs, and Domestic Water Supplies • Keep Container Closed • Handle Concentrate in a Ventilated Area

Statement of Practical Treatment

If Swallowed: Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician immediately. **If On Skin:** In case of contact, remove contaminated clothing and immediately wash skin with soap and water. **If In Eyes:** Flush eyes with plenty of water for 15 minutes. Call a physician.

Note To Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

Physical or Chemical Hazards

COMBUSTIBLE

**Do Not Use or Store Near Heat or Open Flame
Do Not Cut or Weld Container**

Environmental Hazards

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service. DURSBAN 2E is toxic to fish, birds, and other wildlife. Do not apply directly to water. Shrimp and crabs may be killed at application rates recommended on the label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ AND FOLLOW COMPLETE DIRECTIONS FOR
USE AND OTHER IMPORTANT INFORMATION
GIVEN IN BOOKLET ATTACHED TO PRODUCT CONTAINER**

STORAGE AND DISPOSAL

Do not store in uncleanable water. Do not feed by, storage or disposal. Do not store in plastic spray mixture or residue that cannot be used or fully reprocessed. Do not store in plastic spray mixture or residue that cannot be used or fully reprocessed. Do not store in plastic spray mixture or residue that cannot be used or fully reprocessed.

Do not store in plastic spray mixture or residue that cannot be used or fully reprocessed.

Do not store in plastic spray mixture or residue that cannot be used or fully reprocessed.

Do not store in plastic spray mixture or residue that cannot be used or fully reprocessed.

THE DOW CHEM

AND SUBS

MIDLAND MICHIGAN 48640 USA HC
CORAL GABLES FLORIDA 33134, U

Trademark of THE DOW



RO INSECTICIDE
LIQUID N.O.S.
(Chlorpyrifos)
NA 1993
F.P. - 101°F

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

U.S. Patent No. 3,244,586

10685-060-8

26005-F282

ICAL COMPANY

DIARIES
RGEN, SWITZERLAND HONG KONG
SA SARNIA, ONTARIO, CANADA
CHEMICAL COMPANY

SL1210

DURSBAN 2E INSECTICIDE

Controls Numerous Pests of Households, Turf and Ornamental Plants. Also Recommended for Mosquito Control and Area Control of Ticks and Chiggers.

To Be Applied Only by or Under the Supervision of Pest Control Operators, Public Health Organizations, Mosquito Abatement Districts, and Other Trained Personnel Responsible for Insect Control Programs

E.P.A. Registration No. 464-343



Xylene range aromatic solvent

Controls Numerous Pests of Households, Turf and Ornamental Plants. Also Recommended for Mosquito Control and Area Control of Ticks and Chiggers.

USE DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PESTS INDOORS

General Information: Use DURSBAN 2E insecticide to control the pests indicated in the areas listed below by application as a 0.25% or 0.5% chlorpyrifos spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute DURSBAN 2E with water or with a suitable base oil as shown in the following SPRAY DILUTION CHART. Repeat the treatment as needed.

Use only deodorized kerosene for oil-based sprays to be applied in food handling establishments.

NOTE: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrins may be added to the spray mixture containing DURSBAN 2E insecticide. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays; use pyrethrins at 0.25% in water-base and 0.05% in oil-base sprays. Consult labeling for the dichlorvos or pyrethrins product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products. Do not use DURSBAN 2E in combination with dichlorvos (DDVP) or pyrethrins for treatment of food areas of food handling establishments.

Mixture of DURSBAN 2EC and dichlorvos (DDVP) should not be sprayed on carpets.

PESTS OUT-OF-DOORS

General Information: Use DURSBAN 2E insecticide for control of certain pests on outside surfaces of buildings and on lawns and ornamental plants, for area control of ticks and chiggers and for mosquito control. Consult the following directions for dosage and application details to control the pests listed.

Pests on Outside Surfaces of Buildings: Use DURSBAN 2E at the rate of 1/2 fluid ounces per one gallon of water (0.5% chlorpyrifos emulsion) to control ants, clover mites, cockroaches, crickets, earwigs, fleas, flies, millipedes, mosquitos, sowbugs, spiders and ticks by application as a residual spray or with a paint brush to outside surfaces of buildings including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Do not treat or allow the spray to contact plants (lawns, flowering plants, shrubs, trees, evergreens, etc.) because of possible injury. Repeat treatment as needed to maintain effectiveness.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT

PRECAUCION Al Usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada en español.

TRANSLATION: (TO THE USER) If you cannot read English, do not use this product until the label has been fully explained to you.

In case of an emergency endangering life or property involving this product, call collect 517-636-4400

AGRICULTURAL CHEMICAL
Do Not Ship or Store with Food, Feeds, Drugs or Clothing

ACTIVE INGREDIENTS:

Chlorpyrifos (0,0-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate)	24.1%
Aromatic petroleum derivative solvent	59.4
INERT INGREDIENTS	15.6

Contains 2 pounds of chlorpyrifos per gallon
E.P.A. Registration No. 464-343
E.P.A. Est. 464-88-1

CONTENTS

PRECAUTIONARY STATEMENTS	3
Hazards to Humans	3
Statements of Practical Hazards	3
Physical or Chemical Hazards	3
Environmental Hazards	4
PESTS INDOORS	5
General Information	5
Spray Dilution Chart	6
Application Within	6
Food Handling Establishments	7
Residential Buildings	8
Control of Carpet Beetles	9
Control of Brown Dog Ticks	9
PESTS OUT-OF-DOORS	10
General Information	10
Outside Surfaces of Buildings	10
Lawns, Ornamental and Turf Areas	11
Pests of Ornamentals	12
Wood Boring Insects	12
Use on Turf Charts	13-15
Peach Tree Borer	16
Native Elm Bark Beetle	16
Area Control of Ticks and Chiggers	17
Imported Fire Ants	17
Use Charts for Ornamental	19-20
Mosquito Control	21-24
STORAGE AND DISPOSAL	25
NOTICE	25

Application Within Food Handling Establishments (Places Other Than Private Residences in Which Food is Held, Processed, Prepared or Served) Including, But Not Limited to, Restaurants, Grocery Stores, Bakeries, Butchering Plants, Canneries and Grain Mills: Use as a spot treatment in and around both food and nonfood areas to control cockroaches (including strains resistant to other insecticides), ants, crickets, firebrats, silverfish, spiders and stored product insects such as confused flour beetles, saw-toothed grain beetles, Indian meal moths, rice weevils, Mediterranean flour moths and red flour beetles. Apply as a coarse, low pressure (20 psi or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated using special care to avoid unnecessary runoff. Do not introduce the spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners of rooms and closets, along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains, and around plumbing and other utility installations. For ants apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and bases or which occur at points between different elements of construction or between

NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, millipedes, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 1 pint of DURSBAN 2E per 100 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1000 sq. ft. to thoroughly and uniformly wet the band area.

Pests of Lawns and other Ornamental and Recreational Turf Grass Areas: Use DURSBAN 2E to control pests listed in the following table by application at the recommended dosages and in accordance with the directions given below. Dilute DURSBAN 2E in enough water to obtain complete and uniform coverage of pest infested area; and apply as a coarse, low pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf except as noted. For best results, the lawn should be moist at time of treatment.

Pests of Ornamentals: Use DURSBAN 2E insecticide to treat flowers, shrubs, vines, shade and flowering trees, and evergreens found to be infested with pests listed in the following table. Dilute DURSBAN 2E with water according to directions given in the table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage. For best results apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 days intervals if needed. **NOTE:** Do not use on azaleas, camelias, poinsettias, rose bushes, or variegated ivy because of possible injury to these plants.

Wood Boring Insects Infesting Shrubs and Trees such as Dogwood Borer, Lidic Borer, Ash Borer and Rhododendron Borer: Apply DURSBAN 2E to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat in your area. Mix at the rate of two quarts of DURSBAN 2E in 100 gallons of water and apply uniformly as a coarse, low pressure spray.

KEEP OUT OF REACH OF CHILDREN

WARNING

PRECAUTIONARY STATEMENTS

Hazards to Humans

MAY BE FATAL IF SWALLOWED
MAY BE ABSORBED THROUGH SKIN
MAY BE INJURIOUS TO EYES AND SKIN
Do Not Take Internally • Do Not Get in Eyes or on Skin • Wash Thoroughly After Handling • Wash Contaminated Clothing Before Reuse • Flush Contaminated Eyes with Plenty of Water and Get Prompt Medical Attention • Avoid Breathing Vapors and Spray Mist • Avoid Spraying Food Crops • Keep Away from Food, Feedstuffs, and Domestic Water Supplies • Keep Container Closed • Handle Concentrate in a Ventilated Area

Statement of Practical Treatment
If Swallowed: Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician immediately. **If On Skin:** In case of contact, remove contaminated clothing and immediately wash skin with soap and water. **If In Eyes:** Flush eyes with plenty of water for 15 minutes. Call a physician. **Note to Physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

Physical or Chemical Hazards

COMBUSTIBLE

Do Not Use or Store Near Heat or Open Flame
Do Not Cut or Weld Container

equipment and floors. Equipment capable of delivering a pinstream of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back) DURSBAN 2E may be applied after 2 days from last treatment but emergency use should be limited to once per month.

Applications of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

Application Within Residential Buildings Including Homes and Apartment Buildings and Within Nonfood Areas of Industrial, Institutional, and Commercial Buildings Including Hospitals, Stores, Manufacturing Plants and Warehouses:
For Spot Treatment Only—Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets; cracks and crevices in walls, along and behind baseboards; beneath and behind sinks, stoves, refrigerators and cabinets, around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For clover mites use a 0.5% spray only. **Note:** A period of 4 to 7 days is normally required for maximum effect on cockroaches.

For control of carpet beetles, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

For control of brown dog ticks, thoroughly apply a 0.5% spray as a spot treatment to infested areas such as pet beds and resting quarters, nearby cracks and crevices along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. **DO NOT TREAT PETS WITH THIS PRODUCT.**

Attention (Pests Indoors): Do not apply water-base sprays of this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is need to treat such areas use only oil-base sprays. Prior to application de-energize electrical equipment. After application allow time for oil to evaporate before starting motors or placing switches in the "ON" position. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

Pest	Dosage DURSBAN 2E per 1000 sq. ft.	Remarks
Ants	1-1 1/2 oz	
Armyworms	1-1 1/2 oz	
Chinch bugs	1-1 1/2 oz	
Clover mites	1-1 1/2 oz	
Crickets	1-1 1/2 oz	
Earwigs	1-1 1/2 oz	
Flaxs	1-1 1/2 oz	
Grasshoppers	1-1 1/2 oz	
Sod webworms	1-1 1/2 oz	
Ticks	1-1 1/2 oz	
White grubs (including European chaffer, southern masked chaffer and Japanese beetle larvae)	3 to 6 fl oz	For white grubs spray when grubs are young and actively feeding near the soil surface usually during late July and August or as recommended by your local Agricultural Extension Service specialist. If you are unable to determine the feeding area with 1/2 inch of water, wash the insecticide in the underlying soil.
Yucca root rot (Hydrocotyle weevil)	3 fl oz	Spray suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service specialist in your area.

Pest	DOSAGE of DURSABAN 2E per 1000 sq ft.	Remarks
Bluegrass grubbing	3 to 6 fl. oz.	Spray early in the season when adult grubs appear. Retreat when needed.
	7 1/2 fl. oz. per 1000 sq ft.	1 pint per 1000 sq ft.

Peach Tree Borer Apply DURSABAN 2E to flowering trees and shrubs of the Genus *Prunus* in water as a trunk spray before newly hatched larvae enter the trees. Apply as a coarse low pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat your area.

NOTE: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

Attention (Lawn and Ornamental Plant Pests): Keep children and pets off treated areas until spray has dried. Keep cut of lawn and garden fishpots and other bodies of water.

NOTE: Elm Bark Beets. Prevent an outbreak of beetles from overwintering in uninfested trees and to reduce beetle populations in disease free areas, apply DURSABAN 2E insecticide as a 0.5% mixture of chlorpyrifos in water to the bottom 9 feet of trunk. Wet the trunk thoroughly but do not spray to runoff. Care should be taken to apply the spray right to the base of the trunk. Applications can be made with either a backpack blower or a hydraulic pressure sprayer from spring through to the early fall. Prevention of Branch and Twig Feeding on High Value Elm Trees. To reduce the twig and branch feeding on uninfested trees deemed to be high in value apply a 0.5% chlorpyrifos water spray to the tree crown. Application

should be made in the spring or early summer using a sprayer that will give thorough coverage to the tree crown.

Area Control of Ticks and Chiggers. Use DURSABAN 2E insecticide to control ticks and chiggers infesting non-cropland areas such as roadsides, footpaths and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply DURSABAN 2E in water at the rate of 1 pint per acre (equivalent to 1 fl. oz. per 2,700 sq ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer, or other suitable hand or power spray equipment. Treat low underbrush, grassy areas, weeds and ground surface and debris using enough spray volume to obtain thorough coverage usually 40 to 100 gallons per acre.

Attention (Area Control of Ticks and Chiggers) Do not allow public use of treated areas during application or until spray has dried.

Imported Fire Ant. Treatment of potted, containerized and balled and burlapped nursery stock. Use DURSABAN 2E to control imported fire ants in soil attached to roots of these plants by completely submerging the ball of soil in a tank containing DURSABAN 2E diluted at the rate of 8 fl. oz. per 100 gallons of water. Do not remove burlap wrap or plastic containers with drain holes prior to submerging. Keep soil submerged until complete saturation has occurred, normally about 30 seconds. Wear rubber gloves, boots and apron during this operation. An alternate treatment to submerging potted plants is to apply a solution containing

DURSABAN 2E (8 oz per 100 gallons of water) as a substitute for plain water during routine watering activities. This solution should be applied to the potting soil on a twice daily schedule for three consecutive days. Do not remove burlap wrap or container from plants prior to treatment. Application should be made in a well ventilated area.

Treatment of Fire Ant Mounds. Apply DURSABAN 2E as a granular dust at the rate of 1 fl. oz. (30 cc) per 2 gallons of water. Gently sprinkle a total of 1 gallon of the diluted solution over the surface of each mound using a sprinkler can. Thoroughly wet mound and surrounding area to a 4 ft diameter (2 sq ft) for best results apply in cool weather, 65-80 F. or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration reducing product effectiveness.

Pests Controlled	2 Gallons	10 Gallons	100 Gallons
Appl. 25	0.5 fl. oz.	1.6 fl. oz.	16 fl. oz.
Banyan			
Ed. Tree (Lent)			
Cherry (Lent)			
Fall Weebill			
Grasshoppers			
Yellow-necked caterpillar			
Mimosa webworm	1.0 fl. oz.	3.2 fl. oz.	32 fl. oz.
Redhumped caterpillar			
Thrips (exposed)			

Pests Controlled	3 Gallons	10 Gallons	100 Gallons
Ash borer	7.0 fl. oz.	6.4 fl. oz.	64 fl. oz.
Dogwood borer			
Lilac borer			
Rhododendron borer			
White peach borer			

Mosquitoes: Apply DURSABAN 2E insecticide to mosquito infested non-crop areas according to the conditions and rates shown in the following table. Mix DURSABAN 2E with either water or oil such as kerosene or No. 2 fuel oil in sufficient spray volume to give uniform coverage of areas treated. Apply the spray using suitable application equipment such as hand and power ground sprayers, mist applicators and aerial spray equipment.

Warning (Mosquito Control) Read complete Environmental and Use Precautions and consult your State Fish and Game Department before using this product for mosquito control.

Waste Water (High in Organic Content) Use DURSABAN 2E insecticide for control of mosquito larvae in agricultural, industrial, and urban waste waters that contain high levels of organic matter including sewage sludge, ponds and aquaria. Industrial waste water impoundments, cesspools, and septic tanks catch basins, etc. are the most susceptible to mosquito larvae infestation.

DOSAGE for MOSQUITOES in WASTE WATER

Area to be Treated	Dosage of DURSABAN 2E	Use Directions
Sewage drains, ditches, ponds and lagoons	2 fl. oz. per 5000 cu ft. water	Apply DURSABAN 2E to the surface of the water. Do not apply to the bottom of the pond or lagoon.
Industrial waste water impoundments	2 fl. oz. per 100 cu ft. water	Apply DURSABAN 2E to the surface of the water. Do not apply to the bottom of the pond or lagoon.
Catch basins	0.08 fl. oz. per cu ft. water	Apply DURSABAN 2E to the surface of the water. Do not apply to the bottom of the pond or lagoon.

Condition of Area to be Treated	Amount to Use per Acre		Number Acres 1 Gallon of DURSABAN 2E Will Treat
	DURSABAN 2E (fluid ounces)	Chlorpyrifos (pounds)	
Adult Control			
Light to medium vegetative cover	1.5	0.035	32
Medium to heavy vegetative cover	3.2	0.05	4

The higher dosage will also give faster knockdown and longer residual control. Always use this little or no vegetative cover.

Concentration of 0.4 ounces of chlorpyrifos per 100 gallons is equivalent to 8 ounces of chlorpyrifos per 100 gallons. Equivalent to 3 pounds of chlorpyrifos per 100 gallons.

U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) <small>WASHINGTON, D.C. 20460</small>	EPA REGISTRATION NO.	DATE OF ISSUANCE
	TERM OF ISSUANCE	
	NAME OF PESTICIDE PRODUCT	

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
---------------------------------	------