

10F15
pm 12
464-590

APR 3 1985

Ms. Judith Page
Dow Chemical USA
P.O. Box 1706
Midland, MI 48674

Dear Ms. Page:

Subject: Use on Pecans
Dursban 50W
EPA Registration No. 464-590
Your Application Dated March 13, 1985

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A) is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

JOB:85071:RD-85/04:Ellenberger:3/26/85:Del.4/2/85:Kendrick:898-1270:EK

CONCURRENCES

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

20F15

ACCEPTED
with COMMENTS

DRAFT SPECIMEN LABEL

APR 8 1985

464-590

Dursban* 50W

Insecticide

To be Applied Only by or Under the Supervision of Commercial Applicators Responsible for Insect Control

Programs: Not Intended for Sale to or for Use by Homeowners. Do Not Formulate This Product into Other End-Use Products.

Active Ingredient(s):

Chlorpyrifos [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 50%

Inert Ingredients 50%

E.P.A. Registration No. 464-590

E.P.A. Est. 5905-AR-1

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO: PRECAUCION AL USUARIO:

Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Hazards to Humans and Domestic Animals

MAY BE FATAL IF SWALLOWED • HARMFUL IF ABSORBED THROUGH SKIN • CAUSES EYE IRRITATION

PRECAUTIONARY STATEMENT: DO NOT TAKE INTERNALLY •

Avoid Contact With Skin, Eyes, Or Clothing • Avoid Breathing Dust Or Spray Mist • Wash Thoroughly With Soap And Water After Handling And Before Eating Or Smoking • Do Not Get in Eyes, On Skin, Or On Clothing • Remove And Wash Contaminated Clothing Before Reuse

Statements of Practical Treatment if Swallowed:

Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. If On Skin: Wash with plenty of soap and water. Get medical attention if irritation persists. If In Eyes: Flush with plenty of water. Get medical attention if irritation persists. If Inhaled: Remove to fresh air if effects occur. Get medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate degree of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/ protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidotes immediately after establishing an open airway and respiration.

Environmental Hazards

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area. DURSBAN 50W insecticide is highly toxic to fish, birds and other wildlife. Fish and crustaceans may be killed if their waters are contaminated with this product. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Use this product only as specified on this label.

Read the entire label. Use only according to label directions.

Before buying or using this product, read "WARRANTY LIMITATIONS AND DISCLAIMER" below. If terms are not acceptable, return unopened package at once to seller for full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under the Limit of Warranty and Liability.

In case of an emergency endangering life or property involving this product, call collect 517-636-4400

AGRICULTURAL CHEMICAL

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

DURSBAN 50W insecticide is a wettable powder formulation designed for use as a spray to control various pests injurious to trees, turf and ornamental plants. The pests controlled are listed in the accompanying tables. DURSBAN 50W is compatible with insecticides, miticides, and fungicides commonly recommended except for alkaline materials such as Bordeaux mixture and lime. A small amount of spray mixture should be prepared to check for compatibility before a large volume of spray is mixed.

Applicator Safe Handling Procedures: When exposure to spray mist is likely to occur, wear protective equipment and clothing (chemical goggles, long-sleeved coveralls and waterproof gloves) to prevent exposure of eyes and skin. To prevent breathing of spray mist during application in confined areas, wear a respirator and cartridge(s) approved by NIOSH/MSHA for pesticides.

Attention: Do not allow adults, children or pets on treated areas until spray has dried. Keep out of fishpools and other bodies of water. Do not treat vegetable gardens. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings (hay) or seed screenings to livestock nor use hay for livestock bedding. Do not use in poultry houses.

MIXING INSTRUCTIONS

DURSBAN 50W mixes readily with water to form a suspension. Mix the required amount of DURSBAN 50W with a small quantity of water and add this premix while filling the spray tank. Alternatively, fill the tank with the required amount of water and then add the premix. Maintain sufficient agitation during both mixing and application to ensure uniform suspension of the DURSBAN 50W in the spray mixture. DURSBAN 50W contains a wetting agent. Do not use additional wetting agents, spreaders or stickers.

Turf and Other Outdoor Uses

Use DURSBAN 50W to control pests infesting lawns, golf courses, parks, perennial turf grasses, etc., listed in the following table by application at the recommended dosages and in accordance with the directions given below. Dilute DURSBAN 50W in enough water to obtain complete and uniform coverage of pest-infested areas and apply as a coarse, low-pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf, except as noted. For best results, the lawn should be moist at time of treatment. Spray when pests first appear; retreat when needed.

Pest*	Amount of DURSBAN 50W per		Specific Directions
	1000 sq. ft.	Acre	
Ants (including foraging fire ants) Armyworms (Such as: Beet, Fall, Yellowstriped) Centipedes Chiggers(1) Chinch bugs Clover mites Cutworms Crickets Farwigs Fly skipper Fleas Gnats Grasshoppers Greenbugs June beetles Leafhoppers Lucerne moth Millipedes Mites (Such as: Clover, Formula grass, Winter grain) Mosquitoes(2) Sod webworm (lawn moths)(3) Sowbugs Springtails Ticks(1)	3/4 oz.	2 lb.	1. Use DURSBAN 50W insecticide for area control of Ticks and Chiggers infesting non-cropland areas such as roadsides, footpaths, and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or a possible public health problem. Do not allow public use of treated areas during application or until spray has dried. Apply DURSBAN 50W in water at the rate of 1/2 pound per acre (equivalent to 1/2 oz. per 2,700 sq. ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer, or other suitable hand- or power-operated spray equipment. Treat low underbrush, grassy areas, weeds, ground surfaces and debris using enough spray volume to obtain thorough coverage, usually 40 to 100 gallons per acre. 2. Mosquitoes coming to rest on areas treated for control of turf pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions. 3. For sod webworms delay watering or mowing the treated area for 12 to 24 hours after treatment.
European crane fly	1 oz	3 lb	
Turfgrass weevil (<i>Hyperodes</i>)(1)	1 1/2 oz.	4 lb.	1. Spray suspected problem areas in mid-April and again in mid-May or as recommended by your Local Agricultural Extension Service Specialist.

*Numbers in parentheses refer to Specific Directions.

Pest*	Amount of DURSBAN 50W per		Specific Directions
	1000 sq. ft.	acre	
White grubs (Such as: Black turfgrass ataenius, European chafer, Southern masked chafer and Japanese beetle larvae)(1)	1 1/2 to 3 oz.	4 lb. to 8 lb.	1. For white grubs, spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.
Bluegrass billbugs(1)	3/4 to 3 oz.	2 lb. to 8 lb.	1. For bluegrass billbugs spray early in the season when adults first appear. Retreat as needed.
Chiggers (Turkey Pens)(1)	3 oz.	8 lb.	1. For control of chiggers infesting outdoor turkey pens apply DURSBAN 50W as a spray to the soil using 100-150 gallons of water per acre. To prevent turkeys from becoming infested with chiggers, spray the soil in pens just before turkeys are transferred to them. For turkeys already infested with chiggers, spray the soil in pens thoroughly. Allow at least three weeks for chigger lesions on the turkeys to heal before turkeys are slaughtered. For optimum results, spray a 10-15 ft. protective area around the periphery of the treated pens. Treatment is limited to 2 applications to the soil in turkey pens at intervals of not less than 4 weeks. Do not treat soil in turkey pens later than 7 days before turkeys are to be slaughtered. Avoid getting the spray on feed or in waterers. Open feed should be covered during spraying operations, and water troughs should be flushed out immediately after the spraying operations. Do not spray DURSBAN 50W directly on the turkeys.
Imported fire ants in commercial sod(1)	6 oz.	16 lbs.	1. Imported fire ants: Apply DURSBAN 50W as a spray to the area of sod to be cut. Immediately after applying the spray soak in with at least 1/2 inch of water or a sufficient volume to thoroughly soak below the cut line. Do not apply a volume or rate of water to cause runoff. Sod can be cut when vegetation has dried and when soil is moist but not wet. Precaution: Personnel handling treated sod should wear elbow-length neoprene gloves and neoprene apron. Protective clothing and boots should be worn to prevent repeated skin exposure to treated sod.

* Numbers in parentheses refer to Specific Directions.

Certification Period: 42 days from day of

Pest Control on Outside Surfaces of and Around Buildings

Apply DURSBAN 50W by application as a residual spray to outside surfaces of buildings including porches, window frames, eaves, patios, garages, refuse dumps and other areas where pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness. Avoid application to surfaces where visible residues are objectionable.

Pest*	Amount of DURSBAN 50W in Water to Make:			Specific Directions
	1 gal	10 gal	50 gal	
Ants Bees Carpenter ants Clover mites Cockroaches Crickets Earwigs Fleas Flies Hornets Millipedes Mosquitoes Scorpions(2) Silverfish Sowbugs Spiders Symphylans Ticks Wasps Yellowjackets	For Band Treatment (1)			1. To help prevent infestation of buildings, treat a band of soil 6 to 10 feet wide around and adjacent to buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 4 ounces of DURSBAN 50W per 50 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1,000 sq. ft. to thoroughly and uniformly wet the band area. 2. For scorpions treat or remove accumulations of lumber, firewood and other materials that serve as insect harborage sites.
	1/12 oz.	3/4 oz.	4 oz.	
	On Outside Surfaces			
	1 1/3 oz.	13 1/3 oz.	4 lb.	

*Numbers in parentheses refer to Specific Directions.

Ornamentals (Outdoor, Greenhouse, Shadehouse, Nurseries, etc.)

Use DURSBAN 50W insecticide to treat flowers, shrubs, evergreens, vines and shade and flowering trees found to be infested with pests listed in the following table. Dilute DURSBAN 50W with water according to directions given in the table and apply using suitable hand- or power-operated spray equipment in a manner to provide complete and uniform coverage. For best results apply a coarse spray to thoroughly wet both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed.

Consult your State Agricultural Experiment Station or Extension Service Specialist for application timing and other specific use information.

Note: Greenhouse Grown Ornamentals: Environmental factors significantly affect phytotoxicity. DURSBAN 50W has been tested on numerous ornamental plants without causing serious phytotoxicity. However, because of the numerous varieties grown, it is recommended that a small group of the plants be treated at the recommended rate under the anticipated growing conditions and observed for phytotoxic symptoms. Two test sprays should be made 7 days apart and the plants observed for another 7 days after the second treatment. The following species should not be treated:

Plant Species	
Scientific Name	Common Name
<i>Acalyphis wilkesiana</i>	Copperleaf
<i>Aphelandra squarosa</i>	Zebra plant
<i>Brassaia actinophylla</i>	Schefflera
<i>Bucida buceras</i>	Black olive
<i>Carica papaya</i>	Papayas
<i>Cissus antarctica</i>	Cissus
<i>Codiaeum variegatum</i>	Croton
<i>Ficus nitida</i>	Ficus
<i>Ficus pumila</i>	Weeping fig
<i>Ficus retusa</i>	Cuban laurel
<i>Hibiscus calycinus</i>	Yellow hibiscus
<i>Hibiscus rosa-sinensis</i>	Red/Chinese hibiscus
<i>Impatiens spp.</i>	Impatiens
<i>Nephrolepis exaltata</i>	Boston fern, Fluffy ruffle fern
<i>Petunia spp.</i>	Petunia
<i>Rosa spp.</i>	Belinda, Caramia, Lavande, Jack Frost

Pest*	Amount of Dursban 50W in to Make: Water			Specific Directions
	1 gal	3 gal	100 gal	
Adelgids (Such as: Cooley and Eastern spruce galls, Pine bark) Ants (including foraging fire ants) Aphids (Such as: Apple, Chrysanthemum, Cottonwood, Elm leaf, Peach, Rose, Spirea) Armyworms (Fall, Yellowstriped) Bagworms(1) Boxelder bugs Cankerworms Catalpa sphinx Cicadas Citrus mealybug Cockroaches (Such as: American, Brownbanded, German, Oriental, Smokey brown) Elm spanworm Fall webworm Grasshoppers Green fruitworm Hornworms Jackpine budworm Juniper webworm Katydid Lace bugs Leafhoppers Leafrollers(2) Maple leafcutter(3) Mites (Such as: Clover, Red spider, Southern red)(4) Nelder caterpillar Orange tortrix Periodical cicada Plant bugs Poplar tentmaker Psyllids Puss caterpillar Rose chafer Sawflies, exposed (Such as: Pine, Pin oak, Redheaded) Sowbugs Spittlebugs Spring elm caterpillar Springtails Spruce budworm Tent caterpillars (Such as: Eastern, Forest, Western) Thornbug Walnut caterpillar Western spruce budworm Whiteflies Yellownecked caterpillar	1/12 oz.	1/4 oz.	1/2 lb.	1. For bagworms treat when insects are in the crawler stage. 2. For effective control of leafrollers spray should be applied before leaves are tightly rolled. 3. For effective control of maple leafcutter on maple trees apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production. 4. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days in the South or 7 to 10 days in the North after initial treatment to control newly-hatched nymphs.

*Numbers in parentheses refer to Specific Directions.

Pest*	Amount of DURSBAN 50W in Water to Make:			Specific Directions
	1 gal	3 gal	100 gal	
Armyworm (Beet) Beetles (Fuller rose, Native elm bark)(1) Cutworms Leafhoppers Mahogany webworm Mealybugs Mimosa webworm Moths (Such as: Browntail, Cypress tip, Douglas fir tussock, European pine shoot, Gypsy(2), Holly bud, Nantucket pine tip, Pandora, Pitch pine tip, Subtropical pine tip, Tussock) kworms (Such as: California, Orangestriped, Redhumped) Redhumped caterpillar Thrips (exposed) Weevils (Such as: Blackvine(3), Yellow Poplar, Pine reproduction)	1/6 oz.	1/2 oz.	1 lb.	<ol style="list-style-type: none"> 1. To reduce twig and branch feeding by bark beetles applications should be made in the spring or early summer. 2. To kill migrating and invading gypsy moth larvae treat trunks and foliage. 3. Blackvine weevils are night feeders. Late afternoon spraying will give control in some areas.
Beetles (Such as: Cottonwood leaf(1), Elm leaf, Flea, Willow leaf)	1/6 to 1/3 oz.	1/2 to 1 oz.	1 to 2 lb.	<ol style="list-style-type: none"> 1. For cottonwood leaf beetles use DURSBAN 50W in water to control larvae and adults infesting cottonwoods. Make the treatment when field counts indicate damaging beetle populations are developing or present.
Borers(1): Clearwing moths (Such as: Ash, Dogwood, Lesser peachtree, Lilac, Oak, Rhododendron), Metallic wood (Such as: Bronze birch, Flatheaded appletree, Twolined chestnut), Longhorned beetles (Such as: Locust, Red oak) Leafminers Needleminers (Such as: Jeffrey pine, Lodgepole pine, Spruce) Scale crawlers (Such as: Cottonycushion, Cottony maple, Euonymus, Fletcher, Florida wax, Golden oak, Hemispherical, Lecanium, Magnolia, Oak kermes, Oystershell, Pine needle, San Jose, Tea, White birch)	1/3 oz.	1 oz.	2 lb.	<ol style="list-style-type: none"> 1. For borers, apply DURSBAN 50W to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat. Apply uniformly as a coarse low-pressure spray.
Borers (Such as: Cottonwood, Peachtree)(1)	1 oz.	3 oz.	6 lb.	<ol style="list-style-type: none"> 1. For peachtree borers apply DURSBAN 50W in water to flowering trees and shrubs of the genus <i>Prunus</i> as a trunk spray before newly-hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.

*Numbers in parentheses refer to Specific Directions.

Pest*	Amount of DURSBAN 50W in Water to Make:			Specific Directions
	1 gal	3 gal	100 gal	
Beetles (Such as: Ambrosia, Anobiidae, Black turpentine, Cottonwood leaf, Elm leaf, European elm bark, Flea, Fuller rose, Japanese, June, Native elm bark, Pine, Willow leaf)(1)	2 2/3 oz.	8 oz.	16 1/2 lb.	<p>1. For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees.</p> <p>For remedial treatment apply the spray to the main trunk of infested trees when damage occurs but before adult beetles begin to emerge.</p>

*Numbers in parentheses refer to Specific Directions.

Tree Pests (Nurseries, Plantations and Felled Trees)

Use DURSBAN 50W insecticide to treat shade and flowering trees, plantation trees, transplant trees and evergreens found to be infested with pests listed in the following table. Felled trees should be treated as necessary. Dilute DURSBAN 50W with water according to directions given in the table and apply using suitable hand- or power-operated spray equipment in a manner to provide complete and uniform coverage. For best results apply a coarse spray to thoroughly wet both upper and lower leaf surfaces and to infested limb and trunk areas.

Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. Attention: For felled trees, treat after cutting; do not handle until spray has dried or wear protective clothing. Treat only fruit and nut trees identified in the Tree Fruits and Nuts section of this label.

Consult your State Agricultural Experimental Station or Extension Service specialist for application timing and other specific use information.

Pest*	Amount of DURSBAN 50W in Water to Make:			Specific Directions
	1 gal	3 gal	100 gal	
Adelgids (Such as: Cooley and Eastern spruce gall, Pine bark) Aphids (Such as: Apple, Chrysanthemum, Cottonwood, Elm leaf, Peach, Rose, Spirea) Bagworms(1) Boxelder bugs Cankerworms Catalpa sphinx Citrus mealybugs Elm spanworm Fall webworm Fire ants Greenstriped mapleworm Jackpine budworm Juniper webworm Lygids Lace bugs Leafhoppers Leafrollers(2) Maple leafcutter(3) Mites (Such as: Clover, Red spider, Southern red)(4) Oak skeletonizer Poplar tentmaker Puss caterpillar Sawflies, exposed (Such as: Pine, Pin oak) Spring elm caterpillar Spruce budworm Tent caterpillars (Such as: Eastern, Forest, Western) Walnut caterpillar Western spruce budworm Yellownecked caterpillar	1/12 oz.	1/4 oz.	1/2 lb.	1. For bagworms treat when insects are in the crawler stage. 2. For effective control of leafrollers spray should be applied before leave are tightly rolled. 3. For maple leafcutter on maple trees apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production. 4. For effective control of spider mites when large numbers of eggs are present, apply a second spray after 3 to 5 days in the South or 7 to 10 days in the North after initial treatment to control newly-hatched nymphs.

*Numbers in parentheses refer to Specific Directions.

120F15

Pest*	Amount of DURSBAN 50W in Water to Make:			Specific Directions
	1 gal	3 gal	100 gal	
Beetles (Fuller rose, Native elm bark)(1) Leafhoppers Mahogany webworm Mealybugs Mimosa webworm Moths (Such as: Browntail, Cypress tip, Douglas fir tussock, European pine shoot, Gypsy(2), Holly bud, Nantucket pine tip, Pandora, Pitch pine tip, Subtropical pine tip, Tussock) Oakworms (Such as: California, Orangestriped, Redhumped) Redhumped caterpillar Thrips (exposed) Weevils (Such as: Blackvine(3), Yellow poplar, Pine reproduction)	1/6 oz.	1/2 oz.	1 lb.	1. To reduce the twig and branch feeding on uninfested trees deemed to be of high value, apply a water spray to the tree crown. Application should be made in the spring or early summer using a sprayer that will give thorough coverage to the tree crown. 2. To kill migrating and invading gypsy moth larvae treat trunks and foliage. 3. Blackvine weevils are night feeders. Late afternoon spraying will give control in some areas.
Beetles (Such as: Cottonwood leaf(1), Elm leaf, Flea, Willow leaf)	1/6 to 1/3 oz.	1/2 to 1 oz.	1 to 2 lb.	For cottonwood leaf beetles use DURSBAN 50W in water to control larvae and adults infesting cottonwoods. Make the treatment when field counts indicate damaging beetle populations or developing or present. For seedlings use 8 to 20 gallons per spray volume per acre.
Borers(1): Clearwing moths (Such as: Ash, Dogwood, Lesser peachtree, Lilac, Oak, Rhododendron), Metallic wood (Such as: Bronze birch, Flatheaded appletree, Twolined chestnut), Longhorned beetles (Such as: Locust, Red oak) Cranberry girdler larvae(2) Leafminers Needleminers (Such as: Jeffrey pine, Lodgepole pine, Spruce) Scale crawlers (Such as: Cottonycushion, Cottony maple, Euonymus, Fletcher, Florida wax, Golden oak, Hemispherical, Lecanium, Magnolia, Oak kermes, Oystershell, Pine needle, San Jose, Tea, White birch)	1/3 oz.	1 oz.	2 lb.	1. For borers apply DURSBAN 50W to the trunks and lower limbs of trees when the adults begin to emerge. Consult your local State Agricultural Experimental Station or Extension Service specialist for proper time to treat. Apply uniformly as a coarse low-pressure spray. 2. For cranberry girdler larvae infesting Douglas fir seedlings apply 2 lb. of DURSBAN 50W per acre. Direct spray at the lower crown and stems using 50 gallons of water per acre. Irrigate immediately after application for soil penetration of 1-2 inches. Treat after egg laying during the summer.
Borers (Such as: Cottonwood, Peachtree)(1)	1 oz.	3 oz.	6 lb.	1. For peach tree borers apply DURSBAN 50W in water to flowering trees of the genus <i>Prunus</i> as a trunk spray before newly-hatched larvae enter the trees. Apply as a coarse low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.

*Numbers in parentheses refer to Specific Directions

Pest*	Amount of DURSBAN 50W in Water to Make:			Specific Directions
	1 gal	3 gal	100 gal	
Beetles(1) (Such as: Cottonwood leaf(2), Elm leaf, Flea, Fuller rose, Native elm bark(3), Willow leaf)	1 1/3 oz.	4 oz.	8 1/3 lb.	<ol style="list-style-type: none"> 1. For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment apply the spray to the main trunk of infested trees or logs when damage occurs but before adult beetles begin to emerge. For plantation trees apply to individual trees using suitable hand- or power-operated spray equipment. 2. To prevent native elm bark beetles from overwintering in uninfested trees, apply DURSBAN 50W in water to the bottom 9 feet of the trunk. Wet the trunk thoroughly but do not spray to runoff. Care should be taken to apply the spray right to the base of the root flare. Application can be made with either a backpack mistblower or a hydraulic pressure sprayer from spring through to early fall. To reduce the twig and branch feeding on uninfested trees deemed to be of high value, apply a water spray to the tree crown. Application should be made in the spring or early summer using a sprayer that will give thorough coverage to the tree crown.
Beetles (Such as: Ambrosia, Anobiidae, Black turpentine, European elm bark, Japanese, June, Pine, Spruce)(1) Carpenter ants Termites	2 2/3 oz.	8 oz.	16 2/3 lb.	<ol style="list-style-type: none"> 1. For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment apply the spray to the main trunk of infested trees or logs when damage occurs or before adult beetles begin to emerge. For plantation trees apply to individual trees using suitable hand- or power-operated spray equipment.
Weevils (Such as: Northern pine, Pales, Pitch-eating, Twig)(1)	5 1/3 oz.	1 lb.	33 1/3 lb.	<ol style="list-style-type: none"> 1. For Southern pine seedlings treat immediately after transplanting. Treat each seedling with enough spray to thoroughly wet the foliage and stems to the point of runoff but do not use more than 6 gallons of spray per acre.

*Numbers in parentheses refer to Specific Directions.

Tree Fruits and Nuts

Use DURSBAN 50W insecticide at the dosages indicated by application as a foliar spray to control pests listed in the following table. Mix the required dosage of DURSBAN 50W in water. Spray for thorough and complete coverage of the foliage, using suitable hand- or conventional power-operated spray equipment. Treat when pests first appear or in accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service specialist for specific use information. Do not allow livestock to graze in treated areas. **Attention:** In California do not work with or contact trees within 2 days after treatment.

146F15

Crop	Pest	Amount of DURSBAN 50W in Water to Make:			Specific Directions
		1 gallon	3 gallons	100 gallons	
Apples	Apple aphid Apple maggot Codling moth European apple sawfly Eyespotted bud moth Fruittree leafroller Green fruitworm Lesser appleworm Obliquebanded leafroller Orange tortrix Oriental fruit moth Pandemis leafroller Plum curculio Redbanded leafroller Rosy apple aphid San Jose scale Tarnished plant bug Tufted apple budmoth Variegated leafroller Woolly apple aphid	1/2 oz.	1 1/2 oz.	3 lb.	Make no more than 8 applications per season. Do not apply last treatment within 28 days before harvest or apply last two treatments closer than 21 days apart.
Almonds	Navel orangeworm Peach twig borer San Jose scale	2/3 oz.	2 oz.	4 lb.	Make no more than 3 applications per season to almonds and filberts and no more than 2 applications per season to walnuts. Do not apply within 14 days before harvest.
Filberts	Eyespotted bud moth Filbert aphid Filbert leafroller Filbertworm Obliquebanded leafroller Omniverous leaftier Winter moth	1/2 oz. to 2/3 oz.	1 1/2 oz. to 2 oz.	3 to 4 lb.	
Walnuts	Codling moth Walnut scale	2/3 oz.	2 oz.	4 lb.	
Pecans	Black pecan aphid Hickory shuckworm Fecan leaf scorch mite Pecan nut casebearer Pecan weevil <i>Phylloxera spp.</i>	1/3 oz.	1 oz.	2 lb.	Make no more than 5 applications per season. Do not apply within 28 days of harvest. Do not allow livestock to graze in treated orchards.

156F15

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 100° F. Do not stack higher than 2 pallets. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

Pesticide disposal: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container disposal: Completely empty fiber can by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of containers in a sanitary landfill or by incineration if allowed by State and local authorities.

WARRANTY LIMITATIONS AND DISCLAIMER

The Dow Chemical Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions therein under normal conditions of use. **THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL.** Therefore, neither this warranty nor any other warranty or merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.) under abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify in writing The Dow Chemical Company of any claims to be eligible to receive either remedy given below. **The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of The Dow Chemical Company or any other seller will be one of the following, at the election of The Dow Chemical Company:**

- (1) Refund of purchase price paid by buyer or user for product bought.
- (2) Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses. The terms of this Limit of Warranty and Liability cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations and Disclaimer in any manner.

0.91 kg/2 lb net

* Trademark of THE DOW CHEMICAL COMPANY

26075-L4

B584

DOW DIAMOND*

THE DOW CHEMICAL COMPANY
Midland, Michigan 48640 U.S.A.

LOT