

**NOTICE OF PESTICIDE:** REGISTRATION  
 REREGRISTRATION*(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)*

NAME OF PESTICIDE PRODUCT

Dursban 50 DF

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Dow Chemical U.S.A.  
P.O. Box 1706  
Midland, MI 48640-1706

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/dite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase "EPA Registration No. 464-643" to your label before you release the product for shipment.
3. This will acknowledge receipt of your revised and corrected Confidential Statement of Formula dated November 20, 1988. The information is acceptable and it will be placed in our file for your product.
4. Submit five (5) copies of your final label before you release the product for shipment. Refer to the label for a further description of final printing details.

If these conditions are not met, the registration of this product is subject to cancellation in accordance with FIFRA section 3(c)(7)(B) and for shipment of the product constitutes a violation of the Act.

A stamped copy of the label is enclosed for your records.

**BEST AVAILABLE COPY**Dennis H. Edwards, Jr.  
Product Manager (111)  
Institutional Products Division  
Registration Division (375356)

Enclosures

 ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

*Dennis H. Edwards Jr.*

DATE

12/5/89

BE FATAL IF INHALED. DO NOT BREATHE DUST. WEAR A MASK OR PESTICIDE RESPIRATOR JOINTLY APPROVED BY THE MINING ENFORCEMENT AND SAFETY ADMINISTRATION AND THE NATIONAL INSTITUTE FOR OCCUPATIONAL SAFETY AND HEALTH.

# Specimen Label

# Dursban\* 50 DF Insecticide

ACCEPTED  
with COMMENTS  
in EPA Letter Date:

DEC 5 1971

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registration under EPA Reg. No. 1474-1045

To Be Applied Only By or Under the Direct Supervision Of  
Commercial Applicators Responsible For Insect Control Programs.

**Active Ingredient(s):**

Chlorpyrifos (O,O-diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate)

**Inert Ingredient:**

E.P.A. Registration No. 464-

E.P.A. Est. 1471-IN-5 464-MI-1

**KEEP OUT OF REACH OF CHILDREN**

**BEST AVAILABLE COPY**

50%  
50%

Do not contaminate water when disposing of equipment washwaters.

## WARNING

### AVISO:

#### PRECAUCION AL USUARIO:

Si usted no lee estas instrucciones, no use este producto hasta que se asegure de haber sido educado apropiadamente.

#### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

**MAY BE FATAL IF SWALLOWED • HARMFUL IF ABSORBED THROUGH SKIN • CAUSES EYE IRRITATION**

Do Not Get In Eyes, On Skin Or Clothing • Avoid Breathing Spray Mist • Wash Thoroughly With Soap And Water After Handling And Before Eating Or Smoking • Remove Contaminated Clothing And Wash Before Reuse • Keep Away From Food, Feedstuffs And Water Supplies

**STATEMENT OF PRACTICAL TREATMENT:** If Swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. If On Skin: Wash with plenty of soap and water. Get medical attention. If In Eyes: Flush with plenty of water for five minutes. Get medical attention if irritation persists. If Irritated: Remove to fresh air if symptoms of characteristic irritation appear and get medical attention immediately.

**NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, ataxia and red blood cell cholinesterase tests may increase significance of exposure (baseline tests are useful). Atropine, only by injection, is the anticholinergic antidote. Oximes, such as 2-PAM (pralidoxime), may be ineffective if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use atropine immediately after establishing an open airway and respiration.

#### Environmental Hazards

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water by cleaning of equipment or disposal of waste. • This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

#### NOTICE

Read and understand the entire label before using. This only according to label directions. Before buying or using this product, read "WARRANTY LIMITATIONS AND DISCLAIMER" statement on container. If terms are not acceptable, return unopened package at once to seller for full refund of purchase price and. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under WARRANTY LIMITATIONS AND DISCLAIMER.

#### IN CASE OF AN EMERGENCY

endangering life or property involving this product, call collect 817-636-4488  
**DIRECTIONS FOR USE**  
Do not apply product through any type of irrigation system

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
Do not formulate this product into other end-use products.

Dursban\* 50 DF

Do Not Mix or Store with Food, Feeds, Dye or Clothing

**BEST AVAILABLE COPY**

but not more often than once per every 7 days.

**GENERAL INFORMATION**

DURSBAN 50DF insecticide is a wettable granule formulation designed for use as a spray to control various pests injurious to trees, turf and ornamental plants. The pests controlled are listed in the accompanying tables. DURSBAN 50DF is compatible with insecticides, miticides, herbicides and fungicides commonly recommended except for alkaline materials such as Bordeaux mixture and lime. A small amount of spray mixture should be prepared to check for compatibility before a large volume of spray is mixed.

**Applicator Safe Handling Procedures:** When exposure to spray mist is likely to occur, wear protective equipment and clothing (chemical goggles, long-sleeved coveralls and waterproof gloves) to prevent exposure of eyes and skin. To prevent breathing of spray mist during application in confined areas, wear a respirator and cartridge(s) approved by NIOSH/MSHA for pesticides.

**Attention:** Do not allow adults, children or pets on treated areas until spray has dried. Keep out of fishpools and other bodies of water. Do not treat vegetable gardens. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings (hay) or seed screenings to livestock nor use hay for livestock bedding. Do not use in poultry houses.

**MIXING INSTRUCTIONS**

DURSBAN 50DF granules should be mixed with water to form a suspension. Fill the tank with half of the required amount of water, add the recommended amount of DURSBAN 50DF, then add the balance of the water. Maintain sufficient agitation during mixing and application to ensure uniform suspension of DURSBAN 50DF in the spray mixture. **NOTE: Do not use additional wetting agents, surfactants or stickers. If these are used, they may cause compatibility problems. Always use water to check and determine if there is any incompatibility before application.**

**Turf and Other Outdoor Uses**

Use DURSBAN 50DF to control pests infesting lawns, golf courses, parks, perennial turf grasses, etc., listed in the following table by applying at the recommended dosages and in accordance with the directions given below. Dilute DURSBAN 50DF in enough water to obtain complete and uniform coverage of pest-infested areas and apply as a coarse, low-pressure spray using suitable application equipment.

For best results, the lawn should be moist at time of treatment. Spray when pests first appear; retreat when needed.

4. For individual Fire ant mounds, apply DURSBAN 50DF as a drench at a rate of 1.5 oz. per each 2 gallons of water. Gently sprinkle 1 to 2 gallons of dilution over the surface of each mound. Avoid disturbing mounds and treat new mounds as they appear.

Pest	Amount of DURSBAN 50DF per		Specific Directions
	1000 sq. ft.	Acre	
<p>Ants (including foraging fire ants)</p> <p>Armyworms (Such as: Beet, Fall, Yellow-clover)</p> <p>Caterpillars</p> <p>Chiggers(1)</p> <p>Chinch bugs</p> <p>Clover moths</p> <p>Culex moths</p> <p>Crickets</p> <p>Earrings</p> <p>Fiery skipper</p> <p>Fleas</p> <p>Gnats</p> <p>Grasshoppers</p> <p>Greenbugs</p> <p>June beetles</p> <p>Leafhoppers</p> <p>Lusarna moth</p> <p>Millipedes</p> <p>Mites (Such as: Clover, Winter grain)</p> <p>Mosquitoes(2)</p> <p>Sod webworm (lawn moths)(3)</p> <p>Sowbugs</p> <p>Springtails</p> <p>Ticks(1)</p>	<p>3/4 oz.</p>	<p>2 lb.</p>	<p>1. Use DURSBAN 50DF insecticide for area control of Ticks and Chiggers infesting non-cropland areas such as roadsides, footpaths, and trails, picnic and camping areas, parks and other recreational areas where these pests are present and create a nuisance or a possible public health problem. Do not allow public use of treated areas during application or until spray has dried. Apply DURSBAN 50DF in water at the rate of 1/2 pound per acre (equivalent to 1/2 oz. per 2,700 sq. ft.) using a hydraulic sprayer mist applicator, knapsack sprayer, or other suitable hand- or power-operated spray equipment. Treat low-lying, underbrush, grassy areas, weeds, ground surfaces and other areas using enough spray volume to obtain thorough coverage, usually 40 to 100 gallons per acre.</p> <p>2. Mosquitoes coming to rest on areas treated for control of turf pests will be killed for varying periods of time after treatment, depending on size of treated area to weathering conditions.</p> <p>3. For Sod webworm, delay watering and mowing the treated area for 12 to 24 hours after treatment.</p>
<p>European crane fly</p>	<p>1 oz.</p>	<p>3 lbs.</p>	
<p>Turfgrass weevil (Hyperodes)(1)</p>	<p>1 1/2 oz.</p>	<p>4 lb.</p>	<p>1. Spray suspected problem areas in mid-April and again in mid-May or as recommended by your Local Agricultural Extension Service specialist.</p>

Fire ants (mounds)(4)

Pillbugs

Bermuda grass stunt

<sup>1</sup>Numbers in parentheses refer to Specific Directions.

best results

Pest	Amount of DURBAN 50DF per		Specific Directions
	1000 sq. ft.	Acre	
White grubs (Such as: Black turgrass sternus. European crane. Southern masked crane and Japanese beetle larvae)(1)	1 1/2 to 3 oz.	4 lb. to 8 lb.	1. For white grubs, spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.
Bluegrass billbugs(1)	3/4 to 3 oz.	2 lb. to 8 lb.	1. For bluegrass billbugs spray early in the season when adults first appear. Retreat as needed.
Chiggers (Turkey Pens)(1)	3 oz.	8 lb.	1. For control of chiggers infesting outdoor turkey pens apply DURBAN 50DF as a spray to the soil using 100-150 gallons of water per acre.  To prevent turkeys from becoming infested with chiggers, spray the soil in pens just before turkeys are transferred to them. For turkeys already infested with chiggers, spray the soil in pens thoroughly. Allow at least three weeks for chigger lesions on the turkeys to heal before turkeys are slaughtered. For optimum results, spray a 10-15 ft. protective area around the periphery of the treated pens.  Treatment is limited to 2 applications to the soil in turkey pens at intervals of not less than 4 weeks. Do not treat soil in turkey pens later than 7 days before turkeys are to be slaughtered. Avoid getting the spray on feed or in waterers. Open feed should be covered during spraying operations, and water troughs should be flushed out immediately after the spraying operations. Do not spray DURBAN 50DF directly on the turkeys.
Imported fire ants in commercial sod(1)	6 oz.	16 lb.	1. Imported fire ants: Apply DURBAN 50DF as a spray to the area of sod to be cut. Immediately after applying the spray soak in with at least 1/2 acre- inch of water or a sufficient volume to thoroughly soak below the cut line. Do not apply a volume or rate of water to cause runoff. Sod can be cut when vegetation has dried and when soil is moist but not wet.  Precautions: Personnel handling treated sod should wear clothing, including gloves and respirator apron. Protective clothing and boots should be worn to prevent repeated skin exposure to treated sod.  Certification Period: 42 days from day of application.

(Numbers in parentheses refer to Specific Directions.

BEST AVAILABLE COPY

## Pest Control on Outside Surfaces of and Around Buildings

Apply DURSBAN 50DF by application as a residual spray to outside surfaces of buildings including porches, window frames, eaves, eaves, porches, garages, refuse dumps and other areas where pests congregates or have been seen. Repeat treatment as needed to maintain effectiveness. Avoid application to surfaces where visible residues are objectionable.

Pest	Amount of DURSBAN 50DF in Water to Make	Specific Directions	
			1 gallon
1/12 - 1/3 oz.			
3/4 - 3 oz.			
4 - 16 oz.			
	For Perimeter Treatment (1)		
	1/12 oz.	3/4 oz.	4 oz.
	On Outside Surfaces		
	1 1/3 oz.	13 1/3 oz.	4 lb.

1 To help prevent infestation of buildings, treat a band of soil 6 to 10 feet wide around and adjacent to buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 4 ounces of DURSBAN 50DF per 50 gallons of water and spray as a coarse spray at the rate of about 10 gallons spray mixture per 1 000 sq. ft. to thoroughly and uniformly wet the band area.

2 ←

- Boxeider bugs
- Ants
- Bees
- Carpenter ants
- Clover mites
- Cockroaches
- Crickets
- Earwigs
- Firebrats
- Flies
- Flies
- Hornets
- Moths
- Mosquitoes
- Scorpions (2)
- Silverfish
- Sowbugs
- Spiders
- Springtails
- Symonians
- Ticks
- Wasps
- Yellowjackets

\*Numbers in parentheses refer to Specific Directions.

**BEST AVAILABLE COPY**

### Ornamentals (Outdoor, Greenhouse, Shadehouse, Nurseries, etc.)

Use DURSBAN 50DF insecticide to treat flowers, shrubs, evergreens, vines and shade and flowering trees found to be infested with pests listed in the following table. Dilute DURSBAN 50DF with water according to directions given in the table and spray using suitable hand- or power-operated spray equipment in a manner to provide complete and uniform coverage. For best results apply a coarse spray to thoroughly wet both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed.

Consult your State Agricultural Experiment Station or Extension Service specialist for application timing and other specific use information.

#### While

Note: Greenhouse Grown Ornamentals: Environmental factors significantly affect phytotoxicity. DURSBAN 50DF has been tested on numerous ornamental plants without causing serious phytotoxicity, because of the numerous varieties grown, it is recommended that a small group of plants be treated at the recommended rate under the anticipated growing conditions and observed for phytotoxic symptoms. Two test sprays should be made 7 days apart and the plants observed for another 7 days after the second treatment.

### PLANT SPECIES DEMONSTRATING PHYTOTOXICITY TO DURSBAN 50DF

Scientific Name	Common Name
<i>Acrostichum whitehousei</i>	Copernice
<i>Achillea squarosa</i>	Zebra plant
<i>Breasa acrostichylo</i>	Schottiera
<i>Buaya buaya</i>	Black olive
<i>Carica papaya</i>	Papaya
<i>Cissus antarctica</i>	Cissus
<i>Codiaeum variegatum</i>	Croton
<i>Ficus nitida</i>	Ficus
<i>Ficus purpurea</i>	Weeping fig
<i>Ficus retusa</i>	Cuban laurel
<i>Hibiscus caryensis</i>	Yellow hibiscus
<i>Hibiscus rosa-sinensis</i>	Red Chinese hibiscus
<i>Impatiens</i> spp.	Impatiens
<i>Neohortensia exaltata</i>	Boston Fern, Fluffy ruffles fern
<i>Petunia</i> spp.	Petunia
<i>Rosa</i> spp.	Begonia, Caranba, Lavender, Jack Frost

For Scorpions remove accumulations of lumber, firewood, and other materials which serve as harborage sites. DURSBAN 50DF may be applied as a residual spray to surfaces immediately below such materials. Perimeter treatments may reduce pest immigration from surrounding areas.

Do not use additional wetting agents, surfactants or stickers.

Significant phytotoxicity has been observed in the following species following treatment with DURSBAN 50DF. Apply to these species at your own risk.

Pest	Amount of GURBAN 50F in Water to Make			Specific Directions
	1 gallon	3 gallons	100 gallons	
Adelgids (Such as: Cooley and Eastern spruce galls, Pine bark) Ants (including foraging fire ants) Aphids (Such as: Apple, Chrysanthemum, Cottonwood, Elm leaf, Peach, Rose, Sorel) Armyworms (Fall, Yellowstriped) Bagworm(1) Borerworms Caterpillars Caterpillars Caterpillars Chiggers Citrus mealybug Coccinellids (Such as: American, Asian, Wood, Smoky brown) Elm sawworms Fall webworms Grasshoppers Green fruitworms Hornworms Jackpine budworms Juniper webworms Katydid Lace bugs Leafhoppers Leafrollers(2) Maple leafcutters(3) Mites (Such as: Red spider, Southern red(4)) Oleander caterpillars Orange sawfly Periodical cicada Plant bugs Pear sawfly Psyllids Pear caterpillars Rose chafers Sawflies, exochus (Such as: Pine, Pin oak, Redheaded) Sawflies Spittlebugs Spring elm caterpillars Spruce budworms Tent caterpillars (Such as: Eastern, Forest, Western) Thrips Walnut caterpillars Western spruce budworms Whiteflies Yellowheaded caterpillars	1/12 oz.	1/4 oz.	1/2 lb.	1. For best results treat when insects are in the crawler stage. 2. For effective control of leafrollers spray should be applied before leaves are tightly rolled. 3. For effective control of maple leafroller on maple trees spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production. 4. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days in the South or 7 to 10 days in the North after initial treatment to control newly-hatched nymphs. 5. For effective control of Fall webworm direct spray into web and immediately surrounding foliage.
Armyworms (Belt) Beetles (Native elm bark(1)) Cutworms Leafhoppers Mahogany webworms Mealybugs Mine and sawworms Moths (Such as: Brownish, Cypress, G. Douglas fir tussock, European pine shoot, Gypsy(2), Holly bud, Nantucket pine tip, Pandora, Pitch pine tip, Subarctic pine tip, Tussock) Oakworms (Such as: California, Orangeheaded, Redheaded) Redheaded caterpillars Thistle sawflies Worms (Such as: Blackvine(3), Yellow Pecan, Pine reproduction)	1/8 oz.	1/2 oz.	1 lb.	1. To reduce twig and branch breakage, best beetle applications should be made in the spring by early summer. 2. To kill migrating and growing caterpillars, with larvae treat trunk and foliage. 3. Blackvine weevil is night feeder. Late afternoon spraying will give control in some areas.

**BEST AVAILABLE COPY**

\*Numbers in parentheses refer to Specific Directions.








Pest	Amount of DURBAN 50F in Water to Make			Specific Directions
	1 gallon	3 gallons	100 gallons	
Borers (Such as: Cottonwood. Peachtree(1))	1 oz.	3 oz.	6 lb.	1. For peach tree borers apply DURBAN 50F in water to flowering trees of the genus Prunus as a trunk spray before newly-hatched larvae enter the trees. Apply as a coarse low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.
Beetles(1) (Such as: Cottonwood leaf. Elm leaf. Flea. Fuller rose. Native elm bark(2). Willow leaf)	1 1/3 oz.	4 oz.	6 1/3 lb.	1. For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment apply the spray to the main trunk of infested trees or logs when damage occurs but before adult beetles begin to emerge. For plantation trees apply to individual trees using suitable hand- or power-operated spray equipment. 2. To prevent native elm bark beetles from overwintering in uninfested trees, apply DURBAN 50F in water to the bottom 9 feet of the trunk. Wet the trunk thoroughly but do not spray to runoff. Care should be taken to apply the spray right to the base of the root flare. Application can be made with either a backpack mistblower or a hydraulic pressure sprayer from spring through to early fall. To reduce the twig and branch feeding on uninfested trees deemed to be of high value, apply a weak spray to the tree crown. Application should be made in the spring or early summer using a sprayer that will give through coverage to the tree crown.
Beetles (Such as: Ambrosia. Anobias. Black turpentine. European elm bark. Japanese. Juno. Southern pine. Spruce(1))	2 2/3 oz.	8 oz.	16 2/3 lb.	1. For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment apply the spray to the main trunk of infested trees or logs when damage occurs or before adult beetles begin to emerge. For plantation trees apply to individual trees using suitable hand- or power-operated spray equipment.
Worms (Such as: Northern pine. Palea. Pine-needle. Twig)(1)	5 1/3 oz.	1 lb.	33 1/3 lb.	1. For Southern pine seedlings treat immediately after transplanting. Treat each seedling with enough spray to thoroughly wet the foliage and stems to the point of runoff but do not use more than 6 gallons of spray per acre.

(1)Numbers in parentheses refer to Specific Directions.



**BEST AVAILABLE COPY**

## Tree Fruits and Nuts

Use DURSIBAN 50DF insecticide at the dosages indicated by application as a leaf spray to control pests listed in the following table. Mix the required dosage of DURSIBAN 50DF in water. Spray for thorough and complete coverage of the foliage, using suitable hand- or conventional power-operated spray equipment. Treat when pests first appear or in

accordance with local conditions. Consult your State Agricultural Experiment Station or Extension Service specialist for specific use information. Do not allow livestock to graze in treated areas. Attention: in California do not work with or contact trees within 2 days after treatment.

Do not use additional wetting agents, surfactants or adjuvants.

Pest	Amount of DURSIBAN 50DF in Water to Make	Specific Directions		
		1 gallon	3 gallons	100 gallons
Apples Apple aphid Apple maggot Codling moth European apple sawfly Eyedotted bud moth Fruitree leafroller Green fruitworm Lesser appleworm Obliquebanded leafroller Orange tortrix Oriental fruit moth Pangedms leafroller Plum curculio Redbanded leafroller Rosy apple aphid San Jose scale Tarnished plant bug Tulleed apple budmoth Variegated leafroller Woolly apple aphid	1/2 oz.	1 1/2 oz.	3 lb.	Make no more than 8 applications per season. Do not apply last treatment within 28 days before harvest or apply last two treatments closer than 21 days apart.
Almonds Navel orangeworm Peach twig borer San Jose scale	2/3 oz.	2 oz.	4 lb.	Make no more than 3 applications per season to almonds and no more than 2 applications per season to walnuts. Do not apply within 14 days before harvest.
Walnuts Codling moth Walnut scale	2/3 oz.	2 oz.	4 lb.	
Pecans Black pecan aphid Hickory shuckworm Pecan leaf scorch mite Pecan nut casebearer Pecan weevil Phytophthora spp.	1/3 oz.	1 oz.	2 lb.	Make no more than 5 applications per season. Do not apply within 28 days of harvest. Do not allow livestock to graze in treated orchards.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 100°F. If container is damaged or seal occurs, use product immediately or dispose of product and damaged container as indicated below.

**Pesticide disposal:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

### Container disposal:

Do not reuse empty container. Completely empty container into application equipment. Runcture and/or crush empty container and dispose of in a sanitary landfill, or by incineration, or, if allowed by local authorities, by burning.

## WARRANTY LIMITATIONS AND DISCLAIMER

The Dow Chemical Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose stated on the label when used in strict accordance with the directions thereon under normal conditions of use. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT, NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither the Company nor any other warranty of merchantability or fitness for a particular purpose, express or implied, applies to the use of this product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.) or under abnormal conditions (such as excessive rainfall, drought, torpedoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.



**BEST AVAILABLE COPY**

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify in writing The Dow Chemical Company of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of The Dow Chemical Company or any other seller will be one of the following, at the election of The Dow Chemical Company:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations And Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations And Disclaimer in any manner.

26075-L3

**ORM-A**

**RQ/CHLORPYRIFOS. MIXTURE  
NA2783**

**BEST AVAILABLE COPY**



**THE DOW CHEMICAL COMPANY**  
Midland, Michigan 48674 U.S.A.

\* Trademark of THE DOW CHEMICAL COMPANY