

464-700

4/1/2010

1/12

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

APR 1 2010



Office of Pesticide Programs

The Dow Chemical Company
1803 Building
Midland, Mi 48674

Attention: Rhonda Vance-Moeser
Senior Regulatory Specialist

Subject: GLUTEX™ GQ1 Sanitizer and Disinfectant
EPA Registration No. 464-700
Amendment Dated December 10, 2009

The amendment, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable.

Proposed Amendment


1. Change the primary product name
2. Alternate brand name
3. Add a disinfectant claim
4. Delete the egg sanitization claim
5. Revise label language in response to the RED

General Comment

The request to change the alternate brand name from UCARCIDE™ 14 Antimicrobial to "GLUTEX™ GQ1 Sanitizer and Disinfectant" is acceptable.

A stamped copy of the "accepted" label and Efficacy Review is enclosed for your records.

If you have any questions concerning this letter, please contact Martha Terry at (703) 308-6217.

Sincerely

Marshall Swindell
Product Manager (33)
Regulatory Management Branch 1
Antimicrobials Division (7510P)

Enclosure

GLUTEX™ GQ1 Sanitizer and Disinfectant

Active Ingredient: Glutaraldehyde.....	14.0%
Alkyl (C ₁₄ 50%, C ₁₂ 40%, C ₁₆ 10%).....	2.5%
dimethyl benzyl ammonium chloride	
Inert Ingredients:	83.5%
Total	100%

EPA Reg. No. 464-700

EPA Est. No. 10352-WV-2^(A); 464-WV-1^(B) 10352-WV-1^(C)

(A) 5 gallon pails in pallets; all drums, bulk

(B) Sample, 5 gallon pails less than full pallet

(C) Bulk Shipments

ACCEPTED
with COMMENTS
EPA Letter Dated:

APR - 1 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No. 464-700

KEEP OUT OF REACH OF CHILDREN

DANGER

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive • Causes irreversible eye damage • Causes skin irritation • Harmful if inhaled.
Harmful if swallowed • Harmful if absorbed through skin • Prolonged or frequently repeated
skin contact may cause allergic reactions in some individuals • Causes asthmatic signs and
symptoms in hyper-reactive individuals • Do not get in eyes, on skin, on clothing • Avoid
breathing vapor • Do not swallow • Wash thoroughly with soap and water after handling •
Remove contaminated clothing and wash before reuse.

Personal Protective Equipment

Applicators and other handlers must wear: goggles or face shield, long-sleeve shirt, long pants,
shoes, socks, chemical-resistant gloves (such as butyl rubber) and chemical resistant apron.

User Safety Requirements

Follow manufacturers' instructions for cleaning & maintaining PPE if no such instructions exist
for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users must wash hands before eating, drinking, chewing gum, or using the toilet.

GLUTEX™ GQ1 Sanitizer and Disinfectant
 FINAL MASTER LABEL
 December, 2009

ACCEPTED
 with COMMENTS
 EPA Letter Dated:

APR - 1 2010

Users must remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users must remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 464.700

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

FIRST AID	
IF SWALLOWED	<ul style="list-style-type: none"> • If the person is fully alert and cooperative, have the person rinse mouth with plenty of water. have the patient drink 4 to 10 ounces (120-300 mL) of water or milk. • Do not induce vomiting. Do not attempt mouth rinse if the person has respiratory distress, altered mental status, or nausea and vomiting. • Call a physician and/or transport to emergency facility immediately. See Note to Physician.
IF IN EYES	<ul style="list-style-type: none"> • Wash immediately and continuously with flowing water for at least 30 minutes. • Remove contact lenses after the first 5 minutes and continue washing. Obtain prompt medical consultation, preferably from an ophthalmologist. • Call a poison control center or a doctor immediately for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, and then give artificial respiration preferably mouth to mouth if possible. • Call a poison control center or a doctor for further treatment advice
Have product container or label with you when calling a poison control center or doctor or going for treatment.	
HOT LINE NUMBER	
IN CASE OF AN EMERGENCY endangering life or property involving this product, call collect (989) 636-4400.	
NOTE TO PHYSICIAN: Aspiration may cause lung damage. Probable mucosal damage may contraindicate the use of gastric lavage. Have the product container or label with you when calling a poison control center or a doctor, or going for treatment.	

APR - 1 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
commonly used materials
EPA Reg. No. 464-700

STORAGE AND HANDLING

GLUTEX GQ1 Sanitizer and Disinfectant is incompatible with many materials of construction such as steel, galvanized iron, aluminum, tin, and zinc. Sanitizer and Disinfectant can be stored and handled in baked phenolic-lined steel, polyethylene, stainless steel, or reinforced epoxy-plastic equipment. This product freezes at about 27°F (-3°C). Therefore, unless the storage tank is inside or underground, heating and insulation may be required. If heating is needed, exposure to high temperatures should be avoided. For short storage times (up to about 1 month), temperatures of up to 100°F (37.8°C) can be tolerated but the preferred maximum storage temperature is about 80°F (26.7°C).

A stainless steel centrifugal pump is suggested for transfer service. Spiral-wound stainless steel with TEFLON® Polymer is suitable for gaskets and packing.

Handle in a well-ventilated area. If vapors are irritating to the nose or eyes, special ventilation or respiratory protection (MSHA/NIOSH approved air purifying respirator equipped with an organic vapor cartridge) may be required.

The product in its undiluted form must not be used in a spray, fogging or aerosol application.

STORAGE AND DISPOSAL

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or your Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Option to use for labels on nonrefillable rigid containers of all sizes

Nonrefillable container. Do not reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Option to use for labels on refillable rigid tote containers

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning of the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean container before final disposal, empty contents into application equipment and triple rinse. Pour or pump rinsate into application equipment or rinsate collection system. Then offer for recycling if available, or puncture and dispose of in sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

AIR WASHERS AND INDUSTRIAL SCRUBBING SYSTEMS/ RECIRCULATING COOLING AND PROCESS WATER SYSTEMS

This product may be used only in industrial air washers and air washer systems which have mist-eliminating components.

GLUTEX GQ1 Sanitizer and Disinfectant should be added at the application rates described below to a water treatment system at a convenient point of uniform mixing such as the basin area. Addition may be made intermittently (SLUG DOSE) or continuously. Badly fouled systems can be shock treated with GLUTEX GQ1 Sanitizer and Disinfectant. Under these conditions, blowdown should be discontinued for up to 24 hours.

GLUTEX GQ1 Sanitizer and Disinfectant can be used in industrial process water systems that contain ultra filtration units and non-medical reverse osmosis membranes (where approved for compatibility by the membrane manufacturer) and associated distribution systems.

Intermittent (Slug Dose) Method

Initial Dose: When the system is noticeably fouled, apply 4.2 to 8.5 fluid ounces of GLUTEX GQ1 Sanitizer and Disinfectant per 100 gallons of water in the system. Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 1.7 to 4.2 fluid ounces of GLUTEX GQ1 Sanitizer and Disinfectant per 100 gallons of water in the system weekly or as needed to maintain control.

Badly fouled systems must be cleaned before treatment is begun.

Continuous Feed Method

Initial Dose: When the system is noticeably fouled, apply 4.2 to 8.5 fluid ounces of GLUTEX GQ1 Sanitizer and Disinfectant per 100 gallons of water in the system.

Subsequent Dose: Maintain this treatment level by starting a continuous feed of 0.8 to 4.2 fluid ounces of GLUTEX GQ1 Sanitizer and Disinfectant per 100 gallons of water in the system per day.

Badly fouled systems must be cleaned before treatment is begun.

SERVICE WATER AND AUXILIARY SYSTEMS

GLUTEX GQ1 Sanitizer and Disinfectant should be used at the same application rates, and in the same manner as described above. It should be added to the system at a point that will allow for uniform mixing throughout the system.

ACCEPTED
with COMMENTS
EPA Letter Dated:

APR - 7 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 414-700

HEAT TRANSFER SYSTEMS

(Evaporative Condensers, Dairy Sweetwater Systems, Hydrostatic Sterilizers and Retorts, and Pasteurizers and Warmers)

GLUTEX GQ1 Sanitizer and Disinfectant should be used at the same application rates, and in the same manner as described above. It should be added to the system at a point of uniform mixing such as a basin area, sump area or other reservoir or collecting area from which the treated water will be circulated uniformly throughout the system.

INDUSTRIAL WASTEWATER SYSTEMS

(Wastewater Systems, Wastewater Sludge and Wastewater Holding Tanks)

GLUTEX GQ1 Sanitizer and Disinfectant should be added to a wastewater system or sludge at a convenient point of uniform mixing such as the digester. Add 200 to 975 fluid ounces (1.6 to 7.6 gallons) (1,600 to 7,600 ppm product) of GLUTEX GQ1 Sanitizer and Disinfectant per 1,000 gallons of wastewater or sludge.

PAPER MILLS AND PAPER MILL PROCESS WATER SYSTEMS

GLUTEX GQ1 Sanitizer and Disinfectant should be added to the paper making system at a point of uniform mixing such as the beaters, broke chest pump, save-all tank, or white-water tank.

Initial Dose: When the system is noticeably contaminated, add 1.5 to 9.0 lbs of GLUTEX GQ1 Sanitizer and Disinfectant per ton or 0.76 to 4.5 Kg of GLUTEX GQ1 Sanitizer and Disinfectant per metric ton of pulp or paper (dry basis) as a slug dose. Repeat until control is achieved. Heavily fouled systems should be boiled out prior to initial treatment.

Subsequent Dose: When microbial control is evident, add 0.9 to 6.0 lbs of GLUTEX GQ1 Sanitizer and Disinfectant per ton or 0.46 to 3.0 Kg of GLUTEX GQ1 Sanitizer and Disinfectant per metric ton of pulp or paper (dry basis) as a slug dose as necessary to maintain control.

WATER BASED COATINGS FOR PAPER AND PAPERBOARD

NOTE: For use in non-food contact coatings only.

Use from 0.3 to 1.8 lbs. of GLUTEX GQ1 Sanitizer and Disinfectant per 1,000 lbs. of dry powder or 0.3 to 1.8 Kg of GLUTEX GQ1 Sanitizer and Disinfectant per metric ton of dry slurry to produce a concentration from 300 to 1,800 ppm as product (based on slurry solids) in the mixed slurry.

ACCEPTED
with COMMENTS
EPA Letter Dated:

APR -- 1 2010

Under the Federal Insecticide
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 464-700

GLUTEX™ GQ1 Sanitizer and Disinfectant
FINAL MASTER LABEL
December, 2009

ACCEPTED
with COMMENTS
EPA Letter Dated:

APR - 1 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended for the application of
registered under EPA Reg. No. 464-70

WATER FLOODS

GLUTEX GQ1 Sanitizer and Disinfectant should be added to a water flood system at a point of uniform mixing.

Initial Treatment: When the system is noticeably contaminated, add 300 to 15,000 ppm GLUTEX GQ1 Sanitizer and Disinfectant to the system (0.3 to 14.5 gallons GLUTEX GQ1 Sanitizer and Disinfectant per 1,000 gallons flood water). Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 60 to 15,000 ppm GLUTEX GQ1 Sanitizer and Disinfectant (0.06 to 14.5 gallons GLUTEX GQ1 Sanitizer and Disinfectant per 1,000 gallons flood water) to the system weekly, or as needed to maintain control.

FRAC FLUIDS

Not registered for this use in the State of California

GLUTEX GQ1 Sanitizer and Disinfectant reduces bacterial contamination and degradation of fracturing fluids and gels used in oil and gas well stimulations. Add GLUTEX GQ1 Sanitizer and Disinfectant to the frac water storage tanks or directly into the well head injection pipeline as the water is being pumped down-hole.

Dose Range: GLUTEX GQ1 Sanitizer and Disinfectant should be added at a rate of 360 to 17,900 ppm (3.5 - 172.5 gallons per 10,000 gallons) depending on the degree of bacterial fouling in the source water.

DRILLING, COMPLETION, AND WORKOVER FLUIDS

GLUTEX GQ1 Sanitizer and Disinfectant should be added to a drilling fluid system at a point of uniform mixing such as the circulating mud tank.

Initial Treatment: Add 150 to 3,000 ppm GLUTEX GQ1 Sanitizer and Disinfectant (0.6 to 12.2 gallons GLUTEX GQ1 Sanitizer and Disinfectant per 100 barrels of fluid) to a freshly prepared fluid depending on the severity of contamination.

Maintenance Dosage: Maintain a concentration of 150 to 3,000 ppm GLUTEX GQ1 Sanitizer and Disinfectant by adding 0.6 to 12.2 gallons of GLUTEX GQ1 Sanitizer and Disinfectant per 100 barrels of additional fluid, or as needed, depending on the severity of contamination.

PACKER FLUIDS

GLUTEX GQ1 Sanitizer and Disinfectant should be added to a packer fluid at a point of uniform mixing such as a circulating holding tank. Add 150 to 1,800 ppm GLUTEX GQ1 Sanitizer and Disinfectant (0.6 to 7.3 gallons GLUTEX GQ1 Sanitizer and Disinfectant per 100 barrels of fluid) to a freshly prepared fluid depending on the severity of contamination. Seal the treated packer fluid in the wall between the casing and production tube.

OIL AND GAS PRODUCTION AND TRANSMISSION PIPELINE SYSTEMS

GLUTEX GQ1 Sanitizer and Disinfectant should be added to an oil/gas production or transmission line via direct injection. The application should be conducted to ensure maximum distribution of GLUTEX GQ1 Sanitizer and Disinfectant throughout the entire internal pipeline surface by adding a sufficient amount of biocide to detect/measure a residual concentration at the back end of the pipeline system. Criteria for success of the treatment will be a reduction in

GLUTEX™ GQ1 Sanitizer and Disinfectant
FINAL MASTER LABEL
December, 2009

bacterial counts and/or reduced corrosion rates. To facilitate application, it may be desirable to dilute the GLUTEX GQ1 Sanitizer and Disinfectant with an appropriate solvent immediately before use. The concentration in the solvent should not fall below an active concentration range of 500 to 5000 ppm based on the volume of water in the pipeline. Injections to the system should be weekly, or as needed to maintain control.

GAS STORAGE WELLS AND SYSTEMS

Individual injection wells should be treated with sufficient quantity of GLUTEX GQ1 Sanitizer and Disinfectant to produce a concentration of 1,500 to 15,000 ppm GLUTEX GQ1 Sanitizer and Disinfectant when diluted by the water present in the formation.

Injection should take place before gas is injected (during the summer). Injections should be repeated yearly, or as needed to maintain control.

Individual drips should be treated with a sufficient quantity of GLUTEX GQ1 Sanitizer and Disinfectant to produce a concentration of 600 to 6,000 ppm GLUTEX GQ1 Sanitizer and Disinfectant when diluted by the water present in the drip. Injections should be repeated yearly, or as needed to maintain control.

HYDROTESTING

Water used to hydrotest pipelines or vessels should contain 300 to 12,000 ppm GLUTEX GQ1 Sanitizer and Disinfectant (0.3 to 11.8 gallons GLUTEX GQ1 Sanitizer and Disinfectant per 1,000 gallons water), depending on water quality and length of time the equipment will remain idle.

PIPELINE PIGGING AND SCRAPING OPERATIONS

Add GLUTEX GQ1 Sanitizer and Disinfectant to a slug of water immediately following the scraper (ideally this water volume can be kept to a minimum and contained between the scraper and a trailing pig). Sufficient GLUTEX GQ1 Sanitizer and Disinfectant should be added to produce a concentration of 0.3 to 3% (0.29 to 2.9 gallon GLUTEX GQ1 Sanitizer and Disinfectant per 100 gallons water), depending on the length of the pipeline and the severity of biofouling.

ACCEPTED
with COMMENTS
EPA Letter Dated:

APR - 1 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 464-700

DISINFECTING OR SANITIZING NON-FOOD CONTACT SURFACES, FARM, ANIMAL, AND POULTRY HOUSING FACILITIES AND EQUIPMENT

**Not registered for this use in the State of California
DO NOT USE THIS PRODUCT ON FOOD OR FEED**

GLUTEX GQ1 Sanitizer and Disinfectant is suitable for use on a wide variety of surfaces including stainless steel, aluminum, painted concrete, plastic, and glazed tile.

Disinfection

For disinfection against the following pathogens: *Klebsiella pneumoniae, Escherichia coli, Pasturella multocida, Salmonella enterica, Campylobacter jejuni Clostridium perfringens Haemophilus paragallinarum, Mycoplasma hyopneumoniae, Mycoplasma gallisepticum, Staphylococcus aureus, Pseudomonas aeruginosa and Salmonella enterica and for the following viruses:* *Swine Influenza Virua (H1N1), Porcine parvovirus, Avian Infectious Laryngotracheitis Virus and Porcine Respiratory Reproductive Syndrome Virus* prepare disinfection solution according to the following:

Disinfection solution preparation:

1. Mix 1 fluid ounce of GLUTEX GQ1 Sanitizer and Disinfectant per gallon of water (8 mls per liter) to produce a solution containing 0.13% total active ingredients.
2. Allow prepared solution to stand on surface for at least 10 minutes, or until thoroughly dried.
3. Fresh solution should be prepared daily or more often if the solution becomes diluted or soiled

Sanitization

For sanitization against the following pathogens: *Klebsiella pneumoniae, Salmonella cholerasuis, Staphylococcus aureus, and Pseudomonas aeruginosa and for the following viruses:* *Infectious bursal disease, Avian influenza (H6N2 and H5N1), SARS-associated coronavirus, Porcine circovirus* and **fungus:** *Aspergillus fumigatus, prepare sanitizing solution according to the following:*

Sanitizing solution preparation:

1. Mix 0.45 to 1.8 fluid ounces of GLUTEX GQ1 Sanitizer and Disinfectant per gallon of water (3.5 to 14.1 milliliters per liter) to produce a solution containing 0.06 to 0.24% total active ingredients.
2. For application against **Porcine circovirus**, mix 0.9 to 2.2 fluid ounces of GLUTEX GQ1 Sanitizer and Disinfectant per gallon of water (6.9 to 17.3 milliliters per liter) to produce a solution containing 0.12 to 0.293% total active ingredients.
3. Allow to stand for at least five minutes, or until thoroughly dried.
4. Fresh solution should be prepared daily or more often if the solution becomes diluted or soiled.

ACCEPTED
with COMMENTS
EPA Letter Dated:

APR - 7 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 464-700

Farm Equipment and Animal Housing Buildings

(Poultry & turkey grow-out houses, laying houses, swine production and housing, barns and large animal buildings)

1. Remove all poultry and other animals and feed from the premises. Remove all portable equipment and empty all feeders and water troughs. If not removed, all treated feeders and waterers must be thoroughly rinsed with potable water prior to re-use.
2. Remove all litter and manure from floors and other surfaces.
3. Prepare either disinfection or sanitizing solution depending upon the organisms being treated by mixing the appropriate amount of GLUTEX GQ1 Sanitizer and Disinfectant with water. Refer to solution preparation above.
4. Using suitable commercially available equipment, saturate all surfaces with the prepared solution. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
5. Allow to stand for at least 5 minutes for sanitization or 10 minutes for disinfection.
6. Ventilate enclosed spaces.
7. Do not repopulate with poultry or other animals, or use equipment until treatment has been absorbed or dried.

Hatchers, Setters, and Chick Processing Facilities

Not for treatment of hatchers which contain chicks/eggs.

1. For disinfection or sanitizing of environmental surfaces prior to introduction of eggs: Remove all animals and feed from premises/vehicles and enclosures. Clean out feathers, fluff, or other debris.
2. Prepare either disinfection or sanitizing solution depending upon the organisms being treated by mixing the appropriate amount of GLUTEX GQ1 Sanitizer and Disinfectant with water. Refer to solution preparation above.
3. Thoroughly saturate all surfaces with the prepared solution. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
4. Allow to stand for at least 5 minutes for sanitization or 10 minutes for disinfection.
5. Ventilate thoroughly before reuse.
6. **Note:** for treatment of air and surfaces in high humidity hatchers and setters with water spray cooling: Fog or atomize this solution into setters or hatchers, using appropriate application equipment. Fogging of poultry houses should only be done in such a way that the operator is outside the poultry house when applying the fog. Wait at least 3 hours before re-entry after application
7. **Note:** For treatment of air and surfaces in low humidity hatchers with chilled coil cooling: Fog or atomize this product using appropriate application equipment. Fogging of poultry houses should only be done in such a way that the operator is outside the poultry house when applying the fog. Wait at least 3 hours before re-entry after application.

ACCEPTED
with COMMENTS
EPA Letter Dated:
APR - 1 2010

GLUTEX™ GQ1 Sanitizer and Disinfectant
FINAL MASTER LABEL
December, 2009

11
12

Trays, Racks, Carts, Chick Boxes, Cages, and other Hard Surfaces:

1. Remove all filth and heavy debris from surfaces by scraping or washing.
2. Prepare either disinfection or sanitizing solution depending upon the organisms being treated by mixing the appropriate amount of GLUTEX GQ1 Sanitizer and Disinfectant with water Refer to solution preparation above.
3. Apply by spraying or immersion, and allow to drain and air-dry. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
4. Allow to stand for at least 5 minutes for sanitization or 10 minutes or for longer for disinfection,
5. Ventilate enclosed spaces.
6. Do not repopulate with poultry or other animals, or use equipment until treatment has been absorbed or dried.

Trucks and other Vehicles

1. Remove all filth and heavy debris from surfaces by scraping or washing.
2. Prepare either disinfection or sanitizing solution depending upon the organisms being treated by mixing the appropriate amount of GLUTEX GQ1 Sanitizer and Disinfectant with water Refer to solution preparation above.
3. Apply by spraying or immersion, and allow to air-dry.
4. Allow to stand for at least 5 minutes for sanitization or 10 minutes or for longer for disinfection.
5. Ventilate enclosed spaces.
6. Do not repopulate with poultry or other animals, or use equipment until treatment has been absorbed or dried.

**DISINFECTING OR SANITIZING NON-FOOD CONTACT SURFACES,
INDUSTRIAL EQUIPMENT AND BUILDINGS**

Not registered for this use in the State of California

1. Remove all heavy filth and soil deposits from surfaces using a suitable cleaner.
2. Prepare either disinfection or sanitizing solution by mixing the appropriate amount of GLUTEX GQ1 Sanitizer and Disinfectant with water. Refer to solution preparation above.
3. Using suitable commercially available equipment, saturate all surfaces with the prepared solution. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
4. Allow to stand for at least 5 minutes for sanitization or 10 minutes or for longer for disinfection.
5. Ventilate enclosed spaces.
6. Do not repopulate with poultry or other animals, or use equipment until treatment has been absorbed or dried.

ACCEPTED
COMMENTS
EPA Letter Dated:
APR - 1 2010

GLUTEX™ GQ1 Sanitizer and Disinfectant
FINAL MASTER LABEL
December, 2009

12
12

NET Contents/Net Weight:

Container Size	Container Size	Container Size	Container Size
260.8 gallons	55 gallon	5 gallon	1 gallon
Net Weight 2,290 Pounds	Net Weight 470 Pounds	Net Weight 44 Pounds	Net Weight 8.6 Pounds

LOT No.: _____

Produced For (*used when product manufactured by contract manufacturer*)

Dow Diamond TM
THE DOW CHEMICAL COMPANY
Midland, Michigan 48674
989-636-4400

™ Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

ACCEPTED
with COMMENTS
EPA Letter Dated:

APR - 1 2010

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 464-700