


# GURNEY'S APPLE CEDAR RUST SPRAY



## CAUTION

**KEEP OUT OF THE REACH OF CHILDREN**

See back panel for additional cautions.

**ACTIVE INGREDIENTS:**

Zineb (zinc ethylene bisdithiocarbamate) ..... 75%  
(metallic zinc equivalent 17.7%)

**INERT INGREDIENTS** ..... 25%


Total ..... 100%

**NET WEIGHT ONE POUND**

Manufactured for

**Gurney Seed & Nursery Co.**

YANKTON, SOUTH DAKOTA 57078



Authorities rightly maintain that the secret of successful disease control depends largely upon the proper timing of the applications. Foliage growth is rapid in the early spring, so that there is a daily increase in leaf surface exposed to fungus. Keep this new growth covered by frequent and timely applications. Such a program is the best insurance against disease damage during the period of primary infection. Cover upper and under sides of the leaves thoroughly.

## DIRECTIONS

### FRUITS

Recommended concentrations are 2 lbs. per 100 gallons of water, or 1 level tablespoonful per gallon of water (unless otherwise specified).

**APPLES, PEARS:** Fire blight. Make two applications during bloom first at 10% bloom, second at full bloom stage. Under severe conditions continue through petal fall or first cover. **APPLES:** Scab, cedar apple rust, frog eye leaf spot, sooty blotch, fly speck, Brooks' spot. Begin at pre-pink stage and continue through cover sprays. Use 2 lbs. per 100 gallons in pre-pink to petal fall sprays - 1 lb. per 100 gallons beginning 1st and 2nd cover. 1/2 to 1 lb. per 100 gallons water may be combined with sulphur in pre-bloom stage.

**SOUR CHERRIES:** Leaf Spot and shot hole. Apply in regular leaf spot control schedule. Apply first spray when most of petals have fallen, next spray about 10 days after petal fall. Spray at 10 to 12 day intervals thereafter. Mix at the rate of 1 lb. per 100 gallons of water or 1 tablespoonful per gallon. Make no application within 7 days of harvest. After harvest, make one or two foliage applications for adequate leaf spot control.

**GRAPES:** Black rot, downy mildew, ripe rot. Begin just before leaf buds burst. Use at the rate of 1 1/2 lbs. per 100 gallons. Spray regularly at 10 to 14 day intervals until fruit begins to ripen. Make no application within 7 days of harvest.

### CAUTION

May cause irritation of nose, throat and skin! Do not breathe dust or spray mist. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Wash fruits and vegetables before eating. Keep out of reach of children.

**IMPORTANT:** Keep away from fire and sparks. When not in use, keep container tightly closed.

### NON-WARRANTY

The use of this material being beyond our knowledge and control, we make no warranty, express or implied, as to the effect of such use, whether or not in accordance with directions.

26601-MBEPCO

USDA Reg. No. 168