

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

CAUTION
HARMFUL IF SWALLOWED. WASH THOROUGHLY AFTER USING. MAY CAUSE SKIN IRRITATION. AVOID CONTACT WITH EYES, SKIN, AND CLOTHING. DO NOT STORE WITH OR NEAR FERTILIZERS, SEEDS, INSECTICIDES, OR FUNGICIDES. It is almost impossible to remove residues of this product from sprayers and spray equipment, particularly from non-metallic parts, e. g., wood, rubber, polyethylene, or fibre, and the equipment used to measure and apply this weed killer should not be used subsequently for applying other materials to plants or crops.

GURNEY'S LAWN WEED KILLER

Contains 2,4-D in Low-Volatile Esters and SILVEX* (2,4,5-TP)

For Common Chickweed, Mouse-Eared Chickweed, White Clover, Black Medic, and Certain Broadleaf Weeds in Lawns

ACTIVE INGREDIENTS:

Isooctyl Ester of (2,4-dichlorophenoxy-acetic acid) 9.06%
Isooctyl Ester of Silvex (2-(2,4,5-trichlorophenoxy) propionic acid) 4.26%

INERT INGREDIENTS: 86.68%

2,4-Dichlorophenoxyacetic acid equivalent 6.01% by weight or 0.50 lb. per gal.
2-(2,4,5-trichlorophenoxy) propionic acid equivalent 3.11% by weight or 0.27 lb. per gal.

CAUTION
KEEP OUT OF REACH OF CHILDREN. See side panels for additional cautions.

CONTAINS 1 1/2 LB. SILVEX PER GALLON

Net Contents 8 Oz.

Manufactured for
Gurney Seed & Nursery Co.
YANKTON, SOUTH DAKOTA 57078

CAUTION
Considerable caution must be exercised in using this product to avoid injury to crops and desirable plants. Do not apply to vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other crops or desirable plants and do not permit spray mist to drift onto them since even minute quantities may cause severe injury during both growing and dormant periods. Apply only when there is no hazard from drift. Coarse sprays are less likely to drift. High temperatures during or shortly after application of the product may cause vaporization which would injure susceptible plants growing nearby. Do not apply excessive quantities. Excessive amounts in the soil may temporarily inhibit seed germination and plant growth. Do not contaminate domestic or irrigating water supplies.

TO MIX

Place about one-half the required amount of water in the spray tank, add Lawn Weed Killer and stir or agitate thoroughly. Then add remainder of the water and agitate thoroughly. Provide agitation while spraying to prevent separation of the emulsion and to ensure uniform mixture in spray tank. If allowed to stand, reagitrate to uniform mixture before applying the spray.

DIRECTIONS FOR USE — LAWNS

To control common chickweed, mouse-eared chickweed, white clover, plantain, black medic, dandelion, buckhorn, ragweed, and lambsquarter, in established lawns and turf, use 1 1/2 tablespoonfuls (5 3/4 fluid ounces) in 3 gallons of water and apply mixture as a spray to wet uniformly the weed plants on 1,000 sq. ft. (20'x50') of area (equivalent to 2 gallons in 25 to 40 gallons of water per acre). **APPLY ONLY IN THE FALL OR EARLY SPRING**, since application during hot, dry weather may cause excessive damage to grass. Do not use on bentgrass, St. Augustine grass, Dichondra, Carpetgrass or clover lawns. Do not apply to Dichondra or Lippia lawns. Delay re-seeding treated areas for at least 3 weeks. To avoid damage to turf, do not apply in excess of the above stated quantity or rate. Do not overlap. Avoid double coverage when backing up and at turns. Do not spray onto the grass when nozzle is not moving, such as when cleaning or testing nozzles. Use a positive quick shut-off valve on spraying equipment. **APPLY ONLY TO ESTABLISHED GRASS—THOROUGHLY ROOTED AND TILLERED.** Do not apply to newly seeded areas or to seedling grass.

USDA Reg. No. 168-12 7-1964-A-MBLFCO

12-5-68
464-12
1708-17