

Label copy

FRONT PANEL

GURNEY'S

SEVIN\* GARDEN SPRAY

A 50% Wettable Powder for control of Japanese Beetles -  
17 Year Locust (Cicada), many other insects

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) ..... 50%  
INERT INGREDIENTS: ..... 50%

\*Trade Mark of Union Carbide Corp for the active ingredient,  
1-naphthyl N-methylcarbamate


U. S. Pat. No. 2,903,478; 3,009,855

For ORNAMENTALS - FLOWERS - LAWNS - FRUITS - VEGETABLES

(18 pt. type) CAUTION  
(12 pt. type) KEEP OUT OF REACH OF CHILDREN  
See side panel for additional caution.

Net Content 1 lb.

~~XXXXXXXXXX~~  
Packed for  
GURNEY SEED & NURSERY COMPANY  
Yankton, South Dakota



SIDE PANEL

DIRECTIONS


Mix 2 level tablespoonfuls Sevin Garden Spray in 1 gallon of water. To use, mix designated amount of Sevin Garden Spray in a little water to make a thin paste, then add required amount of water. Do not store ~~in~~ diluted spray. Make new dilution for each use. Keep spray mixture well agitated.

ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADY TREES: Japanese Beetles, Aphids, Bagworm, Elm Leaf Beetle, Certain Leafminers, Rose Slug, Periodical Cicada (17-year locust), Scale Insects, Tent Caterpillars, Blister Beetles, Leaf Rollers, Exposed Thrips - Start spraying when insects or their damage first appear. Repeat weekly or as necessary. Do not use on Boston Ivy.

FRUIT TREES, SUCH AS APPLES, PEARS AND PEACHES: Apple Maggot, Plum Curculio, Apple Aphid, Leaf Rollers, Lesser Peach Tree Borer, Oriental Fruit Moth, Codling Moth - Make one application between bud break and bloom, a second application at petal fall, and then at 3 to 4 week intervals for 3 to 4 months as necessary. To control rosy apple aphids apply before leaves are curled. May be applied within 1 day of harvest.

LAWNS - Fleas and Ants: Use 6 ounces in 15 gallons water and cover 850 sq. ft of established lawn. For best results, use lawn sprayer. Mow lawn first and apply after rain or watering. Do not water for at least two days after application. Repeat in 2 or 3 weeks or as necessary.

XXXXXXXXXX  
Packed for  
GURNEY SEED & NURSERY COMPANY  
Yankton, South Dakota



SIDE PANEL

DIRECTIONS

Mix 2 level tablespoonfuls Sevin Garden Spray in 1 gallon of water. To use, mix designated amount of Sevin Garden Spray in a little water to make a thin paste, then add required amount of water. Do not store ~~the~~ diluted spray. Make new dilution for each use. Keep spray mixture well agitated.

ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES: Japanese Beetles, Aphids, Bagworm, Elm Leaf Beetle, Certain Leafminers, Rose slug, Periodical Cicada (17-year locust), Scale Insects, Tent Caterpillars, Blister Beetles, Leaf Rollers, Exposed Thrips - Start spraying when insects or their damage first appear. Repeat weekly or as necessary. Do not use on Boston Ivy.

FRUIT TREES, SUCH AS APPLES, PEARS AND PEACHES: Apple Maggot, Plum Curculio, Apple Aphid, Leaf Rollers, Lesser Peach Tree Borer, Oriental Fruit Moth, Codling Moth - Make one application between bud break and bloom, a second application at petal fall, and then at 3 to 4 week intervals for 3 to 4 months as necessary. To control rosy apple aphids apply before leaves are curled. May be applied within 1 day of harvest.

LAWNS - Fleas and Ants: Use 6 ounces in 15 gallons water and cover 650 sq. ft of established lawn. For best results, use lawn sprayer. Mow lawn first and apply after rain or watering. Do not water for at least two days after application. Repeat in 2 or 3 weeks or as necessary.

GARDEN VEGETABLES

Use 2 tablespoons per gallon of water unless otherwise specified. Use sufficient spray to obtain full coverage. Keep spray mixture well agitated. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary.

POTATO, TOMATO, EGGPLANT and PEPPERS: Use 2 tablespoons per gallon of water for Colorado potato beetle, flea beetles and leafhoppers. Use 2 to 4 tablespoons per gallon for European ~~corn~~ borer, fall plant bugs and stink bugs.

CUCUMBERS, MELONS, PUMPKINS and SQUASH: Pickleworm and melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

CAUTION: Do not breathe dust. Do not take internally. Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid storage near feed and food products. Keep out of reach of children.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Label copy

side panel Gurney's Sevin\* Garden Spray - continued

---

CAUTION: For protection of honeybees avoid, if possible, use during periods when honeybees are not visiting crops. If used when honeybees are visiting crops beekeepers should be warned to locate hives at a safe distance until one week after application. To avoid possible fruit thinning on apples and foliar injury on McIntosh, do not apply within 15 days after petal fall. Dispose of container promptly when empty by burying with wastes.

NOTICE: SELLER makes no warranty, express or implied, concerning this material, except that it conforms to the chemical description on the label. Neither the Manufacturer nor the Seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefore and accepts and uses this material on these conditions.

SDA Reg. No. 468-

MBEPCo