

(LABEL COPY)

(front panel)

GURNEY'S

MALATHION 5% DUST
Insecticide

[468-23]
1 of 2

ACTIVE INGREDIENTS:

Malathion*

5%

INERT INGREDIENTS:

95%

*O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate

USDA Reg. No. 468-

(15 pt. type) CAUTION

(12 pt. type) KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed. Avoid prolonged breathing of dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds.

Do not reuse container. Destroy when empty.

Net Contents 2 lbs.

Manufactured ~~by~~ for
GURNEY SEED & NURSERY COMPANY
Yankton, South Dakota 57078

468 23

(BACK PANEL)

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

DIRECTIONS

(Vegetables, Small Fruit, Ornamentals)

Thorough, full-coverage applications should be made. Dust underneath and top of foliage thoroughly. Apply to both plants and insects. Begin application when plants are well established or when new growth appears. Repeat application as necessary.

CONTROL CHART

Apply before plants are infested and damaged by the following insects:

7-Day Interval Between Last Application And Harvest:

Cabbage, Kale, Brussels Sprouts, Mustard Greens (Aphids, Imported Cabbage Worm, Cabbage Looper); Carrots, Lettuce, Head (Aphids, Leafhoppers); Cauliflower (caterpillars); Radishes, Parsnips, Salsify, Horseradish, Swiss Chard, Spinach (Aphids); Peas (Pea Aphids) Apply to foliage

Do not reuse container. Destroy when empty.

ACCEPTED

Net Contents 2 lbs.

Manufactured by for
GURNEY SEED & NURSERY COMPANY
Yankton, South Dakota 57078

46623

(BACK PANEL)

NOTICE: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes all risk to persons or property arising out of use or handling and accepts the product on these conditions.

DIRECTIONS

(Vegetables, Small Fruit, Ornamentals)

Thorough, full-coverage applications should be made. Dust underneath and top of foliage thoroughly. Apply to both plants and insects. Begin application when plants are well established or when new growth appears. Repeat application as necessary.

CONTROL CHART

Apply before plants are infested and damaged by the following insects:

7-Day Interval Between Last Application And Harvest:

Cabbage, Kale, Brussels Sprouts, Mustard Greens (Aphids, Imported Cabbage Worm, Cabbage Looper); Carrots, Lettuce, Head (Aphids, Leafhoppers); Cauliflower (caterpillars); Radishes, Parsnips, Salsify, Horseradish, Swiss Chard, Spinach (Aphids); Peas (Pea Aphids) Apply to foliage during early bloom period before eggs are laid; repeat as necessary.

5-Day Interval Between Last Application And Harvest:

Corn, Sweet (Aphids);

3-Day Interval Between Last Application And Harvest;

Broccoli, Turnip (Aphids, Imported Cabbage Worm, Cabbage Looper); Onions (Thrips); Pumpkins (Squash Vine Borer) Apply once or twice a week to stems and vines at base of plants. Do not apply to pumpkins unless plants are dry; Strawberries (Lygus bug, Spittlebugs, Field Crickets, Thrips, Potato Leafhopper, Strawberry Leafhopper); Peppers (pepper maggots); Strawberry (Strawberry Root Weevil) Apply to soil surface before planting and work into top 6 to 8 inches of soil.

1-Day Interval Between Last Application And Harvest.

Asparagus (Thrips); Beans (Mexican Bean Beetle, Leafhopper, Cucumber Beetle); Cucumbers, Melons (Aphids, Leafhopper, Spider Mites, Cucumber Beetle) Do not apply to cucumber, melons unless plants are dry; Squash (Aphids, spider Mites, Squash Vine Borer, Cucumber Beetle) Apply weekly to stems and vines at base of plants. Do not apply to squash unless plants are dry; Tomato (Spider Mite, Aphids, Tomato Russet Mite, Drosophila); Blackberries, Boysenberries, (Aphids);

No time limitation:

Potatoes (Aphids, Leafhopper, Mealybug) for Mealybug, apply on foliage of infested plants and repeat as necessary; Sweet Potato (Leafhopper);

(LABEL COPY - CONTINUED) Gurney's Malathion 5% Dust - continued

(BACK PANEL - continued)

[468-23]
2 of 2

Ornamentals (Aphids, Spider Mites) Make thorough, full-coverage applications. Repeat applications as necessary. Injury may occur on certain ferns such as Boston, Mindenhair and on some species of crassula and canaerti juniper.

DIRECTIONS (Animals)

POULTRY: (Northern Fowl Mite, Poultry Lice) Dust individual birds by means of a shaker can or rotary hand duster. Repeat application in 4 to 8 weeks or when necessary.

HOGS: (Lice) Treat animals thoroughly. Repeat in 10 days and thereafter as needed.

HOGS and CATS: (Fleas) Direct Application: Apply sufficient dust to treat animal thoroughly. Use any garden-type duster, puff duster or shaker-type can. If animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animals should be closely examined for fleas between treatment. If the animal is confined to an area free of fleas, one treatment may be sufficient.

CATTLE (Beef and Non-Milking): (Lice) Treat animal thoroughly. Repeat in 10 days and thereafter as needed. (Horn Fly) Use 3 tablespoonfuls per animal. Apply on back and neck of the animal and repeat applications at 10 - 14 day intervals. Note: Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening.

DAIRY CATTLE: (Horn Fly) Use 3 tablespoonfuls per animal. Apply on back and neck of the animal. Do not apply to dairy cattle later than 3 hours before milking or during milking. Repeat applications at 10 - 14 day intervals.

Do not treat animals under one month of age. When applying dusts avoid contamination of feed, food containers and watering troughs.