the state in the second of ear diele and a surger a surger Charlene in the state

ROSE DUST is effective for ----

INSECT CONTROL: For control of Aphids, Red Spider Mites, Blister Beetles, Cucumber Beetle, Common Stalk Borer, Flea Beetles, Lace Bugs, Lygus Bugs, Rose Leaf Beetles, Thrips, Rose Budworms and Rose Leaf Rollers on: Roses, Carnations, Gladioli, Hollyhocks, Lilies, Dahlias, Snapdragons, Violets, Geraniums, Pansies, Zinnias, Chry-santhemums and Phlox.

DISEASE CONTROL: For control of Black Spot, Powdery Mildew, Rust, Leaf Spot.

DIRECTIONS FOR USE

Cover upper and under sides of the leaves thoroughly. Best results will be obtained by dusting in the early morning or evening when the air is calm and the temperature is low.

APPLICATION — Use a good dust gun and dust up and under the leaves.

Authorities rightly maintain that the secret of successful disease and pest control depends largely upon the proper timing of the applications. Foliage growth is rapid in the early spring, so that there is a daily increase in leaf surface exposed to fungus and insect attack. Keep this new growth covered by frequent and timeiy applications. Such a program is the best insurance against disease and insect damage during the period of primary infection and infestation. Apply at 7 to 10 day intervals or immediately after heavy rain.

CAUTION: Harmful if swallowed. Avoid inhalation of dust. Avoid prolonged or repeated contact with skin, wash thoroughly after using. Avoid contamination of feed and foodstuffs. Avoid storage near feed and food products. Keep out of reach of children.

This material should not be used in combination with or closely following bil sprays. Do not apply when temperature is over 85 degrees, ahrenheit or when foliage is wet.

Avoid use of this product on certain ferns including Boston Maidenhair and Pteris, as well as some species of Crassula.

NOTICE: Buyer assumes all r sks of use, storage or handling of this material not in strict accordance with directions given herewith.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Packed by weight not by volume. Product may settle in shipment.

USDA Reg. No. 471-8

Manufactured for HENRY FIELD SEED & NURSERY CO. SHENANDOAH, IOWA / Since 1892

Henry Field's





JAN 27 -----

SOR FOURIER AND A STATUS LER. |

ED UNDER N. 471-8

112.1.

TI DE

· CI

ACTIVE INGREDIENTS: *Methoxychlor, technical Malathion 0,0-dimethyl dithiophosphate of diethyl mereaptosuccinate) Folpet N-(frichloromethylthio) phthalimide** INERT INGREDIENTS:

3.00% 4.00% 7.50% 85.50%

100.00%

10 *Equivalent to 2.64% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 0.36% other isomers and related compounds. **Phaltan

CAUTION: Keep out of reach of children.

See back panel for additional cautions.

NET WEIGHT THREE POUNDS

