

Henry Field's

CRAB GRASS CONTROL

Contains DACTHAL

ACTIVE INGREDIENT:

Dimethyl ester of tetrachloroterephthalic acid*

4.16%

INERT INGREDIENTS

95.84%

*from DACTHAL(R), (R) Diamond Alkali Company, licensed under U.S. Pat. No. 2,923,634.

MAY 31, 1974

471-10

CAUTION: Keep out of reach of children.

See side panel for additional cautions.

NET WEIGHT FOURTEEN OUNCES



19 APR 1974
REGISTRATION DIVISION

DIRECTIONS FOR USE

This package will treat 140 square feet.

Apply on a calm day to avoid wind drift. Apply at the rate of 8 ounces per 80 sq. feet lawn area, 3 lbs. to 475 sq. feet lawn area, 5 lbs. to 800 sq. feet lawn area. In order to kill the crabgrass, or weeds listed below, it is essential to make application very evenly with no skips or misses. It is suggested that the area be covered twice, the second time at right angles to the first to insure thorough coverage.

Do not rake the lawn or disturb soil surface after application. Water in lightly if no rain falls within a week after application.

Crabgrass Control is a pre-emergence herbicide and therefore must be applied in early spring prior to the germination of crabgrass or the weeds listed below: (This usually coincides with the time when Forsythia blossoms start to fall.)

Carpetweed, Common Chickweed, Greenfoxtail, Hairy Crabgrass, Johnson Grass (from seed), Lambsquarters, Purslane, Smooth Crabgrass, Texas Millet, Witchgrass, Yellow Foxtail.

This product does not provide control if applied after such weeds appear.

One application, applied before weed germination in early spring, usually provides up to 3 months control. Lawns will thicken naturally after spring removal of crabgrass and other competing weeds. At the recommended rate of application, Crabgrass Control does not show herbicide injury or discoloration to established common lawn grasses.

Where some spring seeding is necessary, there should be a delay of approximately 60 days after application. This delayed seeding improves the survival percentage of desirable grass seedlings. If reseeding is done between 30 and 60 days after application of Crabgrass Control, the amount of lawn seed sown should be increased by 50% to insure thrifty stands. In no case should reseeding be done earlier than 30 days after application.

Early spring application of Crabgrass Control herbicide may be applied to newly seeded lawns after they have exhibited greening of newly sprouted lawn grass. Such early seeding, followed by Crabgrass Control application, allows control of the crabgrass without injury to newly germinated turf grasses. Early spring application of Crabgrass Control allows fall sowing of permanent grasses.

CAUTION

Harmful if swallowed. Avoid contact with eyes. Wash after handling or using. Do not contaminate feed and foodstuffs.

Do not apply Crabgrass Control to newly seeded lawn areas until after greening appears. Do not use on Dichondra lawns. Do not store near seeds, fertilizers, or other pesticides.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

NOTICE: Seller makes no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning the use of this product or uses which extend beyond the use of the product under normal conditions in accord with the statements on this label.

EPA Reg. No. 471-14

Manufactured for
HENRY FIELD SEED & NURSERY CO.
SHENANDOAH, IOWA 51602 / Since 1892


(LABEL COPY)

(front panel)

HENRY FIELD'S

LAWN
WEED KILLER

2,4-D (Granular)



ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid, Isooctyl ester 2.41%

INERT INGREDIENTS: 97.59%

2,4-Dichlorophenoxyacetic Acid equivalent 1.6% by weight

Net Contents 8 oz.

(18 pt. type) CAUTION
(12 pt. ty) Keep Out Of The Reach Of Children
See back panel for additional cautions

(Back panel)

DIRECTIONS

GRANULAR LAWN WEED KILLER is a convenient form of 2,4-D as low volatile ester for killing certain broad-leaved weeds in lawn and turf. Enters the plant through the leaves and works downward to roots, killing roots and all

LAWN: To kill broad-leaved weeds such as dandelions, plantain and other di-cot (broad-leaved) weeds, apply 1 pound per 425 square feet of lawn area, or 8 ounces per 200 square feet. Do not apply within three feet of shrubs, flowers, or other desirable plants. Repeat application in one month to control weeds that were skipped. If weeds are over 5 to 6 inches tall, they should be sprayed with Henry Field's Liquid Lawn Weed Killer. Do not use on St. Augustine grass, bentgrass, Dichondra, Carpetgrass or clover lawns. Do not use on newly seeded lawns.

CAUTION: Do not take internally. Avoid contact with eyes, skin and clothing. In case of skin contact, wash thoroughly with soap and warm water. Flush eyes with clean water and obtain prompt medical attention. Do not store near seeds, fertilizers, insecticides or fungicides. Avoid drift to susceptible plants as this product may injure cotton, beans, pease, tomatoes, grapes, ornamentals. Do not use in greenhouses or on crops. Do not contaminate water intended for irrigation or domestic use. Vapors from this product may injure susceptible plants in the immediate vicinity.

NOTE: SELLER makes no warranty of any kind, expressed or implied, concerning the use of this product. BUYER assumes all risk of use or handling whether in accordance with directions or not.

EPA Reg. No. 471-

Manufactured by
HENRY FIELD & COMPANY
New York, N.Y.