

POTATO DUST

DIRECTIONS FOR USE

For control of Aphids, False Cinch Bugs, Leaf Hoppers, Mealy Bugs, Early and Late Blight on potatoes. Apply when plants are 2 to 4 inches high or when disease conditions threaten or first appear. Apply at 7 to 10 day intervals or as long as these conditions are present. Apply a light even coating of dust to both the upper and lower leaf surfaces. Apply at the rate of 2 ounces per 100 square feet.

IMPORTANT

Keep foliage covered especially new growth.

NOTICE

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

CAUTION

Harmful if swallowed. May cause irritation of nose and throat. Avoid breathing dust. Avoid contact with skin or clothing. Wash thoroughly after using. Avoid contamination of feed or food stuffs.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste. This product is highly toxic to bees exposed to direct treatment or residue on crops.

For protective information consult State Agricultural Experiment Station or State Extension Service Specialist.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTE TO PHYSICIAN: Malathion upon repeated or prolonged exposure may cause cholinesterase inhibition. Atropine and 2-PAM are antidotal.

USDA Reg. No. 471-75



ACTIVE INGREDIENTS

Malathion	1.00%
Captan (N-Trichloromethylthio-1-hydroxy-2,2-dicarboximide)	1.50%

INERT INGREDIENTS

88.50%

TOTAL

100.00%

10,0-dimethyl phosphorodithioate of diethyl malathionate

CAUTION

Henry Field's is located in the lovely, fertile
Nishna Valley of southwest Iowa where some of the
world's finest seeds and nursery stock are grown.