

FRONT LABEL

BACK LABEL



**DISINFECTANT
TOILET BOWL CLEANER**

ACTIVE INGREDIENTS:
 Hydrogen Chloride 7.00%
 Oxalic Acid 2.00%
 Alkyl (58% C18, 40% C16, 1% C14, 1% C12) Trimethylammonium Chloride 0.25%
INERT INGREDIENTS* 90.75%

* Consisting of Water, Surfactant, 1,3-diethylthiourea, perfume, dye.



DANGER KEEP OUT OF REACH OF CHILDREN

Read antidotes and warning statements on back.

NET 22 FL. OZ. (1 PT. 6 OZ.) SL-2F

ACCEPTED
 02 20 15
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTER-
 ED UNDER NO. 475-69
 TO ACCEPTED COMMENTS.

475-69



LIQUID DISINFECTANT
Sani-FLUSH

CLEANS • DEODORIZES • DISINFECTS
 Fast acting • Removes hardwater rust stains • Just pour, let stand, swab and flush • Will not harm septic tank or hinder bacterial action • Will not injure white or colored bowls.

TOILET BOWLS—Open Spout... Apply one half cupful around sides. Apply small amount to brush and clean around rims. Brush out bowl and flush. For badly stained bowls, repeat and let solution stand for 10 to 15 minutes before brushing.

TRAP OR OUTLET—Put one half cupful in bowl and allow to stand several hours or overnight.

FLUSH BOXES—Shut off water pressure and flush to empty tank. Apply Sani-Flush to brush or swab to clean tank.

Use Sani-Flush weekly to keep toilets stain-free and sanitary. For toilet bowl use only, won't harm septic tanks or plumbing. Safe for all porcelain toilets. Do not use on bathtubs, washstands, sinks or any other enameled or metal fixtures. Avoid contact with fabrics. Do not reuse container—Destroy when empty.

DANGER: CONTAINS HYDROGEN CHLORIDE and OXALIC ACID—CORROSIVE, MAY BE FATAL IF SWALLOWED. Do not get in eyes, on skin or on clothing. May produce chemical burns. Keep out of reach of children. Do not use with chlorine type bleaches, as harmful fumes may result.

ANTIDOTE—EXTERNAL—Flood the surface with water, using soap freely, then cover with moist Magnesia or Baking Soda. If splashed in eyes, wash thoroughly with water. CALL PHYSICIAN.

ANTIDOTE—INTERNAL—Give milk of magnesia, chalk or whiting suspended in much water, milk or raw egg white. CALL PHYSICIAN.

CLOSE SPOUT PROMPTLY AFTER USING

BOYLE-MIDWAY INC. NEW YORK, N. Y. 10017

E.P.A. Registration No. 475-69

©Boyle-Midway Div. 1967, 70

Made in U.S.A.

Net .648 Litre

SFLBC-17