



475-129

11-15-1971

Best Available copy

SNAROL


PELLETS

PUNCH
HERE

NET WT. 1 LB.

SNAROL

PELLETS



KILLS



SNAILS & SLUGS

WARNING: KEEP OUT OF REACH OF CHILDREN
Read warning and directions on side panel before using.

ACTIVE INGREDIENT: Metaldehyde 3.3%
INERT INGREDIENTS 96.7%

SNAROL

PELLETS



SNAROL

PELLETS

KILLS

SNAILS & SLUGS

SPREAD
BEFORE
OR AFTER SPRINKLING

One pound of SNAROL PELLETS is sufficient to treat about 1200 sq. ft. of surface area. SNAROL is also available in MEAL form.

WARNING

May be fatal if swallowed. If swallowed, induce vomiting. Call physician immediately. Harmful dust. Avoid contact with skin, eyes and clothing. Wash hands with soap and water after using. Keep out of reach of children. Keep domestic animals and poultry off treated area. Bait is attractive to dogs and chickens. Do not apply directly on fruits, vegetables or crops, but apply on ground under plants. Dispose of empty container or any spillage promptly—burn or bury. Store in a cool, dry place.

BOYLE-MIDWAY INC.

CRANFORD, N. J. • LOS ANGELES, CALIF.
CHICAGO, ILL. • CHAMBLEE, GA.

U.S.D.A. REGISTRATION NO. 475-129

© 211,521 of Related Co.
BOYLE-MIDWAY SP. Made in U.S.A.

ACCEPTED

NOV 15 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 475-129 SUBJECT
TO ATTACHED COMMENTS.

SNAROL

PELLETS

KILLS

SNAILS & SLUGS

SNAROL PELLETS are a specially prepared bait for the control of snails and slugs. The Metaldehyde in SNAROL PELLETS attracts and kills snails and slugs. These pests are night feeders and inhabit damp, cool places such as flower and vegetable gardens, around fences, hedges and bushes. The presence of snails and slugs can be determined by their shiny mucous-like trails.

SPREAD DIRECTIONS

SNAROL PELLETS late in the day, before or after thoroughly wetting the areas likely to be infested. Care should be taken, when using SNAROL PELLETS, to apply them to the soil and not directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes, apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 5 days of harvesting.

Continuous watering and high temperatures cause SNAROL PELLETS gradually to lose some of their effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control new broods emerging during this period. Three baitings, ten days apart, usually gives control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.

SPREADING