

475-129

12-4-1973

1/3

# SNAROL PELLETS



PUNCH  
HERE

## SNAROL PELLETS

# SNAROL® PELLETS

### KILLS SNAILS & SLUGS

SPREAD BEFORE OR  
AFTER SPRINKLING



SNAROL PELLETS are a specially prepared bait for the control of snails and slugs. The Metaldehyde in SNAROL PELLETS attracts and kills snails and slugs. These pests are night feeders and inhabit damp, cool places such as flower and vegetable gardens, around fences, hedges and bushes. The presence of snails and slugs can be determined by their shiny mucous-like trails.

#### DIRECTIONS

Spread SNAROL PELLETS late in the day, before or after thoroughly wetting the areas likely to be infested. Care should be taken when spreading SNAROL PELLETS, to apply them to the soil and not directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes; apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous watering and high temperatures cause SNAROL PELLETS gradually to lose some of their effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control new broods emerging during this period. Three baitings, ten days apart, usually gives control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.

## KILLS SNAILS & SLUGS


**WARNING** KEEP OUT OF REACH OF CHILDREN.  
KEEP PELLETS AWAY FROM CHILDREN, DOGS &  
OTHER PETS. MAY BE FATAL IF SWALLOWED.


ACTIVE INGREDIENT: Metaldehyde ..... 3.3%  
INERT INGREDIENTS ..... 96.7%

Read and follow all directions.  
Read warning statements on side panel.

**NET WT. 2-1/2 LBS.**

# SNAROL PELLETS





# SNAROL PELLETS

## KILLS SNAILS & SLUGS

One pound of SNAROL PELLETS is sufficient to treat about 1200 sq. ft. of surface area. SNAROL is also available in MEAL form.

### WARNING

May be fatal if swallowed. If swallowed, induce vomiting. Call physician immediately. Harmful dust. Avoid contact with skin, eyes and clothing. Wash hands with soap and water after using.

Keep out of reach of children.

Keep dogs and other pets, domestic animals or poultry away from treated areas.

Do not apply directly on fruits, vegetables or crops, but apply on ground under plants. Dispose of empty container or any spillage promptly—burn or bury. Store in a cool, dry place.

This product is toxic to wildlife and birds. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**BOYLE-MIDWAY INC.**

NEW YORK, N.Y. 10017

© Boyle-Midway Div.

EPA Registration No. 475-129-AA

Made in U.S.A. SP-7

3/3

475-129

**ACCEPTED**  
DEC 4 1973  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO.....SUBJECT  
TO ATTACHED COMMENTS.